

**Chlamydia, Gonorrhea, and Primary and Secondary Syphilis Cases by Year of Diagnosis and Sex
Connecticut, 2007 - 2014**

Year	Disease											
	Chlamydia				Gonorrhea				Syphilis			
	Sex		Sex		Sex		Sex		Sex		Sex	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
No.	Rate*	No.	Rate*	No.	Rate*	No.	Rate*	No.	Rate*	No.	Rate*	
2007	2,882	168	8,628	476	956	56	1,376	76	37	2	2	0
2008	3,246	188	9,202	505	1,108	64	1,685	93	33	2	0	0
2009	3,190	184	8,954	490	1,066	62	1,492	82	64	4	1	0
2010	3,436	197	9,256	504	1,120	64	1,448	79	94	5	5	0
2011	3,829	219	9,833	534	1,074	61	1,376	75	61	3	6	0
2012	3,666	209	9,793	531	1,010	58	1,185	64	68	4	13	1
2013	3,545	202	9,367	508	1,479	84	1,456	79	47	3	6	0
2014	3,717	212	9,341	507	1,167	67	1,052	57	80	5	7	0

Source: Connecticut STD Control Program, STD*MIS Data

Source of population data: CDC Bridged-Race Population Estimates, United States July 1st resident population by state, county, age, sex, bridged-race, and Hispanic origin. Compiled from revised bridged-race 2000-2009 intercensal population estimates (released by NCHS on 7/26/2012); and bridged-race Vintage 2014 (2010-2014) postcensal population estimates (released by NCHS on 10/26/2012); and bridged-race vintage (2010-2014) postcensal population estimates (released by NCHS on 6/30/2015). Available on CDC WONDER Online Database. Accessed at <http://wonder.cdc.gov/bridged-race-v2014.html> on Aug 17, 2015

*Rates = per 100,000 population. Rates based on counts <20 may be unstable and should be interpreted with caution.