

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.88	1	1.0	0.99	2	2.1	1.48
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.5	0.52	1	1.0	1.01	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	4	1.6	0.79	3	2.4	1.36	1	0.8	0.79
30-34	3	1.1	0.66	1	0.8	0.76	2	1.5	1.07
35-39	2	0.9	0.60	0	0.0	0.00	2	1.7	1.20
40-44	7	3.2	1.21	5	4.7	2.09	2	1.8	1.27
45-49	4	2.3	1.13	3	3.4	1.99	1	1.1	1.11
50-54	9	6.3	2.11	5	7.2	3.22	4	5.5	2.75
55-59	7	5.1	1.91	4	6.0	2.99	3	4.2	2.43
60-64	12	8.4	2.42	8	11.8	4.19	4	5.3	2.65
65-69	22	16.6	3.54	14	23.5	6.28	8	11.0	3.87
70-74	29	26.4	4.90	16	34.5	8.62	13	20.5	5.68
75-79	58	70.6	9.27	22	69.3	14.78	36	71.4	11.90
80-84	63	121.3	15.29	26	150.2	29.45	37	106.9	17.58
85+	99	226.7	22.79	26	231.8	45.46	73	225.0	26.33
Total deaths	324			136			188		
Connecticut									
. Crude rate		11.0	-		9.5	-		12.4	-
. AAMR (1940 Std. mil.)		4.9	0.32		5.8	0.53		4.1	0.38
. AAMR (1970 Std. mil.)		7.4	0.42		8.6	0.74		6.5	0.50
. AAMR (2000 Std. mil.)		10.8	0.60		12.3	1.10		9.8	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.0	-		4.6	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.72	0	0.0	0.00	2	2.1	1.48
5-9	1	0.6	0.56	0	0.0	0.00	1	1.2	1.16
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	4	1.8	0.89	2	1.7	1.23	2	1.8	1.27
25-29	19	7.5	1.72	16	12.6	3.14	3	2.4	1.38
30-34	34	12.9	2.22	27	20.6	3.96	7	5.3	2.01
35-39	22	9.4	2.00	20	17.2	3.84	2	1.7	1.20
40-44	26	11.9	2.33	22	20.5	4.38	4	3.6	1.79
45-49	15	8.4	2.18	15	17.2	4.44	0	0.0	0.00
50-54	14	9.9	2.63	11	15.8	4.78	3	4.1	2.39
55-59	3	2.2	1.25	3	4.5	2.59	0	0.0	0.00
60-64	3	2.1	1.21	3	4.4	2.56	0	0.0	0.00
65-69	2	1.5	1.07	1	1.7	1.68	1	1.4	1.37
70-74	1	0.9	0.91	1	2.2	2.16	0	0.0	0.00
75-79	1	1.2	1.22	0	0.0	0.00	1	2.0	1.98
80-84	1	1.9	1.93	1	5.8	5.78	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	148			122			26		
Connecticut									
. Crude rate		5.0	-		8.5	-		1.7	-
. AAMR (1940 Std. mil.)		4.7	0.39		7.9	0.72		1.7	0.34
. AAMR (1970 Std. mil.)		4.1	0.34		6.8	0.63		1.5	0.31
. AAMR (2000 Std. mil.)		4.9	0.41		8.3	0.77		1.7	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.7	-		15.7	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.72	0	0.0	0.00	2	2.1	1.48
5-9	5	2.8	1.26	3	3.3	1.91	2	2.3	1.64
10-14	2	1.2	0.85	1	1.2	1.18	1	1.2	1.23
15-19	3	1.5	0.89	3	3.0	1.74	0	0.0	0.00
20-24	10	4.4	1.40	5	4.4	1.95	5	4.5	2.01
25-29	22	8.7	1.85	11	8.6	2.60	11	8.7	2.64
30-34	37	14.1	2.31	19	14.5	3.32	18	13.6	3.22
35-39	70	29.9	3.57	34	29.2	5.01	36	30.5	5.08
40-44	118	53.9	4.96	53	49.4	6.79	65	58.2	7.22
45-49	185	104.2	7.66	94	107.8	11.12	91	100.7	10.56
50-54	246	173.2	11.04	124	178.6	16.04	122	168.0	15.21
55-59	486	351.5	15.95	246	367.8	23.45	240	336.3	21.71
60-64	754	526.9	19.19	408	603.7	29.89	346	458.2	24.63
65-69	984	742.0	23.65	534	896.1	38.78	450	616.3	29.05
70-74	1,130	1029.0	30.61	600	1293.6	52.81	530	835.5	36.29
75-79	997	1213.3	38.42	503	1584.8	70.66	494	979.5	44.07
80-84	806	1552.5	54.68	388	2241.2	113.78	418	1208.0	59.08
85+	765	1752.1	63.35	308	2745.8	156.46	457	1408.5	65.89
Total deaths	6,622			3,334			3,288		
Connecticut									
. Crude rate		224.5	-		233.1	-		216.3	-
. AAMR (1940 Std. mil.)		128.4	1.73		152.8	2.76		111.9	2.24
. AAMR (1970 Std. mil.)		168.9	2.12		207.8	3.62		144.6	2.65
. AAMR (2000 Std. mil.)		210.8	2.62		266.2	4.81		178.8	3.17
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.3	-		165.4	-		112.2	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	1.3	0.74	2	1.7	1.22	1	0.8	0.85
40-44	2	0.9	0.65	1	0.9	0.93	1	0.9	0.90
45-49	5	2.8	1.26	5	5.7	2.56	0	0.0	0.00
50-54	14	9.9	2.63	10	14.4	4.55	4	5.5	2.75
55-59	14	10.1	2.71	9	13.5	4.49	5	7.0	3.13
60-64	18	12.6	2.96	13	19.2	5.33	5	6.6	2.96
65-69	16	12.1	3.02	13	21.8	6.05	3	4.1	2.37
70-74	14	12.7	3.41	9	19.4	6.47	5	7.9	3.53
75-79	14	17.0	4.55	10	31.5	9.96	4	7.9	3.97
80-84	7	13.5	5.10	4	23.1	11.55	3	8.7	5.01
85+	5	11.5	5.12	2	17.8	12.61	3	9.2	5.34
Total deaths	112			78			34		
Connecticut									
. Crude rate		3.8	-		5.5	-		2.2	-
. AAMR (1940 Std. mil.)		2.7	0.27		4.2	0.49		1.4	0.27
. AAMR (1970 Std. mil.)		3.2	0.31		5.0	0.57		1.7	0.30
. AAMR (2000 Std. mil.)		3.7	0.35		5.8	0.68		2.0	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.7	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	4	1.7	0.85	3	2.6	1.49	1	0.8	0.85
40-44	14	6.4	1.71	6	5.6	2.28	8	7.2	2.53
45-49	17	9.6	2.32	6	6.9	2.81	11	12.2	3.67
50-54	15	10.6	2.73	9	13.0	4.32	6	8.3	3.37
55-59	35	25.3	4.28	18	26.9	6.34	17	23.8	5.78
60-64	82	57.3	6.33	46	68.1	10.04	36	47.7	7.95
65-69	100	75.4	7.54	56	94.0	12.56	44	60.3	9.08
70-74	160	145.7	11.52	94	202.7	20.90	66	104.0	12.81
75-79	146	177.7	14.70	74	233.2	27.10	72	142.8	16.82
80-84	116	223.4	20.75	49	283.0	40.43	67	193.6	23.65
85+	170	389.4	29.86	56	499.2	66.71	114	351.4	32.91
Total deaths	862			419			443		
Connecticut									
. Crude rate		29.2	-		29.3	-		29.1	-
. AAMR (1940 Std. mil.)		14.4	0.55		17.8	0.91		12.1	0.68
. AAMR (1970 Std. mil.)		20.6	0.72		25.8	1.27		17.2	0.87
. AAMR (2000 Std. mil.)		27.6	0.95		34.8	1.78		23.2	1.12
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.0	-		17.0	-		11.7	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	3	1.3	0.74	2	1.7	1.22	1	0.8	0.85
40-44	13	5.9	1.65	6	5.6	2.28	7	6.3	2.37
45-49	15	8.4	2.18	5	5.7	2.56	10	11.1	3.50
50-54	14	9.9	2.63	8	11.5	4.07	6	8.3	3.37
55-59	30	21.7	3.96	16	23.9	5.98	14	19.6	5.24
60-64	69	48.2	5.80	40	59.2	9.36	29	38.4	7.13
65-69	90	67.9	7.15	49	82.2	11.75	41	56.1	8.77
70-74	130	118.4	10.38	74	159.5	18.55	56	88.3	11.80
75-79	126	153.3	13.66	63	198.5	25.01	63	124.9	15.74
80-84	99	190.7	19.17	42	242.6	37.43	57	164.7	21.82
85+	143	327.5	27.39	49	436.8	62.41	94	289.7	29.88
Total deaths	735			356			379		
Connecticut									
. Crude rate		24.9	-		24.9	-		24.9	-
. AAMR (1940 Std. mil.)		12.4	0.51		15.2	0.84		10.5	0.64
. AAMR (1970 Std. mil.)		17.6	0.66		22.0	1.17		14.8	0.80
. AAMR (2000 Std. mil.)		23.5	0.88		29.6	1.65		19.9	1.04
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.7	-		14.2	-		10.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.43	1	0.9	0.86	0	0.0	0.00
40-44	1	0.5	0.46	0	0.0	0.00	1	0.9	0.90
45-49	2	1.1	0.80	1	1.1	1.15	1	1.1	1.11
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	5	3.6	1.62	2	3.0	2.11	3	4.2	2.43
60-64	12	8.4	2.42	6	8.9	3.62	6	7.9	3.24
65-69	10	7.5	2.38	7	11.7	4.44	3	4.1	2.37
70-74	24	21.9	4.46	16	34.5	8.62	8	12.6	4.46
75-79	18	21.9	5.16	9	28.4	9.45	9	17.8	5.95
80-84	15	28.9	7.46	7	40.4	15.28	8	23.1	8.17
85+	23	52.7	10.98	7	62.4	23.59	16	49.3	12.33
Total deaths	112			57			55		
Connecticut									
. Crude rate		3.8	-		4.0	-		3.6	-
. AAMR (1940 Std. mil.)		1.8	0.19		2.4	0.33		1.4	0.23
. AAMR (1970 Std. mil.)		2.7	0.26		3.5	0.47		2.1	0.30
. AAMR (2000 Std. mil.)		3.6	0.34		4.7	0.65		2.9	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.5	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.9	0.60	0	0.0	0.00	2	1.7	1.20
40-44	3	1.4	0.79	3	2.8	1.62	0	0.0	0.00
45-49	9	5.1	1.69	6	6.9	2.81	3	3.3	1.92
50-54	9	6.3	2.11	6	8.6	3.53	3	4.1	2.39
55-59	15	10.8	2.80	11	16.4	4.96	4	5.6	2.80
60-64	37	25.9	4.25	21	31.1	6.78	16	21.2	5.30
65-69	60	45.2	5.84	34	57.1	9.78	26	35.6	6.98
70-74	65	59.2	7.34	32	69.0	12.20	33	52.0	9.06
75-79	55	66.9	9.02	16	50.4	12.60	39	77.3	12.38
80-84	37	71.3	11.72	23	132.9	27.70	14	40.5	10.81
85+	44	100.8	15.19	11	98.1	29.57	33	101.7	17.71
Total deaths	336			163			173		
Connecticut									
. Crude rate		11.4	-		11.4	-		11.4	-
. AAMR (1940 Std. mil.)		6.1	0.37		7.6	0.61		5.0	0.43
. AAMR (1970 Std. mil.)		8.4	0.47		10.2	0.80		7.0	0.55
. AAMR (2000 Std. mil.)		10.5	0.58		12.7	1.03		9.0	0.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		7.5	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	0	0.0	0.00	1	0.9	0.90
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	8	3.4	1.21	7	6.0	2.27	1	0.8	0.85
40-44	18	8.2	1.94	13	12.1	3.36	5	4.5	2.00
45-49	55	31.0	4.18	36	41.3	6.88	19	21.0	4.83
50-54	69	48.6	5.85	45	64.8	9.66	24	33.0	6.75
55-59	164	118.6	9.26	94	140.5	14.50	70	98.1	11.72
60-64	239	167.0	10.80	151	223.4	18.18	88	116.5	12.42
65-69	313	236.0	13.34	182	305.4	22.64	131	179.4	15.67
70-74	330	300.5	16.54	194	418.3	30.03	136	214.4	18.38
75-79	246	299.4	19.09	151	475.8	38.72	95	188.4	19.33
80-84	140	269.7	22.79	90	519.9	54.80	50	144.5	20.43
85+	90	206.1	21.73	51	454.7	63.67	39	120.2	19.25
Total deaths	1,675			1,016			659		
Connecticut									
. Crude rate		56.8	-		71.0	-		43.4	-
. AAMR (1940 Std. mil.)		35.0	0.92		48.5	1.58		24.8	1.06
. AAMR (1970 Std. mil.)		44.5	1.11		63.4	2.00		31.2	1.26
. AAMR (2000 Std. mil.)		52.3	1.29		77.0	2.50		35.8	1.42
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.3	-		57.9	-		25.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	6	2.6	1.05	3	2.6	1.49	3	2.5	1.47
40-44	6	2.7	1.12	4	3.7	1.87	2	1.8	1.27
45-49	1	0.6	0.56	1	1.1	1.15	0	0.0	0.00
50-54	5	3.5	1.57	2	2.9	2.04	3	4.1	2.39
55-59	9	6.5	2.17	3	4.5	2.59	6	8.4	3.43
60-64	8	5.6	1.98	5	7.4	3.31	3	4.0	2.29
65-69	12	9.0	2.61	6	10.1	4.11	6	8.2	3.35
70-74	12	10.9	3.15	8	17.2	6.10	4	6.3	3.15
75-79	8	9.7	3.44	4	12.6	6.30	4	7.9	3.97
80-84	4	7.7	3.85	1	5.8	5.78	3	8.7	5.01
85+	9	20.6	6.87	2	17.8	12.61	7	21.6	8.15
Total deaths	84			42			42		
Connecticut									
. Crude rate		2.8	-		2.9	-		2.8	-
. AAMR (1940 Std. mil.)		1.9	0.23		2.2	0.35		1.7	0.29
. AAMR (1970 Std. mil.)		2.2	0.25		2.5	0.39		2.0	0.32
. AAMR (2000 Std. mil.)		2.7	0.30		3.1	0.49		2.4	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.8	-		2.6	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	-	-	-	1	1.0	1.05
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	2	0.8	0.56	-	-	-	2	1.6	1.12
30-34	8	3.0	1.08	-	-	-	8	6.1	2.14
35-39	15	6.4	1.65	-	-	-	15	12.7	3.28
40-44	27	12.3	2.37	-	-	-	27	24.2	4.65
45-49	22	12.4	2.64	-	-	-	22	24.4	5.19
50-54	36	25.3	4.22	-	-	-	36	49.6	8.26
55-59	52	37.6	5.22	-	-	-	52	72.9	10.10
60-64	83	58.0	6.37	-	-	-	83	109.9	12.06
65-69	73	55.0	6.44	-	-	-	73	100.0	11.70
70-74	100	91.1	9.11	-	-	-	100	157.6	15.76
75-79	73	88.8	10.40	-	-	-	73	144.7	16.94
80-84	61	117.5	15.04	-	-	-	61	176.3	22.57
85+	78	178.6	20.23	-	-	-	78	240.4	27.22
Total deaths	631			-			631		
Connecticut									
. Crude rate		-	-		-	-		41.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		24.0	1.08
. AAMR (1970 Std. mil.)		-	-		-	-		29.2	1.22
. AAMR (2000 Std. mil.)		-	-		-	-		35.5	1.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		23.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	3	1.3	0.74	-	-	-	3	2.5	1.47
40-44	3	1.4	0.79	-	-	-	3	2.7	1.55
45-49	3	1.7	0.98	-	-	-	3	3.3	1.92
50-54	2	1.4	1.00	-	-	-	2	2.8	1.95
55-59	2	1.4	1.02	-	-	-	2	2.8	1.98
60-64	4	2.8	1.40	-	-	-	4	5.3	2.65
65-69	7	5.3	2.00	-	-	-	7	9.6	3.62
70-74	4	3.6	1.82	-	-	-	4	6.3	3.15
75-79	9	11.0	3.65	-	-	-	9	17.8	5.95
80-84	2	3.9	2.72	-	-	-	2	5.8	4.09
85+	4	9.2	4.58	-	-	-	4	12.3	6.16
Total deaths	43			-			43		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.7	0.29
. AAMR (1970 Std. mil.)		-	-		-	-		2.0	0.32
. AAMR (2000 Std. mil.)		-	-		-	-		2.5	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	0	0.0	0.00	-	-	-	0	0.0	0.00
50-54	0	0.0	0.00	-	-	-	0	0.0	0.00
55-59	3	2.2	1.25	-	-	-	3	4.2	2.43
60-64	5	3.5	1.56	-	-	-	5	6.6	2.96
65-69	10	7.5	2.38	-	-	-	10	13.7	4.33
70-74	8	7.3	2.58	-	-	-	8	12.6	4.46
75-79	5	6.1	2.72	-	-	-	5	9.9	4.43
80-84	5	9.6	4.31	-	-	-	5	14.4	6.46
85+	8	18.3	6.48	-	-	-	8	24.7	8.72
Total deaths	44			-			44		
Connecticut									
. Crude rate		-	-		-	-		2.9	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	0.22
. AAMR (1970 Std. mil.)		-	-		-	-		1.8	0.29
. AAMR (2000 Std. mil.)		-	-		-	-		2.2	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.4	0.39	-	-	-	1	0.8	0.79
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	4	1.8	0.91	-	-	-	4	3.6	1.79
45-49	11	6.2	1.87	-	-	-	11	12.2	3.67
50-54	12	8.4	2.44	-	-	-	12	16.5	4.77
55-59	17	12.3	2.98	-	-	-	17	23.8	5.78
60-64	20	14.0	3.13	-	-	-	20	26.5	5.92
65-69	26	19.6	3.85	-	-	-	26	35.6	6.98
70-74	26	23.7	4.64	-	-	-	26	41.0	8.04
75-79	21	25.6	5.58	-	-	-	21	41.6	9.09
80-84	16	30.8	7.70	-	-	-	16	46.2	11.56
85+	15	34.4	8.87	-	-	-	15	46.2	11.94
Total deaths	169			-			169		
Connecticut									
. Crude rate		-	-		-	-		11.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.7	0.58
. AAMR (1970 Std. mil.)		-	-		-	-		8.2	0.66
. AAMR (2000 Std. mil.)		-	-		-	-		9.6	0.76
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.9	0.60	2	1.7	1.22	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.6	0.56	0	0.0	0.00	1	1.1	1.11
50-54	3	2.1	1.22	3	4.3	2.49	0	0.0	0.00
55-59	7	5.1	1.91	5	7.5	3.34	2	2.8	1.98
60-64	11	7.7	2.32	10	14.8	4.68	1	1.3	1.32
65-69	13	9.8	2.72	9	15.1	5.03	4	5.5	2.74
70-74	26	23.7	4.64	21	45.3	9.88	5	7.9	3.53
75-79	27	32.9	6.32	21	66.2	14.44	6	11.9	4.86
80-84	33	63.6	11.07	24	138.6	28.30	9	26.0	8.67
85+	37	84.7	13.93	26	231.8	45.46	11	33.9	10.22
Total deaths	160			121			39		
Connecticut									
. Crude rate		5.4	-		8.5	-		2.6	-
. AAMR (1940 Std. mil.)		2.4	0.22		4.8	0.45		0.9	0.18
. AAMR (1970 Std. mil.)		3.7	0.30		7.6	0.69		1.4	0.24
. AAMR (2000 Std. mil.)		5.2	0.41		11.2	1.06		2.0	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.8	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	2	1.1	0.80	1	1.1	1.10	1	1.2	1.16
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	1	0.5	0.52	1	1.0	1.01	0	0.0	0.00
20-24	3	1.3	0.77	1	0.9	0.87	2	1.8	1.27
25-29	5	2.0	0.88	4	3.1	1.57	1	0.8	0.79
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	2	0.9	0.60	1	0.9	0.86	1	0.8	0.85
40-44	3	1.4	0.79	2	1.9	1.32	1	0.9	0.90
45-49	5	2.8	1.26	4	4.6	2.29	1	1.1	1.11
50-54	6	4.2	1.72	6	8.6	3.53	0	0.0	0.00
55-59	16	11.6	2.89	8	12.0	4.23	8	11.2	3.96
60-64	24	16.8	3.42	14	20.7	5.54	10	13.2	4.19
65-69	20	15.1	3.37	10	16.8	5.31	10	13.7	4.33
70-74	22	20.0	4.27	11	23.7	7.15	11	17.3	5.23
75-79	18	21.9	5.16	5	15.8	7.05	13	25.8	7.15
80-84	11	21.2	6.39	5	28.9	12.92	6	17.3	7.08
85+	5	11.5	5.12	2	17.8	12.61	3	9.2	5.34
Total deaths	146			77			69		
Connecticut									
. Crude rate		4.9	-		5.4	-		4.5	-
. AAMR (1940 Std. mil.)		3.4	0.31		4.3	0.51		2.7	0.36
. AAMR (1970 Std. mil.)		4.1	0.35		5.0	0.58		3.3	0.42
. AAMR (2000 Std. mil.)		4.6	0.39		5.6	0.65		3.8	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.4	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.6	0.56	1	1.1	1.10	0	0.0	0.00
10-14	1	0.6	0.60	0	0.0	0.00	1	1.2	1.23
15-19	2	1.0	0.73	2	2.0	1.42	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	2	0.8	0.56	1	0.8	0.78	1	0.8	0.79
30-34	6	2.3	0.93	4	3.1	1.53	2	1.5	1.07
35-39	8	3.4	1.21	3	2.6	1.49	5	4.2	1.90
40-44	6	2.7	1.12	3	2.8	1.62	3	2.7	1.55
45-49	8	4.5	1.59	5	5.7	2.56	3	3.3	1.92
50-54	6	4.2	1.72	3	4.3	2.49	3	4.1	2.39
55-59	16	11.6	2.89	10	15.0	4.73	6	8.4	3.43
60-64	14	9.8	2.61	9	13.3	4.44	5	6.6	2.96
65-69	30	22.6	4.13	15	25.2	6.50	15	20.5	5.30
70-74	28	25.5	4.82	14	30.2	8.07	14	22.1	5.90
75-79	42	51.1	7.89	18	56.7	13.37	24	47.6	9.71
80-84	36	69.3	11.56	18	104.0	24.51	18	52.0	12.26
85+	27	61.8	11.90	10	89.2	28.19	17	52.4	12.71
Total deaths	234			117			117		
Connecticut									
. Crude rate		7.9	-		8.2	-		7.7	-
. AAMR (1940 Std. mil.)		4.6	0.34		5.6	0.55		3.8	0.42
. AAMR (1970 Std. mil.)		5.9	0.40		7.3	0.69		4.9	0.49
. AAMR (2000 Std. mil.)		7.6	0.50		9.5	0.91		6.4	0.60
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.1	-		6.5	-		3.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	0	0.0	0.00	1	0.9	0.90
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	4	1.5	0.76	3	2.3	1.32	1	0.8	0.76
35-39	4	1.7	0.85	4	3.4	1.72	0	0.0	0.00
40-44	8	3.7	1.29	6	5.6	2.28	2	1.8	1.27
45-49	11	6.2	1.87	5	5.7	2.56	6	6.6	2.71
50-54	12	8.4	2.44	8	11.5	4.07	4	5.5	2.75
55-59	12	8.7	2.51	8	12.0	4.23	4	5.6	2.80
60-64	43	30.0	4.58	26	38.5	7.54	17	22.5	5.46
65-69	56	42.2	5.64	32	53.7	9.49	24	32.9	6.71
70-74	63	57.4	7.23	28	60.4	11.41	35	55.2	9.33
75-79	72	87.6	10.33	37	116.6	19.16	35	69.4	11.73
80-84	69	132.9	16.00	24	138.6	28.30	45	130.0	19.39
85+	73	167.2	19.57	15	133.7	34.53	58	178.8	23.47
Total deaths	429			197			232		
Connecticut									
. Crude rate		14.5	-		13.8	-		15.3	-
. AAMR (1940 Std. mil.)		7.5	0.40		9.0	0.67		6.1	0.48
. AAMR (1970 Std. mil.)		10.4	0.51		12.1	0.87		8.9	0.61
. AAMR (2000 Std. mil.)		13.8	0.67		15.6	1.16		12.1	0.80
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.6	-		12.3	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	0	0.0	0.00	1	1.0	1.05
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.5	0.52	0	0.0	0.00	1	1.1	1.06
20-24	1	0.4	0.44	0	0.0	0.00	1	0.9	0.90
25-29	3	1.2	0.68	2	1.6	1.11	1	0.8	0.79
30-34	6	2.3	0.93	5	3.8	1.71	1	0.8	0.76
35-39	12	5.1	1.48	8	6.9	2.43	4	3.4	1.69
40-44	17	7.8	1.88	12	11.2	3.23	5	4.5	2.00
45-49	28	15.8	2.98	17	19.5	4.73	11	12.2	3.67
50-54	37	26.0	4.28	25	36.0	7.20	12	16.5	4.77
55-59	88	63.7	6.79	54	80.7	10.99	34	47.6	8.17
60-64	195	136.3	9.76	118	174.6	16.07	77	102.0	11.62
65-69	268	202.1	12.34	137	229.9	19.64	131	179.4	15.67
70-74	331	301.4	16.57	171	368.7	28.19	160	252.2	19.94
75-79	372	452.7	23.47	193	608.1	43.77	179	354.9	26.53
80-84	325	626.0	34.72	130	750.9	65.86	195	563.5	40.35
85+	412	943.6	46.49	120	1069.8	97.66	292	900.0	52.67
Total deaths	2,097			992			1,105		
Connecticut									
. Crude rate		71.1	-		69.4	-		72.7	-
. AAMR (1940 Std. mil.)		34.6	0.84		42.9	1.42		28.3	1.01
. AAMR (1970 Std. mil.)		50.0	1.11		61.5	1.96		41.7	1.32
. AAMR (2000 Std. mil.)		67.1	1.48		82.2	2.73		56.9	1.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	3	2.2	1.25	2	3.0	2.11	1	1.4	1.40
60-64	3	2.1	1.21	0	0.0	0.00	3	4.0	2.29
65-69	8	6.0	2.13	4	6.7	3.36	4	5.5	2.74
70-74	13	11.8	3.28	8	17.2	6.10	5	7.9	3.53
75-79	37	45.0	7.40	17	53.6	12.99	20	39.7	8.87
80-84	41	79.0	12.33	16	92.4	23.11	25	72.2	14.45
85+	85	194.7	21.12	22	196.1	41.82	63	194.2	24.46
Total deaths	190			69			121		
Connecticut									
. Crude rate		6.4	-		4.8	-		8.0	-
. AAMR (1940 Std. mil.)		2.1	0.17		2.4	0.29		1.9	0.21
. AAMR (1970 Std. mil.)		3.9	0.29		4.3	0.52		3.6	0.35
. AAMR (2000 Std. mil.)		6.4	0.47		7.1	0.88		6.0	0.55
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		2.4	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	7.1	1.90	7	6.9	2.62	7	7.3	2.77
5-9	2	1.1	0.80	1	1.1	1.10	1	1.2	1.16
10-14	2	1.2	0.85	2	2.4	1.66	0	0.0	0.00
15-19	4	2.1	1.03	2	2.0	1.42	2	2.1	1.50
20-24	4	1.8	0.89	4	3.5	1.75	0	0.0	0.00
25-29	10	3.9	1.25	6	4.7	1.92	4	3.2	1.59
30-34	31	11.8	2.12	22	16.8	3.58	9	6.8	2.27
35-39	47	20.1	2.93	35	30.1	5.09	12	10.2	2.94
40-44	88	40.2	4.29	66	61.6	7.58	22	19.7	4.20
45-49	137	77.2	6.59	103	118.1	11.64	34	37.6	6.45
50-54	198	139.4	9.91	133	191.6	16.61	65	89.5	11.10
55-59	362	261.8	13.76	261	390.2	24.15	101	141.5	14.08
60-64	677	473.1	18.18	444	657.0	31.18	233	308.6	20.21
65-69	1,026	773.7	24.15	643	1079.0	42.55	383	524.5	26.80
70-74	1,436	1307.7	34.51	842	1815.4	62.56	594	936.4	38.42
75-79	1,780	2166.1	51.34	968	3049.9	98.03	812	1610.0	56.50
80-84	1,979	3811.9	85.69	856	4944.5	169.00	1,123	3245.3	96.84
85+	3,817	8742.2	141.50	1,082	9646.1	293.25	2,735	8429.7	161.19
Total deaths	11,614			5,477			6,137		
Connecticut									
. Crude rate		393.7	-		383.0	-		403.7	-
. AAMR (1940 Std. mil.)		166.7	1.78		225.2	3.20		122.5	1.95
. AAMR (1970 Std. mil.)		261.1	2.49		343.1	4.65		203.1	2.76
. AAMR (2000 Std. mil.)		382.4	3.57		489.8	6.94		310.8	3.99
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		197.2	-		260.2	-		147.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.88	0	0.0	0.00	3	3.1	1.81
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	11	4.2	1.26	9	6.9	2.29	2	1.5	1.07
35-39	19	8.1	1.86	17	14.6	3.54	2	1.7	1.20
40-44	51	23.3	3.26	46	42.9	6.33	5	4.5	2.00
45-49	80	45.1	5.04	64	73.4	9.17	16	17.7	4.43
50-54	126	88.7	7.90	97	139.7	14.19	29	39.9	7.42
55-59	244	176.5	11.30	182	272.1	20.17	62	86.9	11.03
60-64	441	308.2	14.68	304	449.8	25.80	137	181.4	15.50
65-69	649	489.4	19.21	419	703.1	34.35	230	315.0	20.77
70-74	913	831.4	27.52	564	1216.0	51.20	349	550.2	29.45
75-79	1,059	1288.7	39.60	599	1887.3	77.11	460	912.0	42.52
80-84	1,157	2228.6	65.52	510	2945.9	130.45	647	1869.7	73.51
85+	2,252	5157.8	108.69	660	5883.9	229.03	1,592	4906.8	122.98
Total deaths	7,006			3,472			3,534		
Connecticut									
. Crude rate		237.5	-		242.8	-		232.5	-
. AAMR (1940 Std. mil.)		101.6	1.39		144.1	2.57		69.2	1.44
. AAMR (1970 Std. mil.)		158.3	1.94		217.4	3.70		116.2	2.07
. AAMR (2000 Std. mil.)		230.0	2.77		307.4	5.47		178.2	3.01
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		125.6	-		174.7	-		87.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.88	3	3.0	1.71	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.5	0.52	0	0.0	0.00	1	1.1	1.06
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	1.1	0.80	2	2.3	1.62	0	0.0	0.00
50-54	2	1.4	1.00	0	0.0	0.00	2	2.8	1.95
55-59	2	1.4	1.02	2	3.0	2.11	0	0.0	0.00
60-64	10	7.0	2.21	7	10.4	3.91	3	4.0	2.29
65-69	19	14.3	3.29	11	18.5	5.57	8	11.0	3.87
70-74	29	26.4	4.90	21	45.3	9.88	8	12.6	4.46
75-79	64	77.9	9.74	36	113.4	18.90	28	55.5	10.49
80-84	61	117.5	15.04	25	144.4	28.88	36	104.0	17.34
85+	171	391.6	29.95	51	454.7	63.67	120	369.9	33.76
Total deaths	365			159			206		
Connecticut									
. Crude rate		12.4	-		11.1	-		13.6	-
. AAMR (1940 Std. mil.)		4.3	0.26		5.8	0.48		3.2	0.29
. AAMR (1970 Std. mil.)		7.6	0.41		10.0	0.80		6.1	0.45
. AAMR (2000 Std. mil.)		12.3	0.65		15.7	1.30		10.3	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.7	-		6.6	-		5.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	2.0	0.88	3	2.4	1.36	2	1.6	1.12
30-34	6	2.3	0.93	2	1.5	1.08	4	3.0	1.52
35-39	4	1.7	0.85	4	3.4	1.72	0	0.0	0.00
40-44	13	5.9	1.65	5	4.7	2.09	8	7.2	2.53
45-49	17	9.6	2.32	10	11.5	3.63	7	7.7	2.93
50-54	19	13.4	3.07	6	8.6	3.53	13	17.9	4.97
55-59	29	21.0	3.90	18	26.9	6.34	11	15.4	4.65
60-64	55	38.4	5.18	22	32.6	6.94	33	43.7	7.61
65-69	105	79.2	7.73	52	87.3	12.10	53	72.6	9.97
70-74	178	162.1	12.15	80	172.5	19.28	98	154.5	15.61
75-79	248	301.8	19.16	122	384.4	34.80	126	249.8	22.26
80-84	353	679.9	36.19	139	802.9	68.10	214	618.4	42.27
85+	608	1392.5	56.47	138	1230.3	104.73	470	1448.6	66.82
Total deaths	1,642			603			1,039		
Connecticut									
. Crude rate		55.7	-		42.2	-		68.4	-
. AAMR (1940 Std. mil.)		21.3	0.61		23.0	0.98		20.2	0.78
. AAMR (1970 Std. mil.)		35.5	0.90		37.8	1.55		33.8	1.11
. AAMR (2000 Std. mil.)		54.8	1.36		57.3	2.43		52.7	1.64
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		28.1	-		31.0	-		26.1	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	0	0.0	0.00	1	1.0	1.05
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	1.1	0.80	2	2.3	1.62	0	0.0	0.00
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.38
55-59	2	1.4	1.02	2	3.0	2.11	0	0.0	0.00
60-64	6	4.2	1.71	4	5.9	2.96	2	2.6	1.87
65-69	9	6.8	2.26	4	6.7	3.36	5	6.8	3.06
70-74	10	9.1	2.88	5	10.8	4.82	5	7.9	3.53
75-79	20	24.3	5.44	9	28.4	9.45	11	21.8	6.58
80-84	25	48.2	9.63	10	57.8	18.27	15	43.3	11.19
85+	134	306.9	26.51	33	294.2	51.21	101	311.3	30.98
Total deaths	210			69			141		
Connecticut									
. Crude rate		7.1	-		4.8	-		9.3	-
. AAMR (1940 Std. mil.)		2.1	0.18		2.4	0.30		2.0	0.22
. AAMR (1970 Std. mil.)		4.2	0.29		4.4	0.53		3.9	0.35
. AAMR (2000 Std. mil.)		7.2	0.50		7.5	0.94		6.9	0.59
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.0	-		3.4	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.5	0.52	1	1.0	1.01	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	0	0.0	0.00	1	0.8	0.79
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	3	1.4	0.79	2	1.9	1.32	1	0.9	0.90
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	5	3.6	1.62	3	4.5	2.59	2	2.8	1.98
60-64	19	13.3	3.05	15	22.2	5.73	4	5.3	2.65
65-69	20	15.1	3.37	16	26.8	6.71	4	5.5	2.74
70-74	25	22.8	4.55	20	43.1	9.64	5	7.9	3.53
75-79	44	53.5	8.07	27	85.1	16.37	17	33.7	8.17
80-84	54	104.0	14.15	37	213.7	35.14	17	49.1	11.92
85+	49	112.2	16.03	15	133.7	34.53	34	104.8	17.97
Total deaths	221			136			85		
Connecticut									
. Crude rate		7.5	-		9.5	-		5.6	-
. AAMR (1940 Std. mil.)		3.2	0.24		5.4	0.48		1.7	0.23
. AAMR (1970 Std. mil.)		5.0	0.35		8.4	0.73		2.8	0.32
. AAMR (2000 Std. mil.)		7.1	0.48		11.8	1.05		4.3	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		5.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	3.6	1.35	2	2.0	1.40	5	5.2	2.34
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.5	0.52	0	0.0	0.00	1	1.1	1.06
20-24	2	0.9	0.63	1	0.9	0.87	1	0.9	0.90
25-29	2	0.8	0.56	2	1.6	1.11	0	0.0	0.00
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	5	2.1	0.95	4	3.4	1.72	1	0.8	0.85
40-44	8	3.7	1.29	4	3.7	1.87	4	3.6	1.79
45-49	8	4.5	1.59	4	4.6	2.29	4	4.4	2.21
50-54	10	7.0	2.23	8	11.5	4.07	2	2.8	1.95
55-59	18	13.0	3.07	13	19.4	5.39	5	7.0	3.13
60-64	25	17.5	3.49	17	25.2	6.10	8	10.6	3.75
65-69	57	43.0	5.69	33	55.4	9.64	24	32.9	6.71
70-74	87	79.2	8.49	59	127.2	16.56	28	44.1	8.34
75-79	105	127.8	12.47	59	185.9	24.20	46	91.2	13.45
80-84	193	371.8	26.76	92	531.4	55.40	101	291.9	29.04
85+	500	1145.2	51.21	165	1471.0	114.52	335	1032.5	56.41
Total deaths	1,030			464			566		
Connecticut									
. Crude rate		34.9	-		32.4	-		37.2	-
. AAMR (1940 Std. mil.)		12.4	0.46		17.0	0.84		9.5	0.53
. AAMR (1970 Std. mil.)		21.6	0.69		29.4	1.37		17.1	0.77
. AAMR (2000 Std. mil.)		34.8	1.09		47.2	2.29		28.4	1.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.7	-		18.2	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
5-9	1	0.6	0.56	0	0.0	0.00	1	1.2	1.16
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	1	0.4	0.43	0	0.0	0.00	1	0.8	0.85
40-44	2	0.9	0.65	2	1.9	1.32	0	0.0	0.00
45-49	7	3.9	1.49	2	2.3	1.62	5	5.5	2.48
50-54	10	7.0	2.23	2	2.9	2.04	8	11.0	3.89
55-59	39	28.2	4.52	18	26.9	6.34	21	29.4	6.42
60-64	68	47.5	5.76	35	51.8	8.75	33	43.7	7.61
65-69	95	71.6	7.35	55	92.3	12.44	40	54.8	8.66
70-74	168	153.0	11.80	84	181.1	19.76	84	132.4	14.45
75-79	206	250.7	17.47	114	359.2	33.64	92	182.4	19.02
80-84	177	340.9	25.63	97	560.3	56.89	80	231.2	25.85
85+	158	361.9	28.79	81	722.1	80.24	77	237.3	27.05
Total deaths	935			493			442		
Connecticut									
. Crude rate		31.7	-		34.5	-		29.1	-
. AAMR (1940 Std. mil.)		14.8	0.53		19.3	0.90		12.2	0.68
. AAMR (1970 Std. mil.)		22.1	0.74		30.6	1.38		17.6	0.88
. AAMR (2000 Std. mil.)		30.0	0.99		43.8	2.06		23.2	1.11
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.5	-		26.9	-		14.8	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----COPD-related (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.72	1	1.0	0.99	1	1.0	1.05
5-9	1	0.6	0.56	0	0.0	0.00	1	1.2	1.16
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	2	0.8	0.54	2	1.5	1.08	0	0.0	0.00
35-39	4	1.7	0.85	1	0.9	0.86	3	2.5	1.47
40-44	4	1.8	0.91	3	2.8	1.62	1	0.9	0.90
45-49	13	7.3	2.03	7	8.0	3.03	6	6.6	2.71
50-54	26	18.3	3.59	13	18.7	5.19	13	17.9	4.97
55-59	75	54.2	6.26	33	49.3	8.59	42	58.9	9.08
60-64	162	113.2	8.89	88	130.2	13.88	74	98.0	11.39
65-69	262	197.6	12.21	151	253.4	20.62	111	152.0	14.43
70-74	398	362.4	18.17	238	513.1	33.26	160	252.2	19.94
75-79	438	533.0	25.47	250	787.7	49.82	188	372.7	27.19
80-84	404	778.2	38.72	229	1322.8	87.41	175	505.7	38.23
85+	443	1014.6	48.21	215	1916.7	130.72	228	702.7	46.54
Total deaths	2,236			1,233			1,003		
Connecticut									
. Crude rate		75.8	-		86.2	-		66.0	-
. AAMR (1940 Std. mil.)		34.7	0.81		48.3	1.43		26.4	0.98
. AAMR (1970 Std. mil.)		52.2	1.13		76.4	2.18		38.7	1.29
. AAMR (2000 Std. mil.)		71.5	1.52		109.4	3.26		51.9	1.65
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.8	0.56	1	0.8	0.78	1	0.8	0.79
30-34	9	3.4	1.14	8	6.1	2.16	1	0.8	0.76
35-39	12	5.1	1.48	10	8.6	2.72	2	1.7	1.20
40-44	15	6.9	1.77	8	7.5	2.64	7	6.3	2.37
45-49	19	10.7	2.45	15	17.2	4.44	4	4.4	2.21
50-54	39	27.5	4.40	24	34.6	7.06	15	20.7	5.33
55-59	33	23.9	4.16	23	34.4	7.17	10	14.0	4.43
60-64	46	32.1	4.74	32	47.3	8.37	14	18.5	4.95
65-69	48	36.2	5.22	22	36.9	7.87	26	35.6	6.98
70-74	31	28.2	5.07	16	34.5	8.62	15	23.6	6.11
75-79	29	35.3	6.55	14	44.1	11.79	15	29.7	7.68
80-84	11	21.2	6.39	7	40.4	15.28	4	11.6	5.78
85+	12	27.5	7.93	5	44.6	19.93	7	21.6	8.15
Total deaths	306			185			121		
Connecticut									
. Crude rate		10.4	-		12.9	-		8.0	-
. AAMR (1940 Std. mil.)		7.8	0.47		10.5	0.80		5.3	0.53
. AAMR (1970 Std. mil.)		8.7	0.51		11.7	0.87		6.1	0.58
. AAMR (2000 Std. mil.)		10.1	0.59		13.7	1.03		7.1	0.66
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.2	-		12.8	-		5.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	0	0.0	0.00	1	0.8	0.79
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	1	0.4	0.43	0	0.0	0.00	1	0.8	0.85
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	4	2.3	1.13	3	3.4	1.99	1	1.1	1.11
50-54	2	1.4	1.00	1	1.4	1.44	1	1.4	1.38
55-59	3	2.2	1.25	3	4.5	2.59	0	0.0	0.00
60-64	16	11.2	2.80	9	13.3	4.44	7	9.3	3.50
65-69	21	15.8	3.46	8	13.4	4.75	13	17.8	4.94
70-74	31	28.2	5.07	16	34.5	8.62	15	23.6	6.11
75-79	41	49.9	7.79	31	97.7	17.54	10	19.8	6.27
80-84	50	96.3	13.62	24	138.6	28.30	26	75.1	14.74
85+	70	160.3	19.16	19	169.4	38.86	51	157.2	22.01
Total deaths	242			116			126		
Connecticut									
. Crude rate		8.2	-		8.1	-		8.3	-
. AAMR (1940 Std. mil.)		3.4	0.25		4.6	0.44		2.7	0.29
. AAMR (1970 Std. mil.)		5.4	0.36		7.2	0.67		4.3	0.40
. AAMR (2000 Std. mil.)		7.9	0.51		10.5	1.01		6.3	0.57
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		5.7	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis-related (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.88	2	2.0	1.40	1	1.0	1.05
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.9	0.63	1	0.9	0.87	1	0.9	0.90
25-29	7	2.8	1.04	2	1.6	1.11	5	4.0	1.78
30-34	8	3.0	1.08	7	5.3	2.02	1	0.8	0.76
35-39	9	3.8	1.28	6	5.2	2.11	3	2.5	1.47
40-44	14	6.4	1.71	12	11.2	3.23	2	1.8	1.27
45-49	26	14.6	2.87	15	17.2	4.44	11	12.2	3.67
50-54	26	18.3	3.59	15	21.6	5.58	11	15.1	4.57
55-59	47	34.0	4.96	31	46.3	8.32	16	22.4	5.61
60-64	105	73.4	7.16	60	88.8	11.46	45	59.6	8.88
65-69	165	124.4	9.69	97	162.8	16.53	68	93.1	11.29
70-74	212	193.1	13.26	132	284.6	24.77	80	126.1	14.10
75-79	259	315.2	19.58	156	491.5	39.35	103	204.2	20.12
80-84	247	475.8	30.27	136	785.6	67.36	111	320.8	30.45
85+	361	826.8	43.52	137	1221.4	104.35	224	690.4	46.13
Total deaths	1,491			809			682		
Connecticut									
. Crude rate		50.5	-		56.6	-		44.9	-
. AAMR (1940 Std. mil.)		23.5	0.69		33.3	1.23		16.9	0.79
. AAMR (1970 Std. mil.)		34.7	0.92		50.2	1.77		25.1	1.02
. AAMR (2000 Std. mil.)		48.3	1.26		71.1	2.62		35.2	1.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	15	7.6	1.97	11	10.9	3.28	4	4.2	2.09
5-9	8	4.5	1.60	6	6.6	2.70	2	2.3	1.64
10-14	15	9.0	2.33	10	11.8	3.72	5	6.2	2.75
15-19	82	42.3	4.68	64	64.4	8.05	18	19.1	4.50
20-24	90	39.8	4.20	74	64.6	7.51	16	14.4	3.59
25-29	56	22.1	2.95	43	33.7	5.15	13	10.3	2.86
30-34	56	21.3	2.85	46	35.1	5.17	10	7.6	2.40
35-39	46	19.6	2.89	40	34.4	5.44	6	5.1	2.08
40-44	35	16.0	2.70	25	23.3	4.66	10	9.0	2.83
45-49	23	13.0	2.70	20	22.9	5.13	3	3.3	1.92
50-54	27	19.0	3.66	20	28.8	6.44	7	9.6	3.64
55-59	31	22.4	4.03	23	34.4	7.17	8	11.2	3.96
60-64	40	28.0	4.42	32	47.3	8.37	8	10.6	3.75
65-69	42	31.7	4.89	22	36.9	7.87	20	27.4	6.12
70-74	56	51.0	6.81	34	73.3	12.57	22	34.7	7.39
75-79	57	69.4	9.19	34	107.1	18.37	23	45.6	9.51
80-84	58	111.7	14.67	29	167.5	31.11	29	83.8	15.56
85+	101	231.3	23.02	33	294.2	51.21	68	209.6	25.42
Total deaths	838			566			272		
Connecticut									
. Crude rate		28.4	-		39.6	-		17.9	-
. AAMR (1940 Std. mil.)		23.1	0.89		35.6	1.57		11.2	0.85
. AAMR (1970 Std. mil.)		24.7	0.90		37.5	1.62		13.0	0.89
. AAMR (2000 Std. mil.)		27.5	0.96		41.4	1.81		15.4	0.95
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		33.8	-		49.4	-		18.9	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.85	2	2.4	1.66	0	0.0	0.00
15-19	2	1.0	0.73	2	2.0	1.42	0	0.0	0.00
20-24	1	0.4	0.44	0	0.0	0.00	1	0.9	0.90
25-29	4	1.6	0.79	3	2.4	1.36	1	0.8	0.79
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	3	1.3	0.74	3	2.6	1.49	0	0.0	0.00
40-44	3	1.4	0.79	2	1.9	1.32	1	0.9	0.90
45-49	1	0.6	0.56	1	1.1	1.15	0	0.0	0.00
50-54	2	1.4	1.00	1	1.4	1.44	1	1.4	1.38
55-59	4	2.9	1.45	3	4.5	2.59	1	1.4	1.40
60-64	11	7.7	2.32	8	11.8	4.19	3	4.0	2.29
65-69	6	4.5	1.85	4	6.7	3.36	2	2.7	1.94
70-74	17	15.5	3.75	12	25.9	7.47	5	7.9	3.53
75-79	17	20.7	5.02	8	25.2	8.91	9	17.8	5.95
80-84	22	42.4	9.03	9	52.0	17.33	13	37.6	10.42
85+	62	142.0	18.03	16	142.6	35.66	46	141.8	20.90
Total deaths	158			75			83		
Connecticut									
. Crude rate		5.4	-		5.2	-		5.5	-
. AAMR (1940 Std. mil.)		2.4	0.24		3.5	0.43		1.5	0.22
. AAMR (1970 Std. mil.)		3.6	0.30		4.8	0.57		2.6	0.31
. AAMR (2000 Std. mil.)		5.3	0.42		6.6	0.81		4.2	0.46
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.7	-		3.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.5	1.14	3	3.0	1.71	2	2.1	1.48
5-9	1	0.6	0.56	1	1.1	1.10	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.5	0.89	2	2.0	1.42	1	1.1	1.06
20-24	4	1.8	0.89	4	3.5	1.75	0	0.0	0.00
25-29	1	0.4	0.39	0	0.0	0.00	1	0.8	0.79
30-34	7	2.7	1.01	7	5.3	2.02	0	0.0	0.00
35-39	4	1.7	0.85	4	3.4	1.72	0	0.0	0.00
40-44	2	0.9	0.65	1	0.9	0.93	1	0.9	0.90
45-49	3	1.7	0.98	3	3.4	1.99	0	0.0	0.00
50-54	1	0.7	0.70	0	0.0	0.00	1	1.4	1.38
55-59	2	1.4	1.02	1	1.5	1.50	1	1.4	1.40
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	1	0.8	0.75	1	1.7	1.68	0	0.0	0.00
70-74	3	2.7	1.58	1	2.2	2.16	2	3.2	2.23
75-79	1	1.2	1.22	0	0.0	0.00	1	2.0	1.98
80-84	2	3.9	2.72	0	0.0	0.00	2	5.8	4.09
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	40			28			12		
Connecticut									
. Crude rate		1.4	-		2.0	-		0.8	-
. AAMR (1940 Std. mil.)		1.3	0.21		1.9	0.37		-	-
. AAMR (1970 Std. mil.)		1.3	0.21		1.8	0.35		-	-
. AAMR (2000 Std. mil.)		1.3	0.21		1.8	0.35		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		3.2	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.60	1	1.2	1.18	0	0.0	0.00
15-19	13	6.7	1.86	11	11.1	3.34	2	2.1	1.50
20-24	29	12.8	2.38	27	23.6	4.54	2	1.8	1.27
25-29	26	10.3	2.01	22	17.3	3.68	4	3.2	1.59
30-34	26	9.9	1.94	22	16.8	3.58	4	3.0	1.52
35-39	23	9.8	2.05	19	16.3	3.75	4	3.4	1.69
40-44	28	12.8	2.42	23	21.5	4.47	5	4.5	2.00
45-49	25	14.1	2.82	20	22.9	5.13	5	5.5	2.48
50-54	14	9.9	2.63	7	10.1	3.81	7	9.6	3.64
55-59	14	10.1	2.71	9	13.5	4.49	5	7.0	3.13
60-64	16	11.2	2.80	13	19.2	5.33	3	4.0	2.29
65-69	13	9.8	2.72	13	21.8	6.05	0	0.0	0.00
70-74	15	13.7	3.53	12	25.9	7.47	3	4.7	2.73
75-79	10	12.2	3.85	6	18.9	7.72	4	7.9	3.97
80-84	5	9.6	4.31	2	11.6	8.17	3	8.7	5.01
85+	6	13.7	5.61	6	53.5	21.84	0	0.0	0.00
Total deaths	264			213			51		
Connecticut									
. Crude rate		8.9	-		14.9	-		3.4	-
. AAMR (1940 Std. mil.)		8.1	0.52		13.4	0.95		3.0	0.44
. AAMR (1970 Std. mil.)		7.7	0.49		13.0	0.91		2.9	0.42
. AAMR (2000 Std. mil.)		8.6	0.54		14.6	1.03		3.3	0.48
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		18.8	-		4.6	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.72	1	1.0	0.99	1	1.0	1.05
5-9	3	1.7	0.98	1	1.1	1.10	2	2.3	1.64
10-14	3	1.8	1.04	2	2.4	1.66	1	1.2	1.23
15-19	15	7.7	2.00	11	11.1	3.34	4	4.2	2.12
20-24	11	4.9	1.47	6	5.2	2.14	5	4.5	2.01
25-29	11	4.3	1.31	10	7.8	2.48	1	0.8	0.79
30-34	19	7.2	1.66	12	9.2	2.64	7	5.3	2.01
35-39	11	4.7	1.42	8	6.9	2.43	3	2.5	1.47
40-44	6	2.7	1.12	3	2.8	1.62	3	2.7	1.55
45-49	11	6.2	1.87	8	9.2	3.24	3	3.3	1.92
50-54	2	1.4	1.00	1	1.4	1.44	1	1.4	1.38
55-59	3	2.2	1.25	2	3.0	2.11	1	1.4	1.40
60-64	6	4.2	1.71	4	5.9	2.96	2	2.6	1.87
65-69	3	2.3	1.31	2	3.4	2.37	1	1.4	1.37
70-74	5	4.6	2.04	1	2.2	2.16	4	6.3	3.15
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	4.6	3.24	0	0.0	0.00	2	6.2	4.36
Total deaths	113			72			41		
Connecticut									
. Crude rate		3.8	-		5.0	-		2.7	-
. AAMR (1940 Std. mil.)		3.9	0.38		5.1	0.61		2.6	0.44
. AAMR (1970 Std. mil.)		3.6	0.36		4.7	0.58		2.6	0.43
. AAMR (2000 Std. mil.)		3.7	0.35		4.7	0.56		2.6	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		9.4	-		14.6	-		4.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	1.02	3	3.0	1.71	1	1.0	1.05
5-9	6	3.4	1.38	4	4.4	2.20	2	2.3	1.64
10-14	9	5.4	1.80	7	8.2	3.11	2	2.5	1.74
15-19	66	34.1	4.20	51	51.3	7.19	15	15.9	4.11
20-24	70	31.0	3.70	57	49.8	6.59	13	11.7	3.24
25-29	40	15.8	2.50	30	23.5	4.30	10	7.9	2.51
30-34	36	13.7	2.28	28	21.4	4.04	8	6.1	2.14
35-39	20	8.5	1.91	19	16.3	3.75	1	0.8	0.85
40-44	19	8.7	1.99	13	12.1	3.36	6	5.4	2.19
45-49	9	5.1	1.69	7	8.0	3.03	2	2.2	1.57
50-54	9	6.3	2.11	7	10.1	3.81	2	2.8	1.95
55-59	9	6.5	2.17	8	12.0	4.23	1	1.4	1.40
60-64	13	9.1	2.52	11	16.3	4.91	2	2.6	1.87
65-69	18	13.6	3.20	8	13.4	4.75	10	13.7	4.33
70-74	15	13.7	3.53	8	17.2	6.10	7	11.0	4.17
75-79	20	24.3	5.44	14	44.1	11.79	6	11.9	4.86
80-84	10	19.3	6.09	4	23.1	11.55	6	17.3	7.08
85+	8	18.3	6.48	1	8.9	8.92	7	21.6	8.15
Total deaths	381			280			101		
Connecticut									
. Crude rate		12.9	-		19.6	-		6.6	-
. AAMR (1940 Std. mil.)		12.8	0.69		19.7	1.21		5.9	0.66
. AAMR (1970 Std. mil.)		12.5	0.68		19.1	1.19		6.0	0.65
. AAMR (2000 Std. mil.)		12.3	0.64		18.7	1.14		6.1	0.62
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		18.6	-		26.1	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.85	2	2.4	1.66	0	0.0	0.00
15-19	2	1.0	0.73	2	2.0	1.42	0	0.0	0.00
20-24	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
25-29	4	1.6	0.79	3	2.4	1.36	1	0.8	0.79
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	4	1.7	0.85	4	3.4	1.72	0	0.0	0.00
40-44	1	0.5	0.46	1	0.9	0.93	0	0.0	0.00
45-49	1	0.6	0.56	1	1.1	1.15	0	0.0	0.00
50-54	2	1.4	1.00	1	1.4	1.44	1	1.4	1.38
55-59	4	2.9	1.45	3	4.5	2.59	1	1.4	1.40
60-64	10	7.0	2.21	7	10.4	3.91	3	4.0	2.29
65-69	6	4.5	1.85	4	6.7	3.36	2	2.7	1.94
70-74	14	12.7	3.41	10	21.6	6.82	4	6.3	3.15
75-79	14	17.0	4.55	6	18.9	7.72	8	15.9	5.61
80-84	16	30.8	7.70	5	28.9	12.92	11	31.8	9.58
85+	49	112.2	16.03	12	107.0	30.88	37	114.0	18.75
Total deaths	131			63			68		
Connecticut									
. Crude rate		4.4	-		4.4	-		4.5	-
. AAMR (1940 Std. mil.)		2.1	0.23		3.1	0.42		1.2	0.19
. AAMR (1970 Std. mil.)		3.1	0.28		4.1	0.52		2.1	0.28
. AAMR (2000 Std. mil.)		4.4	0.38		5.3	0.71		3.4	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	3.0	1.34	4	4.7	2.35	1	1.2	1.23
15-19	23	11.9	2.48	18	18.1	4.27	5	5.3	2.37
20-24	19	8.4	1.93	16	14.0	3.49	3	2.7	1.56
25-29	21	8.3	1.81	18	14.1	3.33	3	2.4	1.38
30-34	20	7.6	1.70	15	11.4	2.95	5	3.8	1.70
35-39	17	7.3	1.76	13	11.2	3.10	4	3.4	1.69
40-44	18	8.2	1.94	14	13.1	3.49	4	3.6	1.79
45-49	18	10.1	2.39	13	14.9	4.13	5	5.5	2.48
50-54	7	4.9	1.86	5	7.2	3.22	2	2.8	1.95
55-59	3	2.2	1.25	2	3.0	2.11	1	1.4	1.40
60-64	12	8.4	2.42	11	16.3	4.91	1	1.3	1.32
65-69	10	7.5	2.38	9	15.1	5.03	1	1.4	1.37
70-74	10	9.1	2.88	8	17.2	6.10	2	3.2	2.23
75-79	2	2.4	1.72	2	6.3	4.46	0	0.0	0.00
80-84	2	3.9	2.72	1	5.8	5.78	1	2.9	2.89
85+	3	6.9	3.97	3	26.7	15.44	0	0.0	0.00
Total deaths	190			152			38		
Connecticut									
. Crude rate		6.4	-		10.6	-		2.5	-
. AAMR (1940 Std. mil.)		6.3	0.47		10.1	0.85		2.6	0.43
. AAMR (1970 Std. mil.)		5.9	0.45		9.8	0.82		2.4	0.41
. AAMR (2000 Std. mil.)		6.2	0.46		10.3	0.86		2.5	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		23.7	-		4.1	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	7	3.6	1.37	6	6.0	2.47	1	1.1	1.06
20-24	11	4.9	1.47	11	9.6	2.89	0	0.0	0.00
25-29	12	4.7	1.37	10	7.8	2.48	2	1.6	1.12
30-34	8	3.0	1.08	5	3.8	1.71	3	2.3	1.31
35-39	11	4.7	1.42	9	7.7	2.58	2	1.7	1.20
40-44	12	5.5	1.58	10	9.3	2.95	2	1.8	1.27
45-49	11	6.2	1.87	8	9.2	3.24	3	3.3	1.92
50-54	6	4.2	1.72	4	5.8	2.88	2	2.8	1.95
55-59	2	1.4	1.02	1	1.5	1.50	1	1.4	1.40
60-64	9	6.3	2.10	9	13.3	4.44	0	0.0	0.00
65-69	9	6.8	2.26	9	15.1	5.03	0	0.0	0.00
70-74	7	6.4	2.41	7	15.1	5.70	0	0.0	0.00
75-79	2	2.4	1.72	2	6.3	4.46	0	0.0	0.00
80-84	2	3.9	2.72	1	5.8	5.78	1	2.9	2.89
85+	3	6.9	3.97	3	26.7	15.44	0	0.0	0.00
Total deaths	112			95			17		
Connecticut									
. Crude rate		3.8	-		6.6	-		1.1	-
. AAMR (1940 Std. mil.)		3.4	0.34		5.9	0.62		1.1	0.27
. AAMR (1970 Std. mil.)		3.3	0.32		5.9	0.61		1.0	0.25
. AAMR (2000 Std. mil.)		3.6	0.35		6.5	0.69		1.2	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		12.0	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	1.8	1.04	2	2.4	1.66	1	1.2	1.23
15-19	14	7.2	1.93	10	10.1	3.18	4	4.2	2.12
20-24	7	3.1	1.17	5	4.4	1.95	2	1.8	1.27
25-29	7	2.8	1.04	6	4.7	1.92	1	0.8	0.79
30-34	10	3.8	1.20	8	6.1	2.16	2	1.5	1.07
35-39	6	2.6	1.05	4	3.4	1.72	2	1.7	1.20
40-44	5	2.3	1.02	3	2.8	1.62	2	1.8	1.27
45-49	6	3.4	1.38	4	4.6	2.29	2	2.2	1.57
50-54	1	0.7	0.70	1	1.4	1.44	0	0.0	0.00
55-59	1	0.7	0.72	1	1.5	1.50	0	0.0	0.00
60-64	3	2.1	1.21	2	3.0	2.09	1	1.3	1.32
65-69	1	0.8	0.75	0	0.0	0.00	1	1.4	1.37
70-74	3	2.7	1.58	1	2.2	2.16	2	3.2	2.23
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	67			47			20		
Connecticut									
. Crude rate		2.3	-		3.3	-		1.3	-
. AAMR (1940 Std. mil.)		2.4	0.30		3.4	0.51		1.4	0.32
. AAMR (1970 Std. mil.)		2.2	0.29		3.2	0.48		1.3	0.32
. AAMR (2000 Std. mil.)		2.2	0.28		3.1	0.46		1.3	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		9.9	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	4	2.1	1.03	3	3.0	1.74	1	1.1	1.06
20-24	2	0.9	0.63	1	0.9	0.87	1	0.9	0.90
25-29	5	2.0	0.88	4	3.1	1.57	1	0.8	0.79
30-34	9	3.4	1.14	7	5.3	2.02	2	1.5	1.07
35-39	12	5.1	1.48	10	8.6	2.72	2	1.7	1.20
40-44	11	5.0	1.52	7	6.5	2.47	4	3.6	1.79
45-49	8	4.5	1.59	6	6.9	2.81	2	2.2	1.57
50-54	8	5.6	1.99	4	5.8	2.88	4	5.5	2.75
55-59	6	4.3	1.77	4	6.0	2.99	2	2.8	1.98
60-64	5	3.5	1.56	3	4.4	2.56	2	2.6	1.87
65-69	3	2.3	1.31	3	5.0	2.91	0	0.0	0.00
70-74	4	3.6	1.82	3	6.5	3.73	1	1.6	1.58
75-79	7	8.5	3.22	3	9.5	5.46	4	7.9	3.97
80-84	3	5.8	3.34	2	11.6	8.17	1	2.9	2.89
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	87			60			27		
Connecticut									
. Crude rate		2.9	-		4.2	-		1.8	-
. AAMR (1940 Std. mil.)		2.6	0.29		3.7	0.49		1.5	0.31
. AAMR (1970 Std. mil.)		2.5	0.28		3.6	0.47		1.5	0.30
. AAMR (2000 Std. mil.)		3.0	0.32		4.2	0.56		1.8	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		6.9	-		3.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
30-34	9	3.4	1.14	8	6.1	2.16	1	0.8	0.76
35-39	12	5.1	1.48	11	9.5	2.85	1	0.8	0.85
40-44	12	5.5	1.58	5	4.7	2.09	7	6.3	2.37
45-49	21	11.8	2.58	17	19.5	4.73	4	4.4	2.21
50-54	23	16.2	3.38	12	17.3	4.99	11	15.1	4.57
55-59	29	21.0	3.90	20	29.9	6.69	9	12.6	4.20
60-64	24	16.8	3.42	19	28.1	6.45	5	6.6	2.96
65-69	23	17.3	3.62	17	28.5	6.92	6	8.2	3.35
70-74	10	9.1	2.88	7	15.1	5.70	3	4.7	2.73
75-79	6	7.3	2.98	4	12.6	6.30	2	4.0	2.80
80-84	2	3.9	2.72	1	5.8	5.78	1	2.9	2.89
85+	5	11.5	5.12	2	17.8	12.61	3	9.2	5.34
Total deaths	177			124			53		
Connecticut									
. Crude rate		6.0	-		8.7	-		3.5	-
. AAMR (1940 Std. mil.)		5.0	0.39		7.4	0.68		2.9	0.42
. AAMR (1970 Std. mil.)		5.3	0.40		7.8	0.71		3.0	0.43
. AAMR (2000 Std. mil.)		6.0	0.46		8.8	0.81		3.5	0.49
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.4	-		11.6	-		3.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1989

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.4	0.44	0	0.0	0.00	1	0.9	0.90
25-29	3	1.2	0.68	3	2.4	1.36	0	0.0	0.00
30-34	5	1.9	0.85	4	3.1	1.53	1	0.8	0.76
35-39	6	2.6	1.05	4	3.4	1.72	2	1.7	1.20
40-44	5	2.3	1.02	3	2.8	1.62	2	1.8	1.27
45-49	1	0.6	0.56	0	0.0	0.00	1	1.1	1.11
50-54	2	1.4	1.00	1	1.4	1.44	1	1.4	1.38
55-59	2	1.4	1.02	1	1.5	1.50	1	1.4	1.40
60-64	1	0.7	0.70	0	0.0	0.00	1	1.3	1.32
65-69	1	0.8	0.75	1	1.7	1.68	0	0.0	0.00
70-74	1	0.9	0.91	1	2.2	2.16	0	0.0	0.00
75-79	4	4.9	2.43	2	6.3	4.46	2	4.0	2.80
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	32			20			12		
Connecticut									
. Crude rate		1.1	-		1.4	-		0.8	-
. AAMR (1940 Std. mil.)		0.9	0.17		1.2	0.27		-	-
. AAMR (1970 Std. mil.)		0.8	0.15		1.1	0.24		-	-
. AAMR (2000 Std. mil.)		1.0	0.19		1.3	0.30		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.1	-		5.5	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower