

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.2	1.31	3	3.1	1.80	3	3.3	1.91
5-9	2	1.0	0.72	1	1.0	0.99	1	1.0	1.04
10-14	3	1.6	0.95	2	2.1	1.52	1	1.1	1.12
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	19	9.8	2.25	11	11.3	3.39	8	8.4	2.95
30-34	46	19.1	2.81	33	27.3	4.75	13	10.8	3.00
35-39	69	26.6	3.20	52	40.1	5.56	17	13.1	3.17
40-44	44	18.7	2.81	38	32.7	5.31	6	5.0	2.05
45-49	53	25.1	3.45	38	36.5	5.92	15	14.0	3.62
50-54	29	17.1	3.18	23	27.7	5.78	6	6.9	2.84
55-59	19	14.8	3.39	11	17.6	5.29	8	12.2	4.30
60-64	22	19.5	4.15	16	29.4	7.35	6	10.2	4.18
65-69	40	33.3	5.27	21	38.1	8.31	19	29.3	6.72
70-74	37	33.2	5.45	25	52.0	10.40	12	18.9	5.46
75-79	61	64.1	8.21	29	75.1	13.95	32	56.6	10.01
80-84	83	130.4	14.32	43	186.0	28.36	40	98.7	15.61
85+	133	241.7	20.96	43	292.6	44.62	90	223.2	23.52
Total deaths	667			390			277		
Connecticut									
. Crude rate		23.1	-		27.7	-		18.7	-
. AAMR (1940 Std. mil.)		13.7	0.61		19.3	1.05		8.5	0.66
. AAMR (1970 Std. mil.)		15.4	0.63		21.3	1.10		10.3	0.69
. AAMR (2000 Std. mil.)		20.5	0.80		28.3	1.46		13.9	0.86
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.0	-		27.2	-		10.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	1.6	0.95	2	2.1	1.52	1	1.1	1.12
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	15	7.8	2.00	9	9.2	3.07	6	6.3	2.56
30-34	42	17.4	2.69	30	24.8	4.53	12	10.0	2.89
35-39	60	23.1	2.98	44	33.9	5.12	16	12.3	3.08
40-44	35	14.8	2.51	31	26.7	4.79	4	3.3	1.67
45-49	36	17.1	2.84	30	28.8	5.26	6	5.6	2.29
50-54	18	10.6	2.51	14	16.9	4.51	4	4.6	2.32
55-59	6	4.7	1.91	2	3.2	2.26	4	6.1	3.04
60-64	3	2.7	1.53	3	5.5	3.18	0	0.0	0.00
65-69	2	1.7	1.18	2	3.6	2.57	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	221			168			53		
Connecticut									
. Crude rate		7.6	-		11.9	-		3.6	-
. AAMR (1940 Std. mil.)		6.9	0.47		10.4	0.81		3.4	0.47
. AAMR (1970 Std. mil.)		5.8	0.40		8.9	0.70		2.8	0.40
. AAMR (2000 Std. mil.)		7.3	0.49		11.1	0.86		3.5	0.48
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.1	-		18.1	-		4.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.93	2	2.1	1.47	1	1.1	1.10
5-9	9	4.6	1.52	8	7.9	2.79	1	1.0	1.04
10-14	5	2.7	1.22	2	2.1	1.52	3	3.4	1.93
15-19	4	2.4	1.19	3	3.5	2.00	1	1.2	1.24
20-24	9	5.7	1.90	6	7.4	3.01	3	3.9	2.26
25-29	15	7.8	2.00	10	10.2	3.24	5	5.2	2.33
30-34	33	13.7	2.38	20	16.5	3.70	13	10.8	3.00
35-39	61	23.5	3.01	26	20.0	3.93	35	26.9	4.55
40-44	100	42.4	4.24	44	37.9	5.71	56	46.8	6.26
45-49	190	90.0	6.53	87	83.5	8.95	103	96.4	9.50
50-54	327	193.1	10.68	153	184.4	14.90	174	201.5	15.27
55-59	369	287.1	14.95	197	314.3	22.39	172	261.3	19.92
60-64	534	472.3	20.44	272	499.9	30.31	262	446.6	27.59
65-69	783	652.5	23.32	414	751.1	36.91	369	568.8	29.61
70-74	1,083	971.1	29.51	574	1193.8	49.83	509	802.3	35.56
75-79	1,168	1227.7	35.92	589	1525.7	62.87	579	1024.2	42.57
80-84	989	1554.2	49.42	504	2179.9	97.10	485	1197.1	54.36
85+	979	1779.2	56.86	417	2837.3	138.94	562	1393.6	58.79
Total deaths	6,661			3,328			3,333		
Connecticut									
. Crude rate		230.5	-		236.5	-		224.8	-
. AAMR (1940 Std. mil.)		119.5	1.68		138.4	2.61		106.7	2.21
. AAMR (1970 Std. mil.)		159.0	2.05		190.9	3.37		138.9	2.62
. AAMR (2000 Std. mil.)		200.6	2.47		248.1	4.36		173.0	3.06
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		127.8	-		153.9	-		108.7	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.42	0	0.0	0.00	1	0.8	0.84
45-49	2	0.9	0.67	2	1.9	1.36	0	0.0	0.00
50-54	6	3.5	1.45	6	7.2	2.95	0	0.0	0.00
55-59	6	4.7	1.91	5	8.0	3.57	1	1.5	1.52
60-64	16	14.2	3.54	9	16.5	5.51	7	11.9	4.51
65-69	12	10.0	2.89	9	16.3	5.44	3	4.6	2.67
70-74	22	19.7	4.21	12	25.0	7.20	10	15.8	4.98
75-79	8	8.4	2.97	5	13.0	5.79	3	5.3	3.06
80-84	7	11.0	4.16	2	8.7	6.12	5	12.3	5.52
85+	7	12.7	4.81	3	20.4	11.79	4	9.9	4.96
Total deaths	87			53			34		
Connecticut									
. Crude rate		3.0	-		3.8	-		2.3	-
. AAMR (1940 Std. mil.)		1.9	0.22		2.7	0.39		1.2	0.23
. AAMR (1970 Std. mil.)		2.4	0.26		3.3	0.46		1.5	0.28
. AAMR (2000 Std. mil.)		2.7	0.29		3.8	0.53		1.8	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.2	-		1.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.41	1	0.8	0.83	0	0.0	0.00
35-39	3	1.2	0.67	2	1.5	1.09	1	0.8	0.77
40-44	8	3.4	1.20	6	5.2	2.11	2	1.7	1.18
45-49	11	5.2	1.57	8	7.7	2.72	3	2.8	1.62
50-54	21	12.4	2.71	15	18.1	4.67	6	6.9	2.84
55-59	29	22.6	4.19	16	25.5	6.38	13	19.7	5.48
60-64	39	34.5	5.52	24	44.1	9.00	15	25.6	6.60
65-69	77	64.2	7.31	47	85.3	12.44	30	46.2	8.44
70-74	119	106.7	9.78	63	131.0	16.51	56	88.3	11.80
75-79	117	123.0	11.37	58	150.2	19.73	59	104.4	13.59
80-84	112	176.0	16.63	50	216.3	30.58	62	153.0	19.44
85+	155	281.7	22.63	56	381.0	50.92	99	245.5	24.67
Total deaths	693			347			346		
Connecticut									
. Crude rate		24.0	-		24.7	-		23.3	-
. AAMR (1940 Std. mil.)		10.8	0.48		13.9	0.81		8.3	0.56
. AAMR (1970 Std. mil.)		15.4	0.62		19.6	1.07		12.2	0.72
. AAMR (2000 Std. mil.)		20.4	0.78		26.2	1.43		16.5	0.91
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.3	-		14.8	-		10.3	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.42	1	0.9	0.86	0	0.0	0.00
45-49	5	2.4	1.06	5	4.8	2.15	0	0.0	0.00
50-54	8	4.7	1.67	6	7.2	2.95	2	2.3	1.64
55-59	5	3.9	1.74	3	4.8	2.76	2	3.0	2.15
60-64	7	6.2	2.34	3	5.5	3.18	4	6.8	3.41
65-69	13	10.8	3.00	5	9.1	4.06	8	12.3	4.36
70-74	23	20.6	4.30	15	31.2	8.06	8	12.6	4.46
75-79	19	20.0	4.58	12	31.1	8.97	7	12.4	4.68
80-84	16	25.1	6.29	4	17.3	8.65	12	29.6	8.55
85+	20	36.3	8.13	7	47.6	18.00	13	32.2	8.94
Total deaths	117			61			56		
Connecticut									
. Crude rate		4.0	-		4.3	-		3.8	-
. AAMR (1940 Std. mil.)		2.0	0.21		2.6	0.36		1.5	0.25
. AAMR (1970 Std. mil.)		2.7	0.26		3.5	0.45		2.1	0.31
. AAMR (2000 Std. mil.)		3.5	0.33		4.5	0.58		2.7	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		2.5	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
35-39	3	1.2	0.67	3	2.3	1.34	0	0.0	0.00
40-44	3	1.3	0.73	2	1.7	1.22	1	0.8	0.84
45-49	14	6.6	1.77	9	8.6	2.88	5	4.7	2.09
50-54	13	7.7	2.13	9	10.8	3.61	4	4.6	2.32
55-59	19	14.8	3.39	9	14.4	4.79	10	15.2	4.80
60-64	27	23.9	4.60	12	22.1	6.37	15	25.6	6.60
65-69	47	39.2	5.71	28	50.8	9.60	19	29.3	6.72
70-74	77	69.0	7.87	36	74.9	12.48	41	64.6	10.09
75-79	74	77.8	9.04	31	80.3	14.42	43	76.1	11.60
80-84	55	86.4	11.65	23	99.5	20.74	32	79.0	13.96
85+	62	112.7	14.31	18	122.5	28.87	44	109.1	16.45
Total deaths	396			182			214		
Connecticut									
. Crude rate		13.7	-		12.9	-		14.4	-
. AAMR (1940 Std. mil.)		6.8	0.39		7.7	0.61		5.9	0.49
. AAMR (1970 Std. mil.)		9.2	0.49		10.4	0.78		8.3	0.62
. AAMR (2000 Std. mil.)		11.8	0.60		13.3	1.00		10.6	0.74
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		7.2	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
35-39	8	3.1	1.09	3	2.3	1.34	5	3.8	1.72
40-44	20	8.5	1.90	9	7.7	2.58	11	9.2	2.77
45-49	37	17.5	2.88	20	19.2	4.29	17	15.9	3.86
50-54	93	54.9	5.69	50	60.2	8.52	43	49.8	7.59
55-59	132	102.7	8.94	81	129.2	14.36	51	77.5	10.85
60-64	188	166.3	12.13	106	194.8	18.92	82	139.8	15.44
65-69	269	224.2	13.67	143	259.4	21.69	126	194.2	17.30
70-74	342	306.7	16.58	212	440.9	30.28	130	204.9	17.97
75-79	330	346.9	19.10	180	466.3	34.75	150	265.3	21.67
80-84	264	414.9	25.53	146	631.5	52.26	118	291.3	26.81
85+	148	269.0	22.11	71	483.1	57.33	77	190.9	21.76
Total deaths	1,834			1,024			810		
Connecticut									
. Crude rate		63.5	-		72.8	-		54.6	-
. AAMR (1940 Std. mil.)		35.1	0.92		44.6	1.49		27.9	1.13
. AAMR (1970 Std. mil.)		46.0	1.12		60.0	1.91		36.0	1.36
. AAMR (2000 Std. mil.)		55.5	1.30		74.2	2.34		43.0	1.54
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.2	-		54.1	-		27.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	2	1.0	0.73	1	1.0	1.02	1	1.0	1.04
30-34	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
35-39	4	1.5	0.77	2	1.5	1.09	2	1.5	1.09
40-44	6	2.5	1.04	3	2.6	1.49	3	2.5	1.45
45-49	6	2.8	1.16	3	2.9	1.66	3	2.8	1.62
50-54	8	4.7	1.67	5	6.0	2.69	3	3.5	2.01
55-59	8	6.2	2.20	5	8.0	3.57	3	4.6	2.63
60-64	7	6.2	2.34	5	9.2	4.11	2	3.4	2.41
65-69	14	11.7	3.12	12	21.8	6.28	2	3.1	2.18
70-74	13	11.7	3.23	9	18.7	6.24	4	6.3	3.15
75-79	13	13.7	3.79	9	23.3	7.77	4	7.1	3.54
80-84	12	18.9	5.44	8	34.6	12.23	4	9.9	4.94
85+	8	14.5	5.14	6	40.8	16.67	2	5.0	3.51
Total deaths	104			71			33		
Connecticut									
. Crude rate		3.6	-		5.0	-		2.2	-
. AAMR (1940 Std. mil.)		2.3	0.25		3.4	0.43		1.4	0.28
. AAMR (1970 Std. mil.)		2.7	0.27		4.2	0.50		1.6	0.29
. AAMR (2000 Std. mil.)		3.2	0.32		5.2	0.62		1.9	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.5	0.52	-	-	-	1	1.0	1.04
30-34	2	0.8	0.59	-	-	-	2	1.7	1.18
35-39	9	3.5	1.15	-	-	-	9	6.9	2.31
40-44	17	7.2	1.75	-	-	-	17	14.2	3.45
45-49	38	18.0	2.92	-	-	-	38	35.6	5.77
50-54	50	29.5	4.18	-	-	-	50	57.9	8.19
55-59	37	28.8	4.73	-	-	-	37	56.2	9.24
60-64	55	48.6	6.56	-	-	-	55	93.8	12.64
65-69	62	51.7	6.56	-	-	-	62	95.6	12.14
70-74	65	58.3	7.23	-	-	-	65	102.5	12.71
75-79	87	91.4	9.80	-	-	-	87	153.9	16.50
80-84	63	99.0	12.47	-	-	-	63	155.5	19.59
85+	78	141.8	16.05	-	-	-	78	193.4	21.90
Total deaths	564			-			564		
Connecticut									
. Crude rate		-	-		-	-		38.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		20.9	1.02
. AAMR (1970 Std. mil.)		-	-		-	-		25.5	1.16
. AAMR (2000 Std. mil.)		-	-		-	-		30.9	1.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		20.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.6	0.63	-	-	-	1	1.3	1.30
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	1	0.4	0.41	-	-	-	1	0.8	0.83
35-39	2	0.8	0.54	-	-	-	2	1.5	1.09
40-44	3	1.3	0.73	-	-	-	3	2.5	1.45
45-49	4	1.9	0.95	-	-	-	4	3.7	1.87
50-54	4	2.4	1.18	-	-	-	4	4.6	2.32
55-59	2	1.6	1.10	-	-	-	2	3.0	2.15
60-64	4	3.5	1.77	-	-	-	4	6.8	3.41
65-69	2	1.7	1.18	-	-	-	2	3.1	2.18
70-74	6	5.4	2.20	-	-	-	6	9.5	3.86
75-79	4	4.2	2.10	-	-	-	4	7.1	3.54
80-84	5	7.9	3.51	-	-	-	5	12.3	5.52
85+	4	7.3	3.63	-	-	-	4	9.9	4.96
Total deaths	42			-			42		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.8	0.32
. AAMR (1970 Std. mil.)		-	-		-	-		2.0	0.33
. AAMR (2000 Std. mil.)		-	-		-	-		2.4	0.38
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is higher
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.4	0.42	-	-	-	1	0.8	0.84
45-49	1	0.5	0.47	-	-	-	1	0.9	0.94
50-54	2	1.2	0.84	-	-	-	2	2.3	1.64
55-59	1	0.8	0.78	-	-	-	1	1.5	1.52
60-64	3	2.7	1.53	-	-	-	3	5.1	2.95
65-69	9	7.5	2.50	-	-	-	9	13.9	4.62
70-74	7	6.3	2.37	-	-	-	7	11.0	4.17
75-79	10	10.5	3.32	-	-	-	10	17.7	5.59
80-84	10	15.7	4.97	-	-	-	10	24.7	7.81
85+	6	10.9	4.45	-	-	-	6	14.9	6.07
Total deaths	50			-			50		
Connecticut									
. Crude rate		-	-		-	-		3.4	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.5	0.25
. AAMR (1970 Std. mil.)		-	-		-	-		2.0	0.31
. AAMR (2000 Std. mil.)		-	-		-	-		2.5	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	1	0.4	0.41	-	-	-	1	0.8	0.83
35-39	1	0.4	0.38	-	-	-	1	0.8	0.77
40-44	2	0.8	0.60	-	-	-	2	1.7	1.18
45-49	6	2.8	1.16	-	-	-	6	5.6	2.29
50-54	10	5.9	1.87	-	-	-	10	11.6	3.66
55-59	9	7.0	2.33	-	-	-	9	13.7	4.56
60-64	12	10.6	3.06	-	-	-	12	20.5	5.91
65-69	14	11.7	3.12	-	-	-	14	21.6	5.77
70-74	22	19.7	4.21	-	-	-	22	34.7	7.39
75-79	27	28.4	5.46	-	-	-	27	47.8	9.19
80-84	12	18.9	5.44	-	-	-	12	29.6	8.55
85+	27	49.1	9.44	-	-	-	27	67.0	12.89
Total deaths	143			-			143		
Connecticut									
. Crude rate		-	-		-	-		9.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		4.8	0.47
. AAMR (1970 Std. mil.)		-	-		-	-		6.1	0.55
. AAMR (2000 Std. mil.)		-	-		-	-		7.5	0.65
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	5	2.5	1.13	4	3.9	1.97	1	1.0	1.04
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	0	0.0	0.00	1	1.2	1.24
20-24	2	1.3	0.89	2	2.5	1.74	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.41	0	0.0	0.00	1	0.8	0.83
35-39	2	0.8	0.54	1	0.8	0.77	1	0.8	0.77
40-44	5	2.1	0.95	2	1.7	1.22	3	2.5	1.45
45-49	11	5.2	1.57	7	6.7	2.54	4	3.7	1.87
50-54	17	10.0	2.43	7	8.4	3.19	10	11.6	3.66
55-59	13	10.1	2.81	8	12.8	4.51	5	7.6	3.40
60-64	13	11.5	3.19	9	16.5	5.51	4	6.8	3.41
65-69	19	15.8	3.63	10	18.1	5.74	9	13.9	4.62
70-74	24	21.5	4.39	11	22.9	6.90	13	20.5	5.68
75-79	10	10.5	3.32	6	15.5	6.35	4	7.1	3.54
80-84	15	23.6	6.09	7	30.3	11.44	8	19.7	6.98
85+	10	18.2	5.75	5	34.0	15.21	5	12.4	5.54
Total deaths	148			79			69		
Connecticut									
. Crude rate		5.1	-		5.6	-		4.7	-
. AAMR (1940 Std. mil.)		3.5	0.32		4.2	0.50		3.0	0.40
. AAMR (1970 Std. mil.)		4.2	0.36		5.0	0.58		3.4	0.44
. AAMR (2000 Std. mil.)		4.7	0.39		5.8	0.65		4.0	0.49
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.4	-		4.4	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.76	1	1.0	1.04	1	1.1	1.10
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	1	0.6	0.63	0	0.0	0.00	1	1.3	1.30
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	4	1.7	0.83	3	2.5	1.43	1	0.8	0.83
35-39	5	1.9	0.86	2	1.5	1.09	3	2.3	1.33
40-44	6	2.5	1.04	4	3.4	1.72	2	1.7	1.18
45-49	8	3.8	1.34	7	6.7	2.54	1	0.9	0.94
50-54	10	5.9	1.87	8	9.6	3.41	2	2.3	1.64
55-59	8	6.2	2.20	4	6.4	3.19	4	6.1	3.04
60-64	15	13.3	3.43	5	9.2	4.11	10	17.0	5.39
65-69	19	15.8	3.63	10	18.1	5.74	9	13.9	4.62
70-74	38	34.1	5.53	25	52.0	10.40	13	20.5	5.68
75-79	49	51.5	7.36	22	57.0	12.15	27	47.8	9.19
80-84	37	58.1	9.56	18	77.9	18.35	19	46.9	10.76
85+	35	63.6	10.75	19	129.3	29.66	16	39.7	9.92
Total deaths	241			132			109		
Connecticut									
. Crude rate		8.3	-		9.4	-		7.4	-
. AAMR (1940 Std. mil.)		4.3	0.32		5.6	0.53		3.3	0.39
. AAMR (1970 Std. mil.)		5.7	0.39		7.5	0.66		4.4	0.47
. AAMR (2000 Std. mil.)		7.3	0.47		9.9	0.88		5.6	0.55
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.2	-		3.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.59	0	0.0	0.00	2	1.7	1.18
35-39	6	2.3	0.94	3	2.3	1.34	3	2.3	1.33
40-44	10	4.2	1.34	5	4.3	1.92	5	4.2	1.87
45-49	14	6.6	1.77	5	4.8	2.15	9	8.4	2.81
50-54	21	12.4	2.71	12	14.5	4.17	9	10.4	3.47
55-59	30	23.3	4.26	15	23.9	6.18	15	22.8	5.88
60-64	42	37.1	5.73	23	42.3	8.81	19	32.4	7.43
65-69	58	48.3	6.35	29	52.6	9.77	29	44.7	8.30
70-74	83	74.4	8.17	43	89.4	13.64	40	63.1	9.97
75-79	108	113.5	10.92	51	132.1	18.50	57	100.8	13.36
80-84	115	180.7	16.85	58	250.9	32.94	57	140.7	18.64
85+	137	249.0	21.27	42	285.8	44.10	95	235.6	24.17
Total deaths	626			286			340		
Connecticut									
. Crude rate		21.7	-		20.3	-		22.9	-
. AAMR (1940 Std. mil.)		9.9	0.46		11.2	0.72		9.0	0.61
. AAMR (1970 Std. mil.)		13.9	0.59		16.1	0.97		12.5	0.75
. AAMR (2000 Std. mil.)		18.6	0.75		21.7	1.30		16.7	0.93
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.8	-		14.9	-		12.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	4	1.7	0.83	1	0.8	0.83	3	2.5	1.44
35-39	8	3.1	1.09	5	3.9	1.72	3	2.3	1.33
40-44	23	9.8	2.03	12	10.3	2.98	11	9.2	2.77
45-49	37	17.5	2.88	21	20.2	4.40	16	15.0	3.74
50-54	48	28.3	4.09	30	36.1	6.60	18	20.8	4.91
55-59	90	70.0	7.38	53	84.6	11.62	37	56.2	9.24
60-64	133	117.6	10.20	77	141.5	16.13	56	95.5	12.76
65-69	234	195.0	12.75	139	252.2	21.39	95	146.4	15.02
70-74	324	290.5	16.14	166	345.3	26.80	158	249.1	19.81
75-79	429	450.9	21.77	211	546.6	37.63	218	385.6	26.12
80-84	446	700.9	33.19	200	865.1	61.17	246	607.2	38.71
85+	542	985.0	42.31	171	1163.5	88.98	371	920.0	47.76
Total deaths	2,318			1,086			1,232		
Connecticut									
. Crude rate		80.2	-		77.2	-		83.1	-
. AAMR (1940 Std. mil.)		34.3	0.84		41.6	1.38		28.6	1.03
. AAMR (1970 Std. mil.)		50.1	1.10		60.8	1.87		42.3	1.33
. AAMR (2000 Std. mil.)		68.0	1.42		82.4	2.54		58.4	1.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.54	1	0.8	0.77	1	0.8	0.77
40-44	1	0.4	0.42	0	0.0	0.00	1	0.8	0.84
45-49	1	0.5	0.47	1	1.0	0.96	0	0.0	0.00
50-54	2	1.2	0.84	2	2.4	1.70	0	0.0	0.00
55-59	1	0.8	0.78	1	1.6	1.60	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	5	4.5	2.01	3	6.2	3.60	2	3.2	2.23
75-79	1	1.1	1.05	1	2.6	2.59	0	0.0	0.00
80-84	1	1.6	1.57	0	0.0	0.00	1	2.5	2.47
85+	1	1.8	1.82	0	0.0	0.00	1	2.5	2.48
Total deaths	15			9			6		
Connecticut									
. Crude rate		0.5	-		0.6	-		0.4	-
. AAMR (1940 Std. mil.)		0.3	0.09		-	-		-	-
. AAMR (1970 Std. mil.)		0.4	0.10		-	-		-	-
. AAMR (2000 Std. mil.)		0.5	0.12		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.4	-		0.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	9	4.8	1.61	6	6.2	2.55	3	3.3	1.91
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
15-19	4	2.4	1.19	2	2.3	1.63	2	2.5	1.75
20-24	8	5.1	1.79	5	6.1	2.75	3	3.9	2.26
25-29	10	5.2	1.63	5	5.1	2.29	5	5.2	2.33
30-34	21	8.7	1.90	18	14.9	3.51	3	2.5	1.44
35-39	39	15.0	2.40	29	22.4	4.15	10	7.7	2.43
40-44	80	33.9	3.79	57	49.1	6.50	23	19.2	4.01
45-49	137	64.9	5.55	104	99.8	9.79	33	30.9	5.38
50-54	187	110.4	8.07	134	161.5	13.95	53	61.4	8.43
55-59	261	203.1	12.57	180	287.2	21.41	81	123.0	13.67
60-64	377	333.4	17.17	257	472.3	29.46	120	204.6	18.67
65-69	640	533.4	21.08	421	763.8	37.22	219	337.6	22.81
70-74	998	894.9	28.33	567	1179.3	49.53	431	679.4	32.72
75-79	1,363	1432.7	38.81	731	1893.6	70.04	632	1118.0	44.47
80-84	1,707	2682.6	64.93	775	3352.1	120.41	932	2300.5	75.36
85+	3,489	6340.9	107.35	998	6790.5	214.95	2,491	6177.0	123.76
Total deaths	9,331			4,290			5,041		
Connecticut									
. Crude rate		322.9	-		304.9	-		339.9	-
. AAMR (1940 Std. mil.)		119.2	1.52		157.8	2.66		87.8	1.67
. AAMR (1970 Std. mil.)		185.4	2.04		237.3	3.68		145.5	2.29
. AAMR (2000 Std. mil.)		271.8	2.82		337.7	5.26		223.5	3.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		134.4	-		179.0	-		98.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Heart disease-related (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	29	15.5	2.88	17	17.7	4.29	12	13.2	3.82
5-9	9	4.6	1.52	6	5.9	2.42	3	3.1	1.80
10-14	6	3.3	1.34	3	3.2	1.86	3	3.4	1.93
15-19	13	7.8	2.15	8	9.2	3.27	5	6.2	2.77
20-24	13	8.2	2.28	8	9.8	3.48	5	6.5	2.92
25-29	19	9.8	2.25	13	13.3	3.69	6	6.3	2.56
30-34	51	21.2	2.96	37	30.6	5.03	14	11.7	3.12
35-39	98	37.7	3.81	65	50.1	6.22	33	25.4	4.42
40-44	141	59.8	5.04	102	87.8	8.69	39	32.6	5.22
45-49	245	116.1	7.42	164	157.5	12.29	81	75.8	8.42
50-54	315	186.0	10.48	208	250.6	17.38	107	123.9	11.98
55-59	439	341.6	16.30	276	440.4	26.51	163	247.6	19.39
60-64	640	566.0	22.37	407	748.0	37.08	233	397.2	26.02
65-69	1,088	906.7	27.49	670	1215.5	46.96	418	644.3	31.52
70-74	1,728	1549.5	37.27	960	1996.7	64.44	768	1210.6	43.68
75-79	2,384	2505.9	51.32	1,226	3175.8	90.70	1,158	2048.5	60.20
80-84	2,804	4406.5	83.22	1,310	5666.1	156.55	1,494	3687.7	95.41
85+	5,380	9777.6	133.30	1,602	10900.2	272.33	3,778	9368.4	152.42
Total deaths	15,402			7,082			8,320		
Connecticut									
. Crude rate		532.9	-		503.3	-		561.1	-
. AAMR (1940 Std. mil.)		203.4	2.01		260.6	3.42		158.2	2.33
. AAMR (1970 Std. mil.)		311.0	2.67		391.0	4.72		251.9	3.10
. AAMR (2000 Std. mil.)		449.6	3.64		555.9	6.73		375.8	4.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									No comparison is available
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.0	0.73	1	1.0	1.02	1	1.0	1.04
30-34	11	4.6	1.38	10	8.3	2.61	1	0.8	0.83
35-39	21	8.1	1.76	18	13.9	3.27	3	2.3	1.33
40-44	63	26.7	3.37	46	39.6	5.84	17	14.2	3.45
45-49	104	49.3	4.83	83	79.7	8.75	21	19.7	4.29
50-54	141	83.3	7.01	110	132.5	12.64	31	35.9	6.45
55-59	193	150.2	10.81	142	226.6	19.01	51	77.5	10.85
60-64	280	247.6	14.80	205	376.8	26.32	75	127.9	14.76
65-69	468	390.0	18.03	322	584.2	32.56	146	225.1	18.63
70-74	746	668.9	24.49	444	923.5	43.83	302	476.0	27.39
75-79	1,020	1072.2	33.57	564	1461.0	61.52	456	806.7	37.77
80-84	1,211	1903.1	54.69	562	2430.8	102.54	649	1602.0	62.88
85+	2,463	4476.2	90.19	731	4973.8	183.96	1,732	4294.9	103.20
Total deaths	6,725			3,240			3,485		
Connecticut									
. Crude rate		232.7	-		230.3	-		235.0	-
. AAMR (1940 Std. mil.)		86.2	1.29		120.0	2.32		58.8	1.33
. AAMR (1970 Std. mil.)		134.1	1.74		179.6	3.20		99.1	1.86
. AAMR (2000 Std. mil.)		195.9	2.40		254.1	4.55		153.7	2.64
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		104.7	-		143.0	-		73.3	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	2	0.9	0.67	0	0.0	0.00	2	1.9	1.32
50-54	3	1.8	1.02	0	0.0	0.00	3	3.5	2.01
55-59	3	2.3	1.35	3	4.8	2.76	0	0.0	0.00
60-64	5	4.4	1.98	3	5.5	3.18	2	3.4	2.41
65-69	17	14.2	3.44	10	18.1	5.74	7	10.8	4.08
70-74	45	40.4	6.02	17	35.4	8.58	28	44.1	8.34
75-79	58	61.0	8.01	26	67.4	13.21	32	56.6	10.01
80-84	99	155.6	15.64	37	160.0	26.31	62	153.0	19.44
85+	271	492.5	29.92	64	435.5	54.43	207	513.3	35.68
Total deaths	503			160			343		
Connecticut									
. Crude rate		17.4	-		11.4	-		23.1	-
. AAMR (1940 Std. mil.)		4.6	0.25		4.5	0.39		4.6	0.32
. AAMR (1970 Std. mil.)		8.6	0.40		8.3	0.66		8.6	0.50
. AAMR (2000 Std. mil.)		14.3	0.64		13.6	1.10		14.5	0.79
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.0	-		6.8	-		5.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.42	1	0.9	0.86	0	0.0	0.00
45-49	1	0.5	0.47	1	1.0	0.96	0	0.0	0.00
50-54	3	1.8	1.02	2	2.4	1.70	1	1.2	1.16
55-59	1	0.8	0.78	1	1.6	1.60	0	0.0	0.00
60-64	6	5.3	2.17	4	7.4	3.68	2	3.4	2.41
65-69	5	4.2	1.86	1	1.8	1.81	4	6.2	3.08
70-74	14	12.6	3.36	3	6.2	3.60	11	17.3	5.23
75-79	20	21.0	4.70	9	23.3	7.77	11	19.5	5.87
80-84	20	31.4	7.03	4	17.3	8.65	16	39.5	9.87
85+	42	76.3	11.78	8	54.4	19.24	34	84.3	14.46
Total deaths	113			34			79		
Connecticut									
. Crude rate		3.9	-		2.4	-		5.3	-
. AAMR (1940 Std. mil.)		1.4	0.16		1.3	0.24		1.4	0.20
. AAMR (1970 Std. mil.)		2.2	0.22		1.9	0.33		2.3	0.29
. AAMR (2000 Std. mil.)		3.3	0.31		2.7	0.47		3.5	0.40
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.5	-		2.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.93	2	2.1	1.47	1	1.1	1.10
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.55	1	1.1	1.07	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	1	0.5	0.52	0	0.0	0.00	1	1.0	1.04
30-34	5	2.1	0.93	4	3.3	1.65	1	0.8	0.83
35-39	5	1.9	0.86	4	3.1	1.54	1	0.8	0.77
40-44	10	4.2	1.34	4	3.4	1.72	6	5.0	2.05
45-49	18	8.5	2.01	8	7.7	2.72	10	9.4	2.96
50-54	17	10.0	2.43	7	8.4	3.19	10	11.6	3.66
55-59	33	25.7	4.47	22	35.1	7.48	11	16.7	5.04
60-64	48	42.5	6.13	30	55.1	10.07	18	30.7	7.23
65-69	95	79.2	8.12	42	76.2	11.76	53	81.7	11.22
70-74	153	137.2	11.09	80	166.4	18.60	73	115.1	13.47
75-79	267	280.7	17.18	118	305.7	28.14	149	263.6	21.59
80-84	398	625.5	31.35	159	687.7	54.54	239	589.9	38.16
85+	775	1408.5	50.59	192	1306.4	94.28	583	1445.7	59.87
Total deaths	1,830			674			1,156		
Connecticut									
. Crude rate		63.3	-		47.9	-		78.0	-
. AAMR (1940 Std. mil.)		20.3	0.59		22.2	0.95		18.8	0.74
. AAMR (1970 Std. mil.)		34.1	0.84		36.2	1.41		32.1	1.04
. AAMR (2000 Std. mil.)		52.7	1.23		54.7	2.15		50.6	1.51
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.5	-		28.5	-		24.6	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.38	1	0.8	0.77	0	0.0	0.00
40-44	1	0.4	0.42	1	0.9	0.86	0	0.0	0.00
45-49	1	0.5	0.47	1	1.0	0.96	0	0.0	0.00
50-54	3	1.8	1.02	3	3.6	2.09	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.9	0.88	0	0.0	0.00	1	1.7	1.70
65-69	3	2.5	1.44	3	5.4	3.14	0	0.0	0.00
70-74	15	13.5	3.47	7	14.6	5.50	8	12.6	4.46
75-79	30	31.5	5.76	13	33.7	9.34	17	30.1	7.29
80-84	32	50.3	8.89	18	77.9	18.35	14	34.6	9.24
85+	118	214.5	19.74	29	197.3	36.64	89	220.7	23.39
Total deaths	205			76			129		
Connecticut									
. Crude rate		7.1	-		5.4	-		8.7	-
. AAMR (1940 Std. mil.)		1.8	0.15		2.2	0.28		1.5	0.16
. AAMR (1970 Std. mil.)		3.4	0.25		3.9	0.45		3.0	0.28
. AAMR (2000 Std. mil.)		5.8	0.41		6.4	0.75		5.3	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.5	-		1.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.93	3	3.1	1.80	0	0.0	0.00
5-9	1	0.5	0.51	0	0.0	0.00	1	1.0	1.04
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.8	1.03	2	2.3	1.63	1	1.2	1.24
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	1	0.4	0.41	0	0.0	0.00	1	0.8	0.83
35-39	5	1.9	0.86	2	1.5	1.09	3	2.3	1.33
40-44	4	1.7	0.85	4	3.4	1.72	0	0.0	0.00
45-49	10	4.7	1.50	3	2.9	1.66	7	6.6	2.48
50-54	7	4.1	1.56	5	6.0	2.69	2	2.3	1.64
55-59	13	10.1	2.81	5	8.0	3.57	8	12.2	4.30
60-64	27	23.9	4.60	14	25.7	6.88	13	22.2	6.15
65-69	39	32.5	5.20	25	45.4	9.07	14	21.6	5.77
70-74	77	69.0	7.87	50	104.0	14.71	27	42.6	8.19
75-79	105	110.4	10.77	58	150.2	19.73	47	83.1	12.13
80-84	204	320.6	22.45	105	454.2	44.32	99	244.4	24.56
85+	576	1046.8	43.62	191	1299.6	94.03	385	954.7	48.66
Total deaths	1,076			468			608		
Connecticut									
. Crude rate		37.2	-		33.3	-		41.0	-
. AAMR (1940 Std. mil.)		10.8	0.42		14.0	0.73		8.8	0.51
. AAMR (1970 Std. mil.)		19.1	0.62		24.8	1.16		15.7	0.72
. AAMR (2000 Std. mil.)		30.9	0.94		40.0	1.89		26.0	1.07
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.0	-		16.3	-		10.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	2	0.8	0.59	0	0.0	0.00	2	1.7	1.18
35-39	4	1.5	0.77	2	1.5	1.09	2	1.5	1.09
40-44	5	2.1	0.95	4	3.4	1.72	1	0.8	0.84
45-49	10	4.7	1.50	5	4.8	2.15	5	4.7	2.09
50-54	11	6.5	1.96	7	8.4	3.19	4	4.6	2.32
55-59	23	17.9	3.73	7	11.2	4.22	16	24.3	6.08
60-64	53	46.9	6.44	23	42.3	8.81	30	51.1	9.34
65-69	118	98.3	9.05	73	132.4	15.50	45	69.4	10.34
70-74	186	166.8	12.23	77	160.1	18.25	109	171.8	16.46
75-79	240	252.3	16.28	112	290.1	27.41	128	226.4	20.01
80-84	238	374.0	24.24	112	484.4	45.77	126	311.0	27.71
85+	277	503.4	30.25	128	870.9	76.98	149	369.5	30.27
Total deaths	1,169			552			617		
Connecticut									
. Crude rate		40.4	-		39.2	-		41.6	-
. AAMR (1940 Std. mil.)		16.1	0.55		18.8	0.87		14.5	0.72
. AAMR (1970 Std. mil.)		24.4	0.75		29.8	1.28		21.6	0.95
. AAMR (2000 Std. mil.)		33.8	0.99		43.2	1.88		29.3	1.20
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		20.9	-		25.8	-		17.6	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	1	0.4	0.41	0	0.0	0.00	1	0.8	0.83
35-39	4	1.5	0.77	2	1.5	1.09	2	1.5	1.09
40-44	2	0.8	0.60	1	0.9	0.86	1	0.8	0.84
45-49	3	1.4	0.82	0	0.0	0.00	3	2.8	1.62
50-54	2	1.2	0.84	1	1.2	1.20	1	1.2	1.16
55-59	3	2.3	1.35	0	0.0	0.00	3	4.6	2.63
60-64	2	1.8	1.25	1	1.8	1.84	1	1.7	1.70
65-69	4	3.3	1.67	1	1.8	1.81	3	4.6	2.67
70-74	6	5.4	2.20	1	2.1	2.08	5	7.9	3.52
75-79	4	4.2	2.10	1	2.6	2.59	3	5.3	3.06
80-84	3	4.7	2.72	1	4.3	4.33	2	4.9	3.49
85+	5	9.1	4.06	2	13.6	9.62	3	7.4	4.30
Total deaths	41			13			28		
Connecticut									
. Crude rate		1.4	-		0.9	-		1.9	-
. AAMR (1940 Std. mil.)		0.9	0.16		-	-		1.1	0.24
. AAMR (1970 Std. mil.)		1.0	0.17		-	-		1.3	0.26
. AAMR (2000 Std. mil.)		1.3	0.20		-	-		1.6	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		1.1	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	7	2.9	1.10	4	3.3	1.65	3	2.5	1.44
35-39	10	3.8	1.22	8	6.2	2.18	2	1.5	1.09
40-44	22	9.3	1.99	18	15.5	3.65	4	3.3	1.67
45-49	31	14.7	2.64	24	23.0	4.70	7	6.6	2.48
50-54	26	15.4	3.01	19	22.9	5.25	7	8.1	3.06
55-59	25	19.5	3.89	18	28.7	6.77	7	10.6	4.02
60-64	22	19.5	4.15	14	25.7	6.88	8	13.6	4.82
65-69	34	28.3	4.86	20	36.3	8.11	14	21.6	5.77
70-74	39	35.0	5.60	22	45.8	9.76	17	26.8	6.50
75-79	30	31.5	5.76	15	38.9	10.03	15	26.5	6.85
80-84	21	33.0	7.20	11	47.6	14.35	10	24.7	7.81
85+	11	20.0	6.03	7	47.6	18.00	4	9.9	4.96
Total deaths	279			181			98		
Connecticut									
. Crude rate		9.7	-		12.9	-		6.6	-
. AAMR (1940 Std. mil.)		6.6	0.43		9.5	0.74		4.0	0.44
. AAMR (1970 Std. mil.)		7.5	0.46		10.8	0.81		4.7	0.50
. AAMR (2000 Std. mil.)		8.9	0.53		12.9	0.97		5.5	0.57
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.5	-		10.7	-		4.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.0	1.04	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.0	0.73	2	2.0	1.45	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	4	1.5	0.77	1	0.8	0.77	3	2.3	1.33
40-44	2	0.8	0.60	2	1.7	1.22	0	0.0	0.00
45-49	4	1.9	0.95	3	2.9	1.66	1	0.9	0.94
50-54	4	2.4	1.18	3	3.6	2.09	1	1.2	1.16
55-59	7	5.4	2.06	1	1.6	1.60	6	9.1	3.72
60-64	11	9.7	2.93	8	14.7	5.20	3	5.1	2.95
65-69	25	20.8	4.17	8	14.5	5.13	17	26.2	6.36
70-74	35	31.4	5.30	18	37.4	8.82	17	26.8	6.50
75-79	48	50.5	7.28	25	64.8	12.95	23	40.7	8.48
80-84	59	92.7	12.07	32	138.4	24.47	27	66.6	12.83
85+	104	189.0	18.53	42	285.8	44.10	62	153.7	19.53
Total deaths	307			147			160		
Connecticut									
. Crude rate		10.6	-		10.4	-		10.8	-
. AAMR (1940 Std. mil.)		4.0	0.28		5.0	0.46		3.4	0.35
. AAMR (1970 Std. mil.)		6.2	0.38		8.0	0.67		5.2	0.46
. AAMR (2000 Std. mil.)		8.9	0.51		11.9	1.00		7.4	0.59
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.3	-		3.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis-related (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.7	1.20	2	2.1	1.47	3	3.3	1.91
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	1	0.6	0.63	0	0.0	0.00	1	1.3	1.30
25-29	3	1.6	0.90	2	2.0	1.45	1	1.0	1.04
30-34	6	2.5	1.02	4	3.3	1.65	2	1.7	1.18
35-39	7	2.7	1.02	2	1.5	1.09	5	3.8	1.72
40-44	10	4.2	1.34	9	7.7	2.58	1	0.8	0.84
45-49	32	15.2	2.68	17	16.3	3.96	15	14.0	3.62
50-54	44	26.0	3.92	25	30.1	6.02	19	22.0	5.05
55-59	28	21.8	4.12	17	27.1	6.58	11	16.7	5.04
60-64	53	46.9	6.44	37	68.0	11.18	16	27.3	6.82
65-69	113	94.2	8.86	63	114.3	14.40	50	77.1	10.90
70-74	186	166.8	12.23	98	203.8	20.59	88	138.7	14.79
75-79	225	236.5	15.77	131	339.3	29.65	94	166.3	17.15
80-84	264	414.9	25.53	146	631.5	52.26	118	291.3	26.81
85+	383	696.1	35.57	162	1102.3	86.60	221	548.0	36.86
Total deaths	1,362			717			645		
Connecticut									
. Crude rate		47.1	-		51.0	-		43.5	-
. AAMR (1940 Std. mil.)		19.2	0.62		25.6	1.06		14.8	0.75
. AAMR (1970 Std. mil.)		28.4	0.81		39.2	1.48		21.7	0.95
. AAMR (2000 Std. mil.)		39.9	1.08		56.4	2.14		30.4	1.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.0	1.04	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.0	0.73	2	2.0	1.45	0	0.0	0.00
30-34	1	0.4	0.41	1	0.8	0.83	0	0.0	0.00
35-39	5	1.9	0.86	1	0.8	0.77	4	3.1	1.54
40-44	2	0.8	0.60	2	1.7	1.22	0	0.0	0.00
45-49	6	2.8	1.16	5	4.8	2.15	1	0.9	0.94
50-54	4	2.4	1.18	3	3.6	2.09	1	1.2	1.16
55-59	11	8.6	2.58	2	3.2	2.26	9	13.7	4.56
60-64	15	13.3	3.43	10	18.4	5.81	5	8.5	3.81
65-69	31	25.8	4.64	11	20.0	6.02	20	30.8	6.89
70-74	42	37.7	5.81	22	45.8	9.76	20	31.5	7.05
75-79	57	59.9	7.94	31	80.3	14.42	26	46.0	9.02
80-84	71	111.6	13.24	39	168.7	27.01	32	79.0	13.96
85+	120	218.1	19.91	50	340.2	48.11	70	173.6	20.75
Total deaths	369			181			188		
Connecticut									
. Crude rate		12.8	-		12.9	-		12.7	-
. AAMR (1940 Std. mil.)		5.0	0.32		6.2	0.52		4.2	0.40
. AAMR (1970 Std. mil.)		7.5	0.42		9.8	0.74		6.3	0.51
. AAMR (2000 Std. mil.)		10.8	0.56		14.6	1.11		8.8	0.65
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.5	-		6.6	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	7.5	2.00	7	7.3	2.75	7	7.7	2.92
5-9	13	6.6	1.82	9	8.9	2.96	4	4.2	2.08
10-14	9	4.9	1.64	5	5.4	2.40	4	4.5	2.23
15-19	34	20.3	3.48	20	23.1	5.17	14	17.3	4.63
20-24	54	34.2	4.65	44	54.1	8.16	10	13.0	4.12
25-29	39	20.2	3.23	27	27.6	5.32	12	12.5	3.62
30-34	55	22.8	3.08	47	38.8	5.67	8	6.7	2.36
35-39	65	25.0	3.10	54	41.6	5.67	11	8.5	2.55
40-44	70	29.7	3.55	60	51.6	6.67	10	8.4	2.64
45-49	48	22.7	3.28	36	34.6	5.76	12	11.2	3.24
50-54	44	26.0	3.92	30	36.1	6.60	14	16.2	4.33
55-59	24	18.7	3.81	17	27.1	6.58	7	10.6	4.02
60-64	33	29.2	5.08	21	38.6	8.42	12	20.5	5.91
65-69	24	20.0	4.08	16	29.0	7.26	8	12.3	4.36
70-74	63	56.5	7.12	33	68.6	11.95	30	47.3	8.63
75-79	69	72.5	8.73	47	121.7	17.76	22	38.9	8.30
80-84	86	135.2	14.57	38	164.4	26.66	48	118.5	17.10
85+	143	259.9	21.73	52	353.8	49.07	91	225.7	23.66
Total deaths	887			563			324		
Connecticut									
. Crude rate		30.7	-		40.0	-		21.8	-
. AAMR (1940 Std. mil.)		22.5	0.89		32.8	1.51		12.7	0.94
. AAMR (1970 Std. mil.)		24.1	0.89		34.5	1.52		14.7	0.97
. AAMR (2000 Std. mil.)		28.7	0.97		41.4	1.77		17.7	1.02
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.5	-		43.2	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.1	1.07	1	1.0	1.04	3	3.3	1.91
5-9	7	3.5	1.34	5	4.9	2.20	2	2.1	1.47
10-14	6	3.3	1.34	5	5.4	2.40	1	1.1	1.12
15-19	27	16.1	3.10	13	15.0	4.17	14	17.3	4.63
20-24	36	22.8	3.80	27	33.2	6.39	9	11.7	3.91
25-29	22	11.4	2.42	13	13.3	3.69	9	9.4	3.13
30-34	17	7.1	1.71	16	13.2	3.31	1	0.8	0.83
35-39	22	8.5	1.81	15	11.6	2.99	7	5.4	2.03
40-44	18	7.6	1.80	14	12.1	3.22	4	3.3	1.67
45-49	19	9.0	2.07	15	14.4	3.72	4	3.7	1.87
50-54	17	10.0	2.43	11	13.3	4.00	6	6.9	2.84
55-59	8	6.2	2.20	6	9.6	3.91	2	3.0	2.15
60-64	18	15.9	3.75	9	16.5	5.51	9	15.3	5.11
65-69	7	5.8	2.20	4	7.3	3.63	3	4.6	2.67
70-74	15	13.5	3.47	8	16.6	5.88	7	11.0	4.17
75-79	15	15.8	4.07	11	28.5	8.59	4	7.1	3.54
80-84	13	20.4	5.67	9	38.9	12.98	4	9.9	4.94
85+	9	16.4	5.45	2	13.6	9.62	7	17.4	6.56
Total deaths	280			184			96		
Connecticut									
. Crude rate		9.7	-		13.1	-		6.5	-
. AAMR (1940 Std. mil.)		9.6	0.63		12.9	1.01		6.5	0.75
. AAMR (1970 Std. mil.)		9.6	0.61		12.7	0.99		6.6	0.74
. AAMR (2000 Std. mil.)		9.7	0.59		13.3	0.98		6.5	0.68
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.3	-		22.4	-		10.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.10
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.16	0	0.0	0.00
20-24	3	1.9	1.10	3	3.7	2.13	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	5	2.1	0.93	4	3.3	1.65	1	0.8	0.83
35-39	3	1.2	0.67	2	1.5	1.09	1	0.8	0.77
40-44	6	2.5	1.04	3	2.6	1.49	3	2.5	1.45
45-49	5	2.4	1.06	5	4.8	2.15	0	0.0	0.00
50-54	3	1.8	1.02	2	2.4	1.70	1	1.2	1.16
55-59	4	3.1	1.56	3	4.8	2.76	1	1.5	1.52
60-64	5	4.4	1.98	5	9.2	4.11	0	0.0	0.00
65-69	9	7.5	2.50	7	12.7	4.80	2	3.1	2.18
70-74	22	19.7	4.21	12	25.0	7.20	10	15.8	4.98
75-79	29	30.5	5.66	20	51.8	11.58	9	15.9	5.31
80-84	34	53.4	9.16	11	47.6	14.35	23	56.8	11.84
85+	82	149.0	16.46	28	190.5	36.00	54	133.9	18.22
Total deaths	212			106			106		
Connecticut									
. Crude rate		7.3	-		7.5	-		7.1	-
. AAMR (1940 Std. mil.)		2.9	0.26		4.2	0.47		1.8	0.24
. AAMR (1970 Std. mil.)		4.2	0.31		5.9	0.59		2.9	0.31
. AAMR (2000 Std. mil.)		6.2	0.43		8.4	0.83		4.7	0.46
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.7	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.1	1.07	3	3.1	1.80	1	1.1	1.10
5-9	1	0.5	0.51	1	1.0	0.99	0	0.0	0.00
10-14	1	0.5	0.55	0	0.0	0.00	1	1.1	1.12
15-19	2	1.2	0.84	2	2.3	1.63	0	0.0	0.00
20-24	3	1.9	1.10	3	3.7	2.13	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	1	0.4	0.41	1	0.8	0.83	0	0.0	0.00
35-39	3	1.2	0.67	2	1.5	1.09	1	0.8	0.77
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.47	1	1.0	0.96	0	0.0	0.00
50-54	2	1.2	0.84	2	2.4	1.70	0	0.0	0.00
55-59	2	1.6	1.10	2	3.2	2.26	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	1	0.8	0.83	0	0.0	0.00	1	1.5	1.54
70-74	1	0.9	0.90	0	0.0	0.00	1	1.6	1.58
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	23			18			5		
Connecticut									
. Crude rate		0.8	-		1.3	-		0.3	-
. AAMR (1940 Std. mil.)		0.9	0.20		1.5	0.36		-	-
. AAMR (1970 Std. mil.)		0.9	0.20		1.4	0.35		-	-
. AAMR (2000 Std. mil.)		0.8	0.18		1.3	0.31		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.5	-		2.3	-		0.4	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.6	0.93	1	1.0	1.04	2	2.2	1.56
5-9	3	1.5	0.88	1	1.0	0.99	2	2.1	1.47
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.6	0.63	1	1.2	1.23	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.8	0.59	2	1.7	1.17	0	0.0	0.00
35-39	2	0.8	0.54	2	1.5	1.09	0	0.0	0.00
40-44	2	0.8	0.60	2	1.7	1.22	0	0.0	0.00
45-49	2	0.9	0.67	2	1.9	1.36	0	0.0	0.00
50-54	3	1.8	1.02	2	2.4	1.70	1	1.2	1.16
55-59	3	2.3	1.35	2	3.2	2.26	1	1.5	1.52
60-64	1	0.9	0.88	0	0.0	0.00	1	1.7	1.70
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	5	4.5	2.01	4	8.3	4.16	1	1.6	1.58
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	4	6.3	3.14	2	8.7	6.12	2	4.9	3.49
85+	2	3.6	2.57	0	0.0	0.00	2	5.0	3.51
Total deaths	33			21			12		
Connecticut									
. Crude rate		1.1	-		1.5	-		0.8	-
. AAMR (1940 Std. mil.)		0.9	0.17		1.2	0.28		-	-
. AAMR (1970 Std. mil.)		1.0	0.19		1.3	0.29		-	-
. AAMR (2000 Std. mil.)		1.1	0.19		1.5	0.32		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.0	-		1.4	-		0.9	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	6	3.3	1.34	5	5.4	2.40	1	1.1	1.12
15-19	9	5.4	1.79	5	5.8	2.58	4	4.9	2.47
20-24	19	12.0	2.76	16	19.7	4.92	3	3.9	2.26
25-29	19	9.8	2.25	18	18.4	4.34	1	1.0	1.04
30-34	33	13.7	2.38	26	21.5	4.21	7	5.8	2.20
35-39	28	10.8	2.04	20	15.4	3.45	8	6.2	2.17
40-44	30	12.7	2.32	27	23.2	4.47	3	2.5	1.45
45-49	18	8.5	2.01	16	15.4	3.84	2	1.9	1.32
50-54	22	13.0	2.77	12	14.5	4.17	10	11.6	3.66
55-59	20	15.6	3.48	17	27.1	6.58	3	4.6	2.63
60-64	11	9.7	2.93	11	20.2	6.10	0	0.0	0.00
65-69	15	12.5	3.23	10	18.1	5.74	5	7.7	3.45
70-74	10	9.0	2.84	9	18.7	6.24	1	1.6	1.58
75-79	7	7.4	2.78	5	13.0	5.79	2	3.5	2.50
80-84	7	11.0	4.16	5	21.6	9.67	2	4.9	3.49
85+	5	9.1	4.06	5	34.0	15.21	0	0.0	0.00
Total deaths	259			207			52		
Connecticut									
. Crude rate		9.0	-		14.7	-		3.5	-
. AAMR (1940 Std. mil.)		8.4	0.55		13.6	0.99		3.3	0.50
. AAMR (1970 Std. mil.)		8.0	0.52		13.1	0.95		3.2	0.47
. AAMR (2000 Std. mil.)		8.8	0.55		14.6	1.02		3.5	0.49
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.8	-		17.9	-		4.0	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.1	1.07	2	2.1	1.47	2	2.2	1.56
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.1	0.77	1	1.1	1.07	1	1.1	1.12
15-19	10	6.0	1.89	8	9.2	3.27	2	2.5	1.75
20-24	9	5.7	1.90	9	11.1	3.69	0	0.0	0.00
25-29	13	6.7	1.86	9	9.2	3.07	4	4.2	2.09
30-34	10	4.1	1.31	8	6.6	2.34	2	1.7	1.18
35-39	7	2.7	1.02	6	4.6	1.89	1	0.8	0.77
40-44	9	3.8	1.27	7	6.0	2.28	2	1.7	1.18
45-49	2	0.9	0.67	0	0.0	0.00	2	1.9	1.32
50-54	11	6.5	1.96	4	4.8	2.41	7	8.1	3.06
55-59	1	0.8	0.78	1	1.6	1.60	0	0.0	0.00
60-64	1	0.9	0.88	0	0.0	0.00	1	1.7	1.70
65-69	3	2.5	1.44	3	5.4	3.14	0	0.0	0.00
70-74	3	2.7	1.55	2	4.2	2.94	1	1.6	1.58
75-79	1	1.1	1.05	0	0.0	0.00	1	1.8	1.77
80-84	1	1.6	1.57	0	0.0	0.00	1	2.5	2.47
85+	1	1.8	1.82	0	0.0	0.00	1	2.5	2.48
Total deaths	88			60			28		
Connecticut									
. Crude rate		3.0	-		4.3	-		1.9	-
. AAMR (1940 Std. mil.)		3.3	0.37		4.7	0.63		1.9	0.38
. AAMR (1970 Std. mil.)		3.1	0.35		4.3	0.58		1.8	0.36
. AAMR (2000 Std. mil.)		3.1	0.33		4.3	0.55		1.9	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.5	-		13.3	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.1	1.07	1	1.0	1.04	3	3.3	1.91
5-9	7	3.5	1.34	5	4.9	2.20	2	2.1	1.47
10-14	6	3.3	1.34	5	5.4	2.40	1	1.1	1.12
15-19	27	16.1	3.10	13	15.0	4.17	14	17.3	4.63
20-24	36	22.8	3.80	27	33.2	6.39	9	11.7	3.91
25-29	21	10.9	2.37	12	12.3	3.54	9	9.4	3.13
30-34	16	6.6	1.66	15	12.4	3.20	1	0.8	0.83
35-39	20	7.7	1.72	13	10.0	2.78	7	5.4	2.03
40-44	18	7.6	1.80	14	12.1	3.22	4	3.3	1.67
45-49	19	9.0	2.07	15	14.4	3.72	4	3.7	1.87
50-54	16	9.4	2.36	10	12.0	3.81	6	6.9	2.84
55-59	8	6.2	2.20	6	9.6	3.91	2	3.0	2.15
60-64	17	15.0	3.65	8	14.7	5.20	9	15.3	5.11
65-69	7	5.8	2.20	4	7.3	3.63	3	4.6	2.67
70-74	15	13.5	3.47	8	16.6	5.88	7	11.0	4.17
75-79	14	14.7	3.93	11	28.5	8.59	3	5.3	3.06
80-84	13	20.4	5.67	9	38.9	12.98	4	9.9	4.94
85+	8	14.5	5.14	2	13.6	9.62	6	14.9	6.07
Total deaths	272			178			94		
Connecticut									
. Crude rate		9.4	-		12.7	-		6.3	-
. AAMR (1940 Std. mil.)		9.4	0.62		12.5	1.00		6.5	0.75
. AAMR (1970 Std. mil.)		9.3	0.61		12.4	0.97		6.5	0.73
. AAMR (2000 Std. mil.)		9.5	0.58		12.9	0.97		6.4	0.67
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.8	-		21.9	-		10.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	0	0.0	0.00	1	1.1	1.10
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	1.9	1.10	3	3.7	2.13	0	0.0	0.00
25-29	1	0.5	0.52	1	1.0	1.02	0	0.0	0.00
30-34	5	2.1	0.93	4	3.3	1.65	1	0.8	0.83
35-39	3	1.2	0.67	2	1.5	1.09	1	0.8	0.77
40-44	6	2.5	1.04	3	2.6	1.49	3	2.5	1.45
45-49	6	2.8	1.16	6	5.8	2.35	0	0.0	0.00
50-54	5	3.0	1.32	3	3.6	2.09	2	2.3	1.64
55-59	3	2.3	1.35	2	3.2	2.26	1	1.5	1.52
60-64	5	4.4	1.98	5	9.2	4.11	0	0.0	0.00
65-69	7	5.8	2.20	6	10.9	4.44	1	1.5	1.54
70-74	19	17.0	3.91	12	25.0	7.20	7	11.0	4.17
75-79	22	23.1	4.93	14	36.3	9.69	8	14.2	5.00
80-84	23	36.1	7.54	7	30.3	11.44	16	39.5	9.87
85+	53	96.3	13.23	16	108.9	27.22	37	91.7	15.08
Total deaths	162			84			78		
Connecticut									
. Crude rate		5.6	-		6.0	-		5.3	-
. AAMR (1940 Std. mil.)		2.5	0.25		3.7	0.45		1.5	0.23
. AAMR (1970 Std. mil.)		3.4	0.29		4.8	0.53		2.3	0.29
. AAMR (2000 Std. mil.)		4.8	0.38		6.4	0.71		3.5	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.6	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.0	1.04	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	5	2.7	1.22	4	4.3	2.14	1	1.1	1.12
15-19	8	4.8	1.69	5	5.8	2.58	3	3.7	2.14
20-24	21	13.3	2.90	20	24.6	5.50	1	1.3	1.30
25-29	19	9.8	2.25	14	14.3	3.83	5	5.2	2.33
30-34	24	10.0	2.03	23	19.0	3.96	1	0.8	0.83
35-39	13	5.0	1.39	11	8.5	2.56	2	1.5	1.09
40-44	18	7.6	1.80	17	14.6	3.55	1	0.8	0.84
45-49	10	4.7	1.50	8	7.7	2.72	2	1.9	1.32
50-54	14	8.3	2.21	7	8.4	3.19	7	8.1	3.06
55-59	6	4.7	1.91	5	8.0	3.57	1	1.5	1.52
60-64	6	5.3	2.17	6	11.0	4.50	0	0.0	0.00
65-69	6	5.0	2.04	6	10.9	4.44	0	0.0	0.00
70-74	8	7.2	2.54	8	16.6	5.88	0	0.0	0.00
75-79	4	4.2	2.10	4	10.4	5.18	0	0.0	0.00
80-84	4	6.3	3.14	4	17.3	8.65	0	0.0	0.00
85+	2	3.6	2.57	2	13.6	9.62	0	0.0	0.00
Total deaths	169			145			24		
Connecticut									
. Crude rate		5.8	-		10.3	-		1.6	-
. AAMR (1940 Std. mil.)		5.9	0.48		10.0	0.88		1.9	0.39
. AAMR (1970 Std. mil.)		5.5	0.45		9.5	0.83		1.7	0.37
. AAMR (2000 Std. mil.)		5.8	0.45		10.2	0.85		1.7	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.8	-		22.3	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	3	1.6	0.95	3	3.2	1.86	0	0.0	0.00
15-19	1	0.6	0.60	0	0.0	0.00	1	1.2	1.24
20-24	11	7.0	2.10	10	12.3	3.89	1	1.3	1.30
25-29	7	3.6	1.37	6	6.1	2.51	1	1.0	1.04
30-34	15	6.2	1.61	15	12.4	3.20	0	0.0	0.00
35-39	8	3.1	1.09	6	4.6	1.89	2	1.5	1.09
40-44	12	5.1	1.47	11	9.5	2.85	1	0.8	0.84
45-49	8	3.8	1.34	8	7.7	2.72	0	0.0	0.00
50-54	7	4.1	1.56	4	4.8	2.41	3	3.5	2.01
55-59	5	3.9	1.74	4	6.4	3.19	1	1.5	1.52
60-64	6	5.3	2.17	6	11.0	4.50	0	0.0	0.00
65-69	6	5.0	2.04	6	10.9	4.44	0	0.0	0.00
70-74	8	7.2	2.54	8	16.6	5.88	0	0.0	0.00
75-79	4	4.2	2.10	4	10.4	5.18	0	0.0	0.00
80-84	4	6.3	3.14	4	17.3	8.65	0	0.0	0.00
85+	2	3.6	2.57	2	13.6	9.62	0	0.0	0.00
Total deaths	107			97			10		
Connecticut									
. Crude rate		3.7	-		6.9	-		0.7	-
. AAMR (1940 Std. mil.)		3.4	0.35		6.1	0.66		-	-
. AAMR (1970 Std. mil.)		3.3	0.33		6.0	0.64		-	-
. AAMR (2000 Std. mil.)		3.6	0.35		6.8	0.69		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		11.3	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.0	1.04	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.1	0.77	1	1.1	1.07	1	1.1	1.12
15-19	7	4.2	1.58	5	5.8	2.58	2	2.5	1.75
20-24	9	5.7	1.90	9	11.1	3.69	0	0.0	0.00
25-29	12	6.2	1.79	8	8.2	2.89	4	4.2	2.09
30-34	8	3.3	1.17	7	5.8	2.19	1	0.8	0.83
35-39	5	1.9	0.86	5	3.9	1.72	0	0.0	0.00
40-44	3	1.3	0.73	3	2.6	1.49	0	0.0	0.00
45-49	2	0.9	0.67	0	0.0	0.00	2	1.9	1.32
50-54	7	4.1	1.56	3	3.6	2.09	4	4.6	2.32
55-59	1	0.8	0.78	1	1.6	1.60	0	0.0	0.00
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	57			43			14		
Connecticut									
. Crude rate		2.0	-		3.1	-		0.9	-
. AAMR (1940 Std. mil.)		2.4	0.32		3.6	0.56		-	-
. AAMR (1970 Std. mil.)		2.1	0.29		3.1	0.50		-	-
. AAMR (2000 Std. mil.)		2.1	0.27		3.1	0.47		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.9	-		9.8	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	1.0	0.73	2	2.0	1.45	0	0.0	0.00
30-34	5	2.1	0.93	2	1.7	1.17	3	2.5	1.44
35-39	9	3.5	1.15	7	5.4	2.04	2	1.5	1.09
40-44	16	6.8	1.70	13	11.2	3.10	3	2.5	1.45
45-49	30	14.2	2.60	25	24.0	4.80	5	4.7	2.09
50-54	17	10.0	2.43	13	15.7	4.34	4	4.6	2.32
55-59	10	7.8	2.46	6	9.6	3.91	4	6.1	3.04
60-64	19	16.8	3.86	13	23.9	6.63	6	10.2	4.18
65-69	18	15.0	3.54	8	14.5	5.13	10	15.4	4.87
70-74	13	11.7	3.23	10	20.8	6.58	3	4.7	2.73
75-79	13	13.7	3.79	8	20.7	7.33	5	8.8	3.96
80-84	3	4.7	2.72	3	13.0	7.49	0	0.0	0.00
85+	4	7.3	3.63	2	13.6	9.62	2	5.0	3.51
Total deaths	159			112			47		
Connecticut									
. Crude rate		5.5	-		8.0	-		3.2	-
. AAMR (1940 Std. mil.)		4.2	0.35		6.3	0.61		2.3	0.36
. AAMR (1970 Std. mil.)		4.5	0.36		6.6	0.64		2.5	0.38
. AAMR (2000 Std. mil.)		5.2	0.41		7.9	0.75		2.8	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		10.1	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1996

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.54	1	1.0	1.04	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	3	1.8	1.03	2	2.3	1.63	1	1.2	1.24
20-24	10	6.3	2.00	8	9.8	3.48	2	2.6	1.84
25-29	15	7.8	2.00	13	13.3	3.69	2	2.1	1.48
30-34	35	14.5	2.45	24	19.8	4.05	11	9.2	2.76
35-39	40	15.4	2.43	30	23.1	4.22	10	7.7	2.43
40-44	36	15.3	2.54	30	25.8	4.71	6	5.0	2.05
45-49	21	10.0	2.17	13	12.5	3.46	8	7.5	2.65
50-54	5	3.0	1.32	3	3.6	2.09	2	2.3	1.64
55-59	5	3.9	1.74	2	3.2	2.26	3	4.6	2.63
60-64	2	1.8	1.25	2	3.7	2.60	0	0.0	0.00
65-69	1	0.8	0.83	0	0.0	0.00	1	1.5	1.54
70-74	1	0.9	0.90	0	0.0	0.00	1	1.6	1.58
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	1	1.6	1.57	1	4.3	4.33	0	0.0	0.00
85+	2	3.6	2.57	0	0.0	0.00	2	5.0	3.51
Total deaths	178			129			49		
Connecticut									
. Crude rate		6.2	-		9.2	-		3.3	-
. AAMR (1940 Std. mil.)		5.8	0.45		8.5	0.77		3.0	0.45
. AAMR (1970 Std. mil.)		4.9	0.39		7.2	0.66		2.6	0.40
. AAMR (2000 Std. mil.)		5.9	0.45		8.6	0.76		3.2	0.46
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.3	-		7.3	-		3.1	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is higher