

the next stage, and then seek a host. Other residents may pick up lone star ticks while visiting or vacationing on Long Island or areas further south.

Background on the Lone Star Tick

The lone star tick is the most common human biting tick in the southeastern United States with roughly 90-95% or more of the ticks either collected from vegetation or reported attached to humans being *A. americanum*. This tick is widely distributed through the southeastern United States as far west as mid-Texas and north to southern parts of Iowa, Illinois, Indiana, Ohio, Pennsylvania, and Wisconsin. Along the Atlantic coast, its northern range extends to the middle and southern half of New Jersey and Long Island, New York. It is also abundant on Prudence Island, RI and recently appears to have become established on parts of Cape Cod, MA. Their northern range may be increasing due, in part, to the milder winters the northeast has been experiencing over the past few years. Drs. Stafford and Scott Williams at CAES have found that the lone star tick can overwinter in Connecticut.

Lone star ticks are reddish brown in color. The tick gets its name from a conspicuous spot on the back of female ticks. Male ticks have faint white markings at the edge of the body. All stages are active during the summer months, but adult tick activity peaks in the spring and early summer, while nymphs are active from April throughout the summer. Larval ticks are most abundant in the late summer and early fall. Female ticks can lay 1,000 to 8,000 eggs with an average of around 3,000 eggs, so populations can build to tremendous numbers under suitable conditions. In some areas in the southeast where this tick has long been established, 80-100% of white-tailed deer can be infested with tick burdens exceeding 300 ticks per deer. Heavy infestations have been known to kill fawns and young animals.

Long considered primarily a noxious pest, the lone star tick is associated with a number of human and animal diseases such as ehrlichiosis, Southern tick-associated rash illness (STARI), spotted fever rickettsiosis, tularemia, Heartland virus, theileriosis in deer (related to human babesiosis), and more recently, red meat allergy. Bites from lone star ticks may produce or generate a hypersensitivity allergic response that causes a food allergy to a red meat protein called alpha-gal that is found in most mammals, but not humans or other primates. Lone star ticks do NOT transmit the pathogens that cause Lyme disease, babesiosis or anaplasmosis, which are transmitted by the blacklegged tick, *Ixodes scapularis*.

Protective Measures for People

Personal protection measures against the lone star tick are the same as that for the blacklegged tick. The use of a repellent or pesticide, correctly applied to clothing and on gear following specific product label instructions, is considered the best tick bite prevention measure (see CAES fact sheet *The Prevention of Tick Bite and Tick-Borne Disease: Tick Checks and the Use of Insect Repellents*; www.ct.gov/caes). Wearing light colored clothing, inspecting clothing, gear, and pets, conducting a full body tick check, and showering after being outdoors are all suggested steps toward preventing tick bites. The primary disease associated with the lone star tick will be ehrlichiosis. Previous studies have indicated about 8-13% of these ticks are infected with *Ehrlichia chaffeensis* and cases may occur after tick bite. Symptoms are similar to anaplasmosis and antibiotic treatment is the same. An article on the discovery by DEEP wildlife biologist Andrew LaBonte will appear in the September/October issue of Connecticut Wildlife. Questions can be directed to Dr. Kirby Stafford at CAES (203-974-8485 or Kirby.Stafford@ct.gov).

Captions for attached photos:

Photo 1: Female lone star tick (photo by J. Gathany, CDC)

Photo 2: Lone star ticks on white-tailed deer in South Norwalk, CT (J. Nivolo)

Photo 3: Close up of ticks on the white-tailed deer in South Norwalk, CT (K. Stafford)

Photo 4: Female lone star ticks on sock (K. Stafford)







