



CAES

The Connecticut Agricultural Experiment Station
Putting Science to Work for Society since 1875

Founded in 1875
Putting science to work for society

Dr. Yonghao Li
Department of Plant Pathology and Ecology
The Connecticut Agricultural Experiment Station
123 Huntington Street, P. O. Box 1106
New Haven, CT 06504

Phone: (203) 974-8601
Fax: (203) 974-8502
Email: Yonghao.Li@ct.gov
Website: www.portal.ct.gov/caes

BEECH LEAF DISEASE

Beech leaf disease was first discovered in Ohio in 2012. It has since been found in Pennsylvania, New York, and Ontario, Canada in North America. In Connecticut, this disease was first detected in August 2019 (Figure 1). The disease has been observed mainly in forests, but also in landscaped areas. Beech leaf disease causes premature leaf drop and thin canopies, and also makes the trees more susceptible to other pests.




Figure 1. A beech sapling affected by beech leaf disease (arrow).

SYMPTOMS AND DIAGNOSTICS

Dark green striping between leaf veins is a characteristic symptom of this disease, which is especially noticeable when viewing upward into the canopy (Figure 2) or viewing against light (Figure 3). The symptom appears when leaves form in the spring. The initial symptoms are darkening and wrinkling of small portion of leaf tissues between veins (Figures 3 and 4). As disease progresses, the infected leaf tissues turn yellow, slightly raised, crinkly, and leathery (Figure 5). Heavily infected leaves are curled downward, shrunken, and are prematurely defoliated (Figure 6). Sapling and young trees are more susceptible to the disease and can die within three years after symptoms are observed, which can reduce the proportion of American beech in the affected forest areas. Symptoms of other



Figure 2. Dark green striping between veins on beech leaves.

pests, such as beech blight aphid, European beech scale, eriophyid mites, and anthracnose, can resemble beech leaf disease, which stresses the need for a laboratory examination.

DISEASE DEVELOPMENT

A foliar nematode species, *Litylenchus crenatae*, that was first described in Japan is associated with beech leaf disease and pathogenic to American beech (*Fagus grandifolia*), European beech (*F. sylvatica*), and oriental beech (*F. orientalis*). However, the origin of the pathogen and its distribution is unclear. Emerging leaves can be infected by the pathogen in the spring. But, survival and dissemination of the pathogens are unknown.

DISEASE MANAGEMENT

Because little is known about the biology of the pathogen and epidemiology of the disease, no effective control or eradication measures have been developed. In general, the spread of invasive species can be prevented by restricting the movement of plant materials and monitoring trees closely for signs and symptoms. Beech leaf disease has spread very quickly eastward in the United States. Management efforts for the disease should focus on preventing the introduction of this invasive pathogen. Quarantines and regulations can be used to prevent further spread of beech leaf disease.

August 2019




Figure 3. Darkening of leaf tissues between veins.



Figure 5. Yellow striping between veins on the upper surface of beech leaves.



Figure 3. Wrinkling of leaf tissues between veins.



Figure 6. Curling and browning of leaves.