

**Connecticut Department of Public Health HIV Surveillance Program
Recently Diagnosed HIV cases by Risk, Sex, Race, and Age Group, New Haven, 2015 - 2019**

	Total		Mode of Transmission											
			MSM		PWID		Heterosexual contact		Perinatal		Presumed Heterosexual contact		Unknown	
	N	% of total	N	% of row total	N	% of row total	N	% of row total	N	% of row total	N	% of row total	N	% of row total
Total	147	100.0	93	63.3	4	2.7	35	23.8	1	0.7	4	2.7	10	6.8
Sex														
Male	121	82.3	93	76.9	4	3.3	15	12.4	0	0	0	0	9	7.4
Female	26	17.7	0	0	0	0	20	76.9	1	3.8	4	15.4	1	3.8
Race/ethnicity														
Black/African American	88	59.9	44	50.0	2	2.3	29	33.0	1	1.1	4	4.5	8	9.1
Hispanic/Latino	40	27.2	33	82.5	0	0	5	12.5	0	0	0	0	2	5.0
White	16	10.9	14	87.5	1	6.3	1	6.3	0	0	0	0	0	0
Other races	3	2.0	2	66.7	1	33.3	0	0	0	0	0	0	0	0
Age at diagnosis														
<20	13	8.8	12	92.3	0	0	0	0	1	7.7	0	0	0	0
20-29	59	40.1	46	78.0	1	1.7	10	16.9	0	0	1	1.7	1	1.7
30-39	31	21.1	19	61.3	0	0	9	29.0	0	0	1	3.2	2	6.5
40-49	20	13.6	7	35.0	1	5.0	6	30.0	0	0	1	5.0	5	25.0
50+	24	16.3	9	37.5	2	8.3	10	41.7	0	0	1	4.2	2	8.3

This table presents HIV Infection cases using the year the person was first diagnosed.

The term HIV Infection is used to refer to three categories of diagnoses collectively:

- 1 - persons with a diagnosis of HIV infection (not AIDS)
- 2 - a diagnosis of HIV infection and a later diagnosis of AIDS
- 3 - concurrent diagnoses of HIV and AIDS.

Source: HIV Surveillance Registry for cases reported through December 23, 2020