

Viral suppression ≤ 6 months of diagnosis of HIV infection among persons aged ≥13 years, Connecticut, 2019

Characteristics	Total Diagnosis No.	VL<200 copies ml ^a	
		No.	%
Sex at Birth			
Male	168	130	77.4
Female	51	38	74.5
Gender			
Men	168	130	77.4
Women	50	37	74.0
Transgender persons	1	1	100.0
Age at Diagnosis (yr)			
13-24	48	37	77.1
25-34	74	56	75.7
35-44	25	20	80.0
45-54	38	27	71.1
55+	34	28	82.4
Race/ethnicity			
Asian	7	7	100.0
Black/African American	102	76	74.5
Hispanic/Latino	51	41	80.4
White	58	44	75.9
Multiple races	1	0	0.0
Transmission Category^e			
Male-to-male sexual contact	119	94	79.0
Injection drug use			
Male	6	2	33.3
Female	8	5	62.5
Male-to-male sexual contact and injection drug use	2	2	100.0
Heterosexual contact ^f			
Male	17	14	82.4
Female	28	25	89.3
Other ^g			
Male	24	18	75.0
Female	15	8	53.3
Total	219	168	76.7

^a A viral load test result of < 200 copies/mL indicates HIV viral suppression. Viral load test results are within 6 months (≤182 days) of diagnosis of HIV infection during the specified year.

^b "Transgender male-to-female" includes individuals who were assigned "male" sex at birth but have ever identified as "female" gender. "Transgender female-to-male" includes individuals who were assigned "female" sex at birth but have ever identified as "male" gender.

^e Transmission category data presented based on sex at birth and include transgender persons.

^f Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^g Other transmission category includes persons whose infection was attributed to hemophilia, blood transfusion, perinatal exposure or whose risk factor was not reported or not identified. Data not displayed because the numbers were too small to be meaningful.