

Receipt of HIV medical care among people living with HIV aged ≥13 years, Connecticut, 2019

Characteristics	PLWH	≥1 care visit ^a		≥2 care visits ^a	
	No.	No.	%	No.	%
Gender					
Men	6,888	5,491	79.7	3,975	57.7
Women	3,509	2,870	81.8	2,117	60.3
Transwomen ^b	71	63	88.7	51	71.8
Transmen ^b	8	7	87.5	6	75.0
Additional Gender Identity ^c	3	3	100.0	3	100.0
Age in 2018					
13-24	230	182	79.1	128	55.7
25-34	1,070	837	78.2	558	52.1
35-44	1,525	1,191	78.1	812	53.2
45-54	3,027	2,455	81.1	1,802	59.5
55+	4,627	3,769	81.5	2,852	61.6
Race/ethnicity					
American Indian/Alaska Native	13	9	69.2	7	53.8
Asian	106	84	79.2	49	46.2
Black/African American	3,548	2,862	80.7	2,092	59.0
Hispanic/Latino	3,552	2,790	78.5	2,136	60.1
Native Hawaiian/Other Pacific Islander	7	6	85.7	3	42.9
White	3,116	2,563	82.3	1,785	57.3
Multiple races	137	120	87.6	80	58.4
Transmission Category^e					
Male-to-male sexual contact	3,295	2,744	83.3	1,927	58.5
Injection drug use					
Male	1,723	1,323	76.8	1,027	59.6
Female	989	806	81.5	621	62.8
Male-to-male sexual contact and injection drug use	247	213	86.2	152	61.5
Heterosexual contact ^f					
Male	964	771	80.0	565	58.6
Female	1,938	1,642	84.7	1,195	61.7
Other ^g					
Male	733	506	69.0	358	48.8
Female	590	429	72.7	307	52.0
Total	10,479	8,434	80.5	6,152	58.7

Note: includes persons aged ≥13 years and diagnosed with HIV through 2018 and living as residents of Connecticut through 2019.

^a Receipt of care any care is defined as at least one care visit during the calendar year. Retention in care is defined as two or more visits at least 3 months (≤91 days) apart in the calendar year.

^b Transwomen includes individuals who were assigned male sex at birth but have ever identified as female gender. Transmen includes individuals who were assigned female sex at birth but have ever identified as male gender.

^c Additional gender identity examples include "bigender," "gender queer," and "two-spirit."

^d Hispanics/Latinos can be of any race.

^e Transmission category data presented based on sex at birth and include transgender persons.

^f Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^g Other transmission category includes persons whose infection was attributed to hemophilia, blood transfusion, perinatal exposure or whose risk factor was not reported or not identified. Data not displayed because the numbers were too small to be meaningful.