

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.23	1	0.5	0.47	0	0.0	0.00
30-34	6	1.1	0.46	3	1.1	0.66	3	1.1	0.65
35-39	13	2.2	0.61	9	3.1	1.03	4	1.4	0.68
40-44	47	8.6	1.25	25	9.3	1.87	22	7.9	1.68
45-49	95	20.7	2.12	53	23.6	3.24	42	17.9	2.76
50-54	180	44.6	3.33	88	44.8	4.78	92	44.4	4.63
55-59	262	86.2	5.33	149	101.3	8.30	113	72.0	6.78
60-64	403	165.3	8.23	226	193.6	12.88	177	139.2	10.47
65-69	544	220.0	9.43	330	290.2	15.98	214	160.2	10.95
70-74	665	287.9	11.16	378	378.7	19.48	287	218.8	12.92
75-79	666	329.8	12.78	358	434.6	22.97	308	257.7	14.68
80-84	510	369.7	16.37	263	515.7	31.80	247	284.1	18.08
85+	331	274.6	15.09	169	506.2	38.94	162	185.8	14.60
Total deaths	3,723			2,052			1,671		
Connecticut									
. Crude rate		56.9	-		64.6	-		49.6	-
. AAMR (1940 Std. mil.)		33.0	0.60		41.5	0.98		26.5	0.74
. AAMR (1970 Std. mil.)		43.3	0.74		55.7	1.25		34.3	0.90
. AAMR (2000 Std. mil.)		52.3	0.86		68.8	1.53		41.2	1.02
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		37.4	-		50.4	-		27.0	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	6	1.1	0.46	-	-	-	6	2.2	0.92
35-39	20	3.4	0.76	-	-	-	20	6.8	1.51
40-44	49	9.0	1.28	-	-	-	49	17.5	2.51
45-49	70	15.2	1.82	-	-	-	70	29.8	3.56
50-54	113	28.0	2.64	-	-	-	113	54.6	5.13
55-59	81	26.7	2.96	-	-	-	81	51.6	5.74
60-64	92	37.7	3.93	-	-	-	92	72.4	7.55
65-69	122	49.3	4.47	-	-	-	122	91.3	8.27
70-74	172	74.5	5.68	-	-	-	172	131.1	10.00
75-79	130	64.4	5.65	-	-	-	130	108.8	9.54
80-84	139	100.8	8.55	-	-	-	139	159.9	13.56
85+	198	164.2	11.67	-	-	-	198	227.1	16.14
Total deaths	1,192			-			1,192		
Connecticut									
. Crude rate		-	-		-	-		35.4	-
. AAMR (1940 Std. mil.)		-	-		-	-		19.6	0.65
. AAMR (1970 Std. mil.)		-	-		-	-		24.1	0.75
. AAMR (2000 Std. mil.)		-	-		-	-		29.6	0.87
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		19.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.2	0.18	-	-	-	1	0.4	0.36
45-49	3	0.7	0.38	-	-	-	3	1.3	0.74
50-54	1	0.2	0.25	-	-	-	1	0.5	0.48
55-59	5	1.6	0.74	-	-	-	5	3.2	1.43
60-64	6	2.5	1.00	-	-	-	6	4.7	1.93
65-69	7	2.8	1.07	-	-	-	7	5.2	1.98
70-74	11	4.8	1.44	-	-	-	11	8.4	2.53
75-79	17	8.4	2.04	-	-	-	17	14.2	3.45
80-84	15	10.9	2.81	-	-	-	15	17.3	4.45
85+	18	14.9	3.52	-	-	-	18	20.6	4.87
Total deaths	84			-			84		
Connecticut									
. Crude rate		-	-		-	-		2.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.1	0.14
. AAMR (1970 Std. mil.)		-	-		-	-		1.5	0.18
. AAMR (2000 Std. mil.)		-	-		-	-		1.9	0.22
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.2	0.23	-	-	-	1	0.5	0.46
30-34	2	0.4	0.27	-	-	-	2	0.7	0.53
35-39	2	0.3	0.24	-	-	-	2	0.7	0.48
40-44	7	1.3	0.48	-	-	-	7	2.5	0.95
45-49	11	2.4	0.72	-	-	-	11	4.7	1.41
50-54	34	8.4	1.45	-	-	-	34	16.4	2.82
55-59	24	7.9	1.61	-	-	-	24	15.3	3.12
60-64	33	13.5	2.36	-	-	-	33	26.0	4.52
65-69	49	19.8	2.83	-	-	-	49	36.7	5.24
70-74	60	26.0	3.35	-	-	-	60	45.7	5.91
75-79	50	24.8	3.50	-	-	-	50	41.8	5.92
80-84	44	31.9	4.81	-	-	-	44	50.6	7.63
85+	58	48.1	6.32	-	-	-	58	66.5	8.74
Total deaths	375			-			375		
Connecticut									
. Crude rate		-	-		-	-		11.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.0	0.36
. AAMR (1970 Std. mil.)		-	-		-	-		7.6	0.42
. AAMR (2000 Std. mil.)		-	-		-	-		9.2	0.48
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.2	0.23	0	0.0	0.00	1	0.5	0.46
30-34	6	1.1	0.46	4	1.5	0.76	2	0.7	0.53
35-39	13	2.2	0.61	9	3.1	1.03	4	1.4	0.68
40-44	14	2.6	0.68	9	3.4	1.12	5	1.8	0.80
45-49	29	6.3	1.17	16	7.1	1.78	13	5.5	1.54
50-54	49	12.2	1.74	28	14.3	2.70	21	10.1	2.21
55-59	53	17.4	2.40	30	20.4	3.73	23	14.7	3.06
60-64	103	42.2	4.16	55	47.1	6.35	48	37.8	5.45
65-69	139	56.2	4.77	79	69.5	7.82	60	44.9	5.80
70-74	171	74.0	5.66	88	88.2	9.40	83	63.3	6.95
75-79	197	97.6	6.95	101	122.6	12.20	96	80.3	8.20
80-84	204	147.9	10.35	90	176.5	18.60	114	131.1	12.28
85+	301	249.7	14.39	99	296.5	29.80	202	231.7	16.30
Total deaths	1,280			608			672		
Connecticut									
. Crude rate		19.6	-		19.1	-		20.0	-
. AAMR (1940 Std. mil.)		9.5	0.31		11.4	0.50		8.1	0.39
. AAMR (1970 Std. mil.)		13.3	0.39		15.9	0.66		11.4	0.49
. AAMR (2000 Std. mil.)		17.6	0.49		21.1	0.87		15.2	0.60
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		14.8	-		12.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.2	0.17	1	0.3	0.34	0	0.0	0.00
40-44	2	0.4	0.26	1	0.4	0.37	1	0.4	0.36
45-49	2	0.4	0.31	1	0.4	0.44	1	0.4	0.43
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	2	0.7	0.47	1	0.7	0.68	1	0.6	0.64
60-64	1	0.4	0.41	1	0.9	0.86	0	0.0	0.00
65-69	1	0.4	0.40	1	0.9	0.88	0	0.0	0.00
70-74	1	0.4	0.43	0	0.0	0.00	1	0.8	0.76
75-79	2	1.0	0.70	1	1.2	1.21	1	0.8	0.84
80-84	4	2.9	1.45	3	5.9	3.40	1	1.2	1.15
85+	4	3.3	1.66	0	0.0	0.00	4	4.6	2.29
Total deaths	20			10			10		
Connecticut									
. Crude rate		0.3	-		0.3	-		0.3	-
. AAMR (1940 Std. mil.)		0.2	0.04		-	-		-	-
. AAMR (1970 Std. mil.)		0.2	0.05		-	-		-	-
. AAMR (2000 Std. mil.)		0.3	0.06		-	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.5	-		0.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.5	0.35	2	1.0	0.68	0	0.0	0.00
20-24	2	0.6	0.39	2	1.1	0.77	0	0.0	0.00
25-29	12	2.8	0.81	8	3.7	1.32	4	1.9	0.93
30-34	21	3.9	0.86	15	5.7	1.47	6	2.2	0.92
35-39	50	8.5	1.21	38	13.1	2.12	12	4.1	1.17
40-44	126	23.0	2.05	100	37.3	3.73	26	9.3	1.83
45-49	209	45.5	3.15	165	73.4	5.72	44	18.7	2.83
50-54	340	84.3	4.57	265	135.0	8.30	75	36.2	4.18
55-59	490	161.2	7.28	346	235.3	12.65	144	91.8	7.65
60-64	615	252.2	10.17	438	375.3	17.93	177	139.2	10.47
65-69	981	396.7	12.66	647	569.1	22.37	334	250.0	13.68
70-74	1,366	591.4	16.00	844	845.5	29.10	522	398.0	17.42
75-79	2,029	1004.9	22.31	1,129	1370.4	40.79	900	752.9	25.10
80-84	2,466	1787.6	36.00	1,200	2352.9	67.92	1,266	1456.1	40.92
85+	5,120	4246.8	59.35	1,476	4421.1	115.08	3,644	4180.1	69.25
Total deaths	13,829			6,675			7,154		
Connecticut									
. Crude rate		211.3	-		210.0	-		212.6	-
. AAMR (1940 Std. mil.)		83.2	0.86		114.9	1.55		57.2	0.91
. AAMR (1970 Std. mil.)		128.4	1.16		170.3	2.12		95.4	1.26
. AAMR (2000 Std. mil.)		186.5	1.59		238.5	2.96		147.2	1.77
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		100.4	-		137.4	-		70.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.23	1	0.5	0.46	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	0.9	0.47	2	0.9	0.66	2	0.9	0.66
30-34	7	1.3	0.50	5	1.9	0.85	2	0.7	0.53
35-39	21	3.6	0.78	12	4.1	1.19	9	3.0	1.02
40-44	45	8.2	1.23	37	13.8	2.27	8	2.9	1.01
45-49	58	12.6	1.66	40	17.8	2.81	18	7.7	1.81
50-54	55	13.6	1.84	41	20.9	3.26	14	6.8	1.81
55-59	53	17.4	2.40	37	25.2	4.14	16	10.2	2.55
60-64	63	25.8	3.26	45	38.6	5.75	18	14.2	3.34
65-69	67	27.1	3.31	40	35.2	5.56	27	20.2	3.89
70-74	69	29.9	3.60	41	41.1	6.41	28	21.3	4.03
75-79	66	32.7	4.02	21	25.5	5.56	45	37.6	5.61
80-84	50	36.2	5.13	23	45.1	9.40	27	31.1	5.98
85+	21	17.4	3.80	9	27.0	8.99	12	13.8	3.97
Total deaths	580			354			226		
Connecticut									
. Crude rate		8.9	-		11.1	-		6.7	-
. AAMR (1940 Std. mil.)		6.3	0.28		8.7	0.48		4.0	0.30
. AAMR (1970 Std. mil.)		7.2	0.31		9.8	0.53		4.8	0.34
. AAMR (2000 Std. mil.)		8.4	0.35		11.4	0.61		5.8	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.3	-		10.5	-		4.5	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	31	7.2	1.30	22	10.0	2.14	9	4.3	1.44
5-9	16	3.5	0.86	11	4.6	1.40	5	2.2	0.99
10-14	27	6.1	1.18	14	6.2	1.66	13	6.0	1.67
15-19	102	25.2	2.49	69	33.0	3.98	33	16.8	2.93
20-24	115	31.9	2.97	88	47.8	5.09	27	15.3	2.94
25-29	126	29.3	2.61	102	47.6	4.71	24	11.1	2.27
30-34	174	32.7	2.48	131	49.6	4.33	43	16.1	2.45
35-39	187	31.9	2.33	140	48.1	4.06	47	15.9	2.32
40-44	158	28.9	2.30	120	44.8	4.09	38	13.6	2.21
45-49	137	29.8	2.55	100	44.5	4.45	37	15.8	2.59
50-54	75	18.6	2.15	55	28.0	3.78	20	9.7	2.16
55-59	60	19.7	2.55	34	23.1	3.97	26	16.6	3.25
60-64	67	27.5	3.36	41	35.1	5.49	26	20.5	4.01
65-69	98	39.6	4.00	64	56.3	7.04	34	25.4	4.36
70-74	115	49.8	4.64	63	63.1	7.95	52	39.6	5.50
75-79	157	77.8	6.21	83	100.7	11.06	74	61.9	7.20
80-84	201	145.7	10.28	96	188.2	19.21	105	120.8	11.79
85+	312	258.8	14.65	134	401.4	34.67	178	204.2	15.30
Total deaths	2,158			1,367			791		
Connecticut									
. Crude rate		33.0	-		43.0	-		23.5	-
. AAMR (1940 Std. mil.)		25.2	0.62		36.2	1.06		14.7	0.66
. AAMR (1970 Std. mil.)		26.4	0.61		37.5	1.06		16.5	0.67
. AAMR (2000 Std. mil.)		31.3	0.68		44.9	1.23		19.9	0.73
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.1	-		42.8	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	1.2	0.52	2	0.9	0.64	3	1.4	0.83
5-9	10	2.2	0.68	8	3.4	1.19	2	0.9	0.63
10-14	15	3.4	0.88	5	2.2	0.99	10	4.6	1.46
15-19	76	18.8	2.15	49	23.5	3.35	27	13.8	2.65
20-24	72	20.0	2.35	54	29.3	3.99	18	10.2	2.40
25-29	66	15.4	1.89	55	25.7	3.46	11	5.1	1.54
30-34	69	13.0	1.56	48	18.2	2.62	21	7.8	1.71
35-39	55	9.4	1.26	32	11.0	1.94	23	7.8	1.62
40-44	42	7.7	1.18	27	10.1	1.94	15	5.4	1.39
45-49	46	10.0	1.48	31	13.8	2.48	15	6.4	1.65
50-54	32	7.9	1.40	23	11.7	2.44	9	4.3	1.45
55-59	17	5.6	1.36	6	4.1	1.67	11	7.0	2.11
60-64	22	9.0	1.92	14	12.0	3.21	8	6.3	2.22
65-69	31	12.5	2.25	19	16.7	3.83	12	9.0	2.59
70-74	35	15.2	2.56	17	17.0	4.13	18	13.7	3.23
75-79	36	17.8	2.97	17	20.6	5.00	19	15.9	3.65
80-84	35	25.4	4.29	16	31.4	7.84	19	21.9	5.01
85+	26	21.6	4.23	11	32.9	9.93	15	17.2	4.44
Total deaths	690			434			256		
Connecticut									
. Crude rate		10.5	-		13.7	-		7.6	-
. AAMR (1940 Std. mil.)		10.3	0.43		13.8	0.70		6.8	0.48
. AAMR (1970 Std. mil.)		10.0	0.41		13.3	0.67		6.9	0.47
. AAMR (2000 Std. mil.)		10.5	0.40		13.9	0.67		7.3	0.46
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.0	-		21.7	-		10.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									
. The U.S. AAMR (1940 Std. mil.):									
. THE U.S. HP 2000 target:									

Comparison of the CT Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.2	0.22	1	0.4	0.42	0	0.0	0.00
10-14	1	0.2	0.23	1	0.4	0.44	0	0.0	0.00
15-19	2	0.5	0.35	1	0.5	0.48	1	0.5	0.51
20-24	4	1.1	0.55	4	2.2	1.09	0	0.0	0.00
25-29	2	0.5	0.33	2	0.9	0.66	0	0.0	0.00
30-34	7	1.3	0.50	6	2.3	0.93	1	0.4	0.37
35-39	3	0.5	0.30	3	1.0	0.59	0	0.0	0.00
40-44	5	0.9	0.41	5	1.9	0.83	0	0.0	0.00
45-49	7	1.5	0.58	4	1.8	0.89	3	1.3	0.74
50-54	9	2.2	0.74	7	3.6	1.35	2	1.0	0.68
55-59	6	2.0	0.81	5	3.4	1.52	1	0.6	0.64
60-64	8	3.3	1.16	6	5.1	2.10	2	1.6	1.11
65-69	26	10.5	2.06	18	15.8	3.73	8	6.0	2.12
70-74	25	10.8	2.16	15	15.0	3.88	10	7.6	2.41
75-79	52	25.8	3.57	32	38.8	6.87	20	16.7	3.74
80-84	81	58.7	6.52	40	78.4	12.40	41	47.2	7.36
85+	162	134.4	10.56	65	194.7	24.15	97	111.3	11.30
Total deaths	401			215			186		
Connecticut									
. Crude rate		6.1	-		6.8	-		5.5	-
. AAMR (1940 Std. mil.)		2.4	0.15		3.6	0.28		1.4	0.14
. AAMR (1970 Std. mil.)		3.6	0.19		5.4	0.37		2.4	0.19
. AAMR (2000 Std. mil.)		5.4	0.27		7.9	0.55		3.8	0.28
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.6	-		1.7	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is not significantly different
- . The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3, E922.8-.9, E955.0-.4, E965.0-.4, E970, E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.2	0.22	0	0.0	0.00	1	0.4	0.44
10-14	2	0.5	0.32	2	0.9	0.63	0	0.0	0.00
15-19	41	10.1	1.58	38	18.2	2.95	3	1.5	0.88
20-24	60	16.6	2.15	54	29.3	3.99	6	3.4	1.39
25-29	50	11.6	1.65	47	21.9	3.20	3	1.4	0.80
30-34	42	7.9	1.22	40	15.1	2.40	2	0.7	0.53
35-39	45	7.7	1.14	34	11.7	2.00	11	3.7	1.12
40-44	34	6.2	1.07	28	10.4	1.97	6	2.1	0.88
45-49	14	3.0	0.81	11	4.9	1.48	3	1.3	0.74
50-54	21	5.2	1.14	19	9.7	2.22	2	1.0	0.68
55-59	15	4.9	1.27	13	8.8	2.45	2	1.3	0.90
60-64	14	5.7	1.53	13	11.1	3.09	1	0.8	0.79
65-69	11	4.4	1.34	10	8.8	2.78	1	0.7	0.75
70-74	8	3.5	1.22	8	8.0	2.83	0	0.0	0.00
75-79	5	2.5	1.11	5	6.1	2.71	0	0.0	0.00
80-84	10	7.2	2.29	10	19.6	6.20	0	0.0	0.00
85+	7	5.8	2.19	6	18.0	7.34	1	1.1	1.15
Total deaths	380			338			42		
Connecticut									
. Crude rate		5.8	-		10.6	-		1.2	-
. AAMR (1940 Std. mil.)		6.2	0.34		11.2	0.64		1.3	0.21
. AAMR (1970 Std. mil.)		5.7	0.31		10.4	0.59		1.2	0.20
. AAMR (2000 Std. mil.)		5.9	0.30		10.8	0.59		1.3	0.20
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.2	-		21.1	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is lower

Connecticut Resident Age-Specific Mortality Rates by Sex, 1997-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.2	0.23	0	0.0	0.00	1	0.5	0.48
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	10	2.5	0.78	7	3.4	1.27	3	1.5	0.88
20-24	32	8.9	1.57	24	13.0	2.66	8	4.5	1.60
25-29	45	10.5	1.56	35	16.3	2.76	10	4.6	1.47
30-34	92	17.3	1.80	67	25.4	3.10	25	9.3	1.87
35-39	127	21.7	1.92	99	34.0	3.42	28	9.5	1.79
40-44	115	21.0	1.96	84	31.3	3.42	31	11.1	1.99
45-49	68	14.8	1.79	45	20.0	2.98	23	9.8	2.04
50-54	22	5.5	1.16	14	7.1	1.91	8	3.9	1.37
55-59	14	4.6	1.23	7	4.8	1.80	7	4.5	1.69
60-64	5	2.1	0.92	1	0.9	0.86	4	3.1	1.57
65-69	3	1.2	0.70	0	0.0	0.00	3	2.2	1.30
70-74	8	3.5	1.22	5	5.0	2.24	3	2.3	1.32
75-79	7	3.5	1.31	4	4.9	2.43	3	2.5	1.45
80-84	7	5.1	1.92	2	3.9	2.77	5	5.8	2.57
85+	5	4.1	1.85	2	6.0	4.24	3	3.4	1.99
Total deaths	561			396			165		
Connecticut									
. Crude rate		8.6	-		12.5	-		4.9	-
. AAMR (1940 Std. mil.)		7.9	0.35		11.5	0.59		4.4	0.36
. AAMR (1970 Std. mil.)		6.8	0.30		9.7	0.51		4.0	0.33
. AAMR (2000 Std. mil.)		8.3	0.35		11.9	0.60		4.7	0.37
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.6	-		7.8	-		3.3	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-
Comparison of the CT Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is higher
. THE U.S. HP 2000 target:									CT rate is higher