# Ellington Quadrangle Bedrock Geology w/Explanation

M.H. Pease, Jr.

**Explanation** 

Map

## NOTICE !


This document contains unpublished maps, cross- sections, and related information archived by the State Geological and Natural History Survey of Connecticut as part of the Survey Library Collection.


These materials have not been reviewed for accuracy, consistency, or completeness. For many geographic areas, more current information exists, either in published or unpublished form.

For the most part, these materials were developed under research and mapping agreements between the State Geological Survey and individual scientists, academic institutions, or graduate students. Some of these materials have been received by the State Geological Survey as donations. The veracity of the information contained within these documents is the responsibility of the authorship. The State Geological and Natural History Survey of Connecticut, does not promote or endorse this content, nor does the State Survey attest as to its level of accuracy.

These materials have been preserved under a cooperative agreement between the State Geological Survey and the US Geological Survey as part of the National Geological and Geophysical Data Preservation Program. <u>www.datapreservation.usgs.gov</u>


These materials are offered in the spirit of open government. Reproduction of these manuscripts was conducted to the highest practical degree, within the parameters of the funding mechanism. Original documents are available for inspection by contacting the <u>Connecticut State Geologist</u>.





- ARMA

The .



40 7

B

BEDROCK LITHOLOGY

25 \$

G

ELLINGTON QUADRANGLE, CONNECTICUT BY M.H.PEASE, JR. 1972

U.S. Geological Survey OPEN FILE MAP This map is preliminary and has not been addied or reviewed for conformity with Geocheal Survey standards or nemenclature.

1 41 . 52 . 30"

72 221 301

12

get.

## EXPLANATION

The purpose of this map is to describe geologic map units in terms of physical properties useful for engineering and land management. Map units are derived from the published geologic map by Collins (1954), modified from data obtained by recent detailed management in parts of this quadrangle and adjacent quadrangles, and from interpretation of adronagnetic data. Unlike a geologic map, this map is not concerned with relative stratigraphic dosint tion, and geologic hames are omitted. Units are designated by letter and most units include a variety of cock types each with significantly different physical properties. These variations are too small to delineate at the map scale. This explanation is keyed to a bedrock-properties matrix (Table 1) that classifies the bedrock in terms of significant physical properties. Rock terms <u>underlined</u> below correspond to specific rock types listed and described in Table 1.

Bedrock in the EMIngton quadrangle is separable into two major divisions by the prominent north-south trending fault near the center of the quadrangle. Rocks to the west are relatively soft, pervious sedimentary rocks almost entirely covered by placial debris. Rocks east of the fault are hard, relatively impervious metamorphic and igneous rocks covered by a patchy mantle of thin glacial debris.

Color terms describe fresh tock.



Lithology: light-gray, poorly lawered gneiss. Occurrence: extreme southeast corner of quadrangle, two small exposures.

Lithology: dark-greenish-gray to greenish-black <u>platy schist</u>, layers 1 to 10 feet thick, interlayered with medium-greenish-gray <u>layered gneiss</u> in about equal proportions. Dark-greenish-gray massive gneiss common a locally.

Occurrence: in southeast corner of guadrangle, a cortheast-trending band mostly of <u>platy</u> schist rests on A; good exposure in road cuts on Baid Hill Boad where it crosses ridge (701, 800 E: 382, 400 w - Connecticut State Coordinates, in feet). Block immediately to west mostly of <u>massive gneiss</u>, intertongues northeast with C. Well exposed on Soapstone Mountain and in several smaller areas near center of quadrangle, mostly <u>layered gneiss</u> with <u>massive gneiss</u> porticularly near creat of mountain; these areas enclosed of unit 6. Several large blocks along west side of major fault and at morthern border of quadrangle, mostly interlayered <u>platy schist</u> and <u>layered gneiss</u>; representative outcrops in scarps at south end Bald Mountain just north of County Road (690, 150 E; 419, 700 N) where contact with unit 6 is exposed.

Prospects and quarries: at least two prospects on southeast slope Soapstone Mountain from which an inferior grade of soapstone has been quarried.

	10000				
1.45		10.1			
1.1					
1.1					
- A.			Canal -		
1.0			5.7		
558.0					
2.281			340		
e ga					
1.081					
100					

Lithology: mostly medium-gray, medium-to-fine-grained layered gneiss with misor light-gray. Timer-grained layered cneiss. Distinguished from unit 8, with which it intertongues, by lack of greenish color; less subject to parting along layers than unit 8. Physical properties similar to that of unit 6, but less massive.

Occurrence southeast corner of quadrangle, fine-grained, granular layered meiss, intertongues with unit 8. Northeast frending band through Newell Hill and area of hill 663 north of Hopkins Road, mostly medium-grained layered gneiss to massive gneiss. In two other structural blocks outcrops are poor and presence is inferred. Representative exposure on crest of hill 663.

Lithology: blocks in which 8 and C are interlayered or undifferentiated Occurrence: interlayered 8 and C exposed in triangular shaped block west and southwest of Crystal Lake. Representative exposures: I. In gravel pit between Burbank Road and State Highway 30 (699, 400 E; 400, 750 N), greenish gray to black <u>platy schist</u> and <u>layere6 gneiss</u> interlayered with Hight-gray <u>layered gneiss</u>. 2 road cuts on State Highway 30 at Crystal Lake (701, 300 E; 403, 800 N). Exposures of unit 8 west of road highly contorted by faulting: thinly <u>layered gneiss</u> of unit C exposed east of road. Undifferentiated 8,C in blocks with no exposed bedrock on southeast side of fault along Martins Brook. Presence inferred.



Dithelogy: light-gray flaggy quartzite in beds as much as one foot thick interlayered with 1/16 to one inch-thick beds of light-to-medium-gray mica-rich schust.

Occurrence: two bands in southeast corner of guadrangle well exposed in hogback ridges. Unit commonly has high magnetic susceptibility and occurrence in area of no exposure inferred in part from magnetic anomaly. Representative exposure in small abandened quarry at road intersection (elevation 816) on Hunter Rd. 3008 ft. southeast of Crystal Lake Ed. (701, 400 Fr 397, 300 N).

Prospects and quarries: unit has been quarried for flagstone and building stone at several localities; no operating quarries in quadrangle.



Lithology: gray platy schist interlayered with gray mica-rich schist in beds 1/16 to 6 inches thick; thin <u>quartzite</u> lenses amounting to less than 5 percent of the unit. Occurrence:northeast trending band about 1/2 mile wide along northwest side of fault in southeast corner of quadrangle. Generally poorly exposed; representative outcrop just west of Hunter Road (543, 500 E; 389, 600 N) but here the layered rocks are more conspicuously folded than elsewhere.



Lithology: brown-weathering mica-rich schist about 75 percent and rusty-weathering sullidic schist about 25 percent, interlayered on scale of 1/16 Inch to 5 feet Occurrance:mostly burled by sufficial debris, but exposed in 3 small, widely separated outcrops. Aerial extent of blocks on map largely interpretive. 1. Outcrop on State Highway 30 just north of Grant Brook (697, 950 E; 394, 300 N). 2. Outcrop on State Highway 190 just west of ScullyRoad intersection (686, 700 E; 419, 500 N). Here unit F-is in contact with greenish-gray platy schist of unit B. 3. North border of guadrahole on west side of major fault.



Lithology: medium-to light-gray, medium-grained massive gneiss with minor thickly layered gneiss.

Occurrence: occurs in most of the large structural blocks between the major fault and the fault along Martins Brook. Unit 5 is bordered by and encloses units 8 and C and contains many more inclusions of 8 and 6, too small to show on map. Accessible representative exposures crop out along Shenipsit Lake; extensive outcrops also along the crest and east slope of Baid Mountain.



Lithology: mottled dark-and light-greenish-gray, medium-to-coarse-grained <u>massive gneiss</u>. <u>Occurrence</u>:exposed on several small knobs within unit G in northeast part of the quadrangle.

Lithology: rock breccia more or less cemented by silica; i.e. silicitied rock. <u>Occurrence</u>:mapped in two narrow slivers along major fault. Probably occurs at other localities along major fault and may be present along other faults. Representative exposures in small excavation north of Mountain View Road (683, 500 E: 415, 250 N): greenishgray platy schist fragments cemented by silica. Quartz veins as much as 6 inches wide are common.

Lithology: dark+oray, dense diabase; crayish-brown weathered rind componly about 1 ch. thick Occurrence:narrow linear features less than 200 feet wide extending in a north to portheast direction. Principal body extends for a distance of about 4 miles along a fault zone that passes just east of Soapstone Mountain. Representative exposure on east side of eastern road to Soapstone Mountain lookout (692, 150 E; 410, 000 N).

Prospects and quarries: diabase is quarried for road metal and crushed stone aggregate outside of quadrangle: no operating quarries in quadrangle.



Lithology: consists of reddish-brown sandstone, 60 percent; and <u>conclonerate</u> with peobles as large as 3 to 4 inches, 30 percent; with minor amounts of <u>slitstone</u> and <u>shale</u>. Exposures elsewhere in Connecticut suggest that <u>conclonerate</u> becomes more common and coarser toward the major fault.

Occurrence:underlies entire area west of major fault. Crops out only in small exposures in southwest corner of quadrangle and a few exposures to the north. Also encountered in several water wells

Prospects and quarries: sandstone has been quarried for dimension stone at many localities in the Connecticut Valley, but no commercial quarries are known to have operated in the Ellington quadrangle.

### Symbols



Boundaries between mapped geologic units. The symbol may represent a sharp lithologic break or, more commonly, a gradational zone. Contacts are rarely exposed.



Boundary between blocks of bedrock along which displacement probably has occurred. Hay be narrow surface or a zone of rupture several hundred feet wide within which rock generally is weak, and joints and fractures are classify spaced tending to decrease in abundance outward. Zone of weakness may extend to depths of **hundreds** of feet and act as channels for groundwater or mineral-bearing solutions. Breccia (severely fragmented rock) and couge (pulverized clayey material) may be common. Fault traces generally form troughs of low relief or scarps carved in weak fractured rock and subsequently covered by overburden.

## 20

#### Strike and dip of foliation

Symbol defines the attitude at a specific locality of a surface or surfaces formed by preferred orientation of minerals in metamorphic rocks. In the Ellington quadrangle this follation is parallel to layering. The line is oriented parallel to the strike of foliation and indicates the direction in which the foliation trends on a horizontal (or map) surface: the solid triangle indicates the direction in which the layering or foliation is inclined; the number represents degrees of inclination. The known data concerning distribution and

attitude of exposed bedrock is shown by these symbols on the geologic man.

## Erike and dip of bedding

Symbol defines the attitude of bedding (layering) in sedimentery rocks. The line tepresents the strike and indicates the direction that the bedding trends on a horizontal (or map) surface. The short line represents the dip at right angles to the **strike** and indicates the direction in which the bedding or layering is inclined; the number represents degrees of inclination.

### References

Collins, G. E., 1954. The bedrock geology of the Ellington quadrangle, Connecticut: State Geological and Natural History Survey, Quadrangle Report No. 4.

Pitkin, J. A., Philbin, P. W., and Gilbert, F. P., 1969, Aeromagnetic map of the Ellington quadrangle and part of the Rockville quadrangle, Hartford and Tolland Counties, Connecticut: U. S. Geological Survey Geophysical Investigations Map GP+648.

U.S. Geological Survey OPEN FILE REPORT This report is preliminary and has not been edited or reviewed for conformity with Geological Survey standards or nomenclature.

To secompany: Plate 1 Bedrock Lithology Ellington Quadrangle, Connecticut by M.H. Pease, Jr. 1972