Page: Run Date: 8 / 20 / 02

Connecticut **Summary of Demographic Indicators** Children Aged < 5 Years (2,3)

Total Records	Number
Individual Children	51,410
Total Records (All Visits)	51,410

Source of Data	Number	%	
WIC EPSDT Child Health-MCH Child Health-HMO Headstart School Health Other		51,410 0 0 0 0 0 0	100.0 0.0 0.0 0.0 0.0 0.0 0.0
Total <i>Excluded Unknown:</i> 0	0.0 %	51,410	100.0

Race/Ethnic Distribution	Race/Ethnic Distribution			
White, Not Hispanic Black, Not Hispanic Hispanic American Indian/Alaskan Native Asian/Pacific Islander All Other		15,215 14,263 19,724 92 623 1,031	29.9 28.0 38.7 0.2 1.2 2.0	
Total		50,948	100.0	
Excluded Unknown: 462	0.9 %			

Age Distribution (4)	Number	%
0 - 5 Months 6 - 11 Months 12 - 23 Months 24 - 35 Months 3 - 4 Years	11,696 6,156 10,865 8,470 14,223	22.8 12.0 21.1 16.5 27.7
Total	51,410	100.0

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.(3) Excludes records with unknown data or errors.

⁽⁴⁾ Records rejected if date of birth and date of visit unknown.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 2C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Health Indicators Children Aged < 5 Years (2)

	Summary of Hea	alth Indicato	rs ⁽³⁾		
		Connect Prevale		Natio Preva (Prior	lence
		Number	%	Number	%
Birthweight (4) Low High	< 2500 g > 4000 g	13,868 13,868	9.0 7.9	1,204,78 1,204,78	
Height and Weight Short Stature (5)	< 5th < 10th	50,033 50,033	6.2 11.3	4,428,86 4,428,86	
Underweight (6)	< 5th < 10th	50,033 50,033	5.2 8.5	4,428,86 4,428,86	
Overweight (6) ≥ 2 Yrs Overweight (6)	≥ 95th 85th - < 95th ≥ 95th	50,033 22,084 22,084	16.3 17.3 18.7	4,428,86 1,796,49 1,796,49	3 15.1
Anemia (7) Low Hb Low Hct Low Hb/Hct		37,291 12,432 39,670	10.7 13.4 10.4	2,687,45 530,84 3,082,39	0 14.5 6 9.6
Breastfeeding (8) Ever Breastfed Breastfed At Least 6 Mol Breastfed At Least 12 Mol		13,868 8,784 7,977	49.3 23.4 17.7	875,42 333,89 319,82	2 20.6

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Infants born during the reporting period included in the analysis.
- (5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th < 95th percentile category identifies childrisk of overweight.
- (7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- (8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- * Percentages are not calculated if < 100 records are available for analysis after exclusions.



Page:

Table 3C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Breastfeeding Indicators Children Aged < 5 Years (2)

Connecticut

Summary of Breastfeeding Indicators (3)								
	Connec Prevale		National Prevalence (Prior Year)					
	Number	%	Number	%				
Breastfeeding Duration (4	4,5)							
Ever Breastfed	13,868	49.3	875,423	47.5				
Breastfed 1+ Weeks Breastfed 2+ Weeks Breastfed 3+ Weeks Breastfed 4+ Weeks Breastfed 6+ Weeks	6,559 5,949 5,803 5,915 6,113	51.3 44.1 41.4 39.5 35.9	565,388 434,296 346,229 292,204 230,422	50.1 48.5 47.1 45.1 39.9				
Breastfed 2+ Months Breastfed 3+ Months Breastfed 4+ Months Breastfed 5+ Months Breastfed 6+ Months	6,631 7,300 8,253 8,940 8,784	31.2 28.0 26.3 24.9 23.4	207,623 224,692 267,785 316,450 333,892	34.0 28.6 25.1 22.2 20.6				
Breastfed 9+ Months Breastfed 12+ Months	7,020 7,977	19.5 17.7	287,084 319,827	14.9 12.3				
Breastfed 18+ Months	4,258	12.3	187,037	7.3				

⁽¹⁾ Reporting period is January 1 through December 31.



Page:

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. For each breastfeding duration category, the analysis includes only children who turned that age during the reporting period by/on their date of visit. For example, infants who turned 9 months old during the reporting period by/on their date of visit are included in the 9+ months analysis.

⁽⁵⁾ It is possible the analysis will show that slightly more infants are breastfed at older ages, e.g. 1+ week of age, than at younger ages, e.g. at birth (ever breastfed), due to variation in infants included in each breastfeeding analysis category.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

2001 Pediatric Nutrition Surveillance (1) Connecticut

Comparison of Racial and Ethnic Distribution by Contributor to Nation Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution										
Contributor Number Included in Analysis (3) Number Included in Analysis (3) White, Not Hispanic Hispanic % Black, Not Hispanic Hispanic % Hispanic % Hispanic Malaskan Native % American Indian/ Asian/Pacific Islander Unknown %										
Connecticut	51,410	29.6	27.7	38.4	0.2	1.2	2.9			
Nation (Prior Year)	4,820,831	42.2	21.9	28.3	1.5	2.9	3.3			



Page:

⁽¹⁾Reporting period is January 1 through December 31. (2)Analyses based on one record per child. (3)Excludes records with errors.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

2001 Pediatric Nutrition Surveillance (1) Connecticut

Comparison of Age Distribution by Contributor to Nation Children Aged < 5 Years (2)

Comparison of Age Distribution									
Contributor	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %					
Connecticut	51,410	34.7	21.1	44.1					
Nation (Prior Year)	4,820,831	38.4	21.7	39.9					

Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.(3) Excludes records with unknown data or errors.

⁽⁴⁾ Records rejected if date of birth and date of visit unknown.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 1 Run Date: 8 / 20 / 02

Comparison of Growth and Anemia Indicators by Contributor to Nation Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)													
		Birthweigh	t (4)	Height and Weight							- An	One or More Low/High	
		Low	High		Short Stature (5)	Underweight (6)			≥ 2 Yrs Over- weight (6	S)	Low	Health Indicators (8)	
State	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th- <95	% th <u>≥</u> 95th	Number	%	%
Connecticut	13,868	9.0	7.9	50,033	6.2	5.2	16.3	22,084	17.3	18.7	39,670	10.4	41.1
Nation (Prior Year)	1,204,782	8.9	8.3	4,428,86	3 6.1	5.4	12.8	1,796,49	93 15.1	13.0	3,082,394	1 13.6	38.2



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

⁽⁸⁾ One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators. LBW, HBW, short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated. Not calculated if one or more health indicators are missing data.

^{*} Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7C

2001 Pediatric Nutrition Surveillance (1)

Connecticut

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor to Nation Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing (4) and Smoking in Household									
		Breastfeeding (5)							
State	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months			
Connecticut	13,868	49.3	8,784	23.4	7,977	17.7			
Nation (Prior Year)	875,423	47.5	333,892	20.6	319,827	12.3			

⁽⁵⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6

* Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ TV Viewing and Smoking in Household data are not reported by all contributers.

⁽³⁾ Analyses based on one record per child.

⁽⁴⁾ Excludes records with unknown data or errors.

Table 8C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 1 Run Date: 8 / 20 / 02

Growth and Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

	Growth and Anemia Indicators by Race/Ethnicity and Age (3)											
	Bir	thweight (4	1)			Hei	ght and W	/eight				
		Low High % %			Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Ane Low H	b/Hct
	Number	<2500 g	>4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number	% 85th-<95th	% ı <u>></u> 95th	Number	%
Race/Ethnicity												
White, Not Hispanic	4,325	8.4	9.9	14,869	6.4	5.6	13.7	6,259	17.5	15.2	11,413	7.7
Black, Not Hispanic	3,815	11.5	6.2	13,810	6.3	5.5	15.7	6,132	16.7	16.5	11,119	14.3
Hispanic	5,078	7.5	7.6	19,205	6.0	4.5	19.2	8,850	17.7	23.1	15,470	9.7
American Indian/Alaskan Native	29	*	*	90	*	*	*	36	*	*	70	*
Asian/Pacific Islander	217	10.6	6.0	608	6.7	8.7	12.7	222	14.0	16.7	446	9.6
All Other/Unknown	404	9.4	7.7	1,451	5.7	7.0	12.3	585	17.4	13.2	1,152	10.9
Total	13,868	9.0	7.9	50,033	6.2	5.2	16.3	22,084	17.3	18.7	39,670	10.4
Age												
0 - 11 Months	_	_	_	17,237	6.8	7.5	11.7	0	*	*	6,148	15.8
12 - 23 Months	-	-	-	10,712	8.2	4.2	18.9	0	*	*	10,849	10.1
24 - 35 Months	-	-	-	8,294	4.8	5.1	16.4	8,294	15.7	16.4	8,462	11.2
36 - 47 Months	-	-	-	6,971	5.1	3.4	19.7	6,971	18.7	19.7	7,171	8.9
48 - 59 Months	-	-	-	6,819	4.5	2.9	20.4	6,819	17.9	20.4	7,040	6.9
Total	-	-	_	50,033	6.2	5.2	16.3	22,084	17.3	18.7	39,670	10.4



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusion

2001 Pediatric Nutrition Surveillance (1)

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity and Age (2) Children Aged < 5 Years (3)

Breastfeeding, TV Viewing and Smoking in Household by Race/Ethnicity and Age (4)									
	Breastfeeding (5)								
	Number E	% Ever Breastfed	Number	Breastfed At Least 12 Months					
Race/Ethnicity White, Not Hispanic Black, Not Hispanic Hispanic American Indian/Alaskan Native Asian/Pacific Islander All Other/Unknown Total	4,325 3,815 5,078 29 217 404 13,868	44.7 43.0 56.4 * 48.8 69.1 49.3	2,510 2,551 3,296 13 125 289 8,784	20.2 23.1 24.6 * 18.4 41.2 23.4	2,368 2,150 3,084 15 86 274 7,977	15.7 16.8 18.3 * * 31.8			
Age 0 - 11 Months 12 - 23 Months 24 - 35 Months 36 - 47 Months 48 - 59 Months	- - - -	- - - -	- - - -	- - - -		- - - -			
Total	13,868	49.3	8,784	23.4	7,977	17.7			



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ TV Viewing and Smoking in Household are not reported by all contributors.

⁽³⁾ Analyses based on one record per child.

⁽⁴⁾ Excludes records with unknown data or errors.

⁽⁵⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Summary of Trends in Racial and Ethnic Distribution

Page: Run Date: 8 / 20 / 02

Children Aged < 5 Years (2)

		Summary of Tr	ends in Racial	and Ethnic D	istribution		
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/ Alaskan Native %	Asian/Pacific Islander %	All Other/ Unknown %
2001	51,410	29.6	27.7	38.4	0.2	1.2	2.9
2000	857	35.8	28.5	32.1	0.0	0.8	2.8
1999	13	*	*	*	*	*	*
1998	53,177	31.8	29.7	35.2	0.2	1.2	2.0
1997	49,384	32.9	29.3	34.7	0.2	1.3	1.6
1996	0	*	*	*	*	*	*
1995	5,838	36.4	25.9	36.2	1.2	0.3	0.0
1994	6,823	38.8	26.2	33.8	1.1	0.2	0.0
1993	6,625	39.3	26.2	33.3	1.0	0.3	0.0
1992	3,587	37.9	29.5	31.7	0.8	0.2	0.0
1991	6,882	42.9	26.0	30.3	0.6	0.1	0.0
1990	6,002	42.8	28.4	28.1	0.5	0.1	0.0
1989	5,867	44.7	28.7	25.6	0.7	0.2	0.0
1988	5,907	51.0	26.3	21.7	0.6	0.4	0.0
1987	5,234	54.2	25.1	19.9	0.6	0.3	0.0
1986	2,396	59.2	28.1	11.7	0.7	0.3	0.0
1985	2,344	65.9	23.6	9.4	0.8	0.4	0.0
1984	2,529	70.1	19.4	9.6	0.6	0.4	0.0
1983	2,009	77.1	11.1	10.0	0.4	1.3	0.0

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.(3) Excludes records with errors.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 10C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Racial and Ethnic Distribution Children Aged < 5 Years (2)

	Summary of Trends in Racial and Ethnic Distribution										
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/ Alaskan Native %	Asian/Pacific Islander %	All Other/ Unknown %				
1982	3,232	57.3	17.8	23.2	0.3	1.4	0.0				
1981	40	*	*	*	*	*	*				



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.(3) Excludes records with errors.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Summary of Trends in Age Distribution

Children Aged < 5 Years (2)

	Summa	ary of Trends in Age	Distribution (3)	
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %
2001	51,410	34.7	21.1	44.1
2000	857	45.6	15.5	38.9
1999	13	*	*	*
1998	53,177	34.8	20.4	44.8
1997	49,384	26.8	22.5	50.7
1996	0	*	*	*
1995	5,838	34.8	18.6	46.6
1994	6,823	34.7	18.3	47.0
1993	6,625	33.8	18.9	47.4
1992	3,587	33.0	20.2	46.8
1991	6,882	35.3	20.0	44.6
1990	6,002	36.2	19.5	44.3
1989	5,867	35.0	19.6	45.4
1988	5,907	35.0	20.2	44.8
1987	5,234	33.0	19.3	47.7
1986	2,396	37.1	22.3	40.7
1985	2,344	42.0	21.2	36.7
1984	2,529	53.6	16.5	29.9

⁽¹⁾ Reporting period is January 1 through December 31.

Page:

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Records rejected if date of birth and date of visit unknown.

 ⁽⁴⁾ Excludes records with unknown data or errors.
 Percentages are not calculated if < 100 records are available for analysis after exclusions.

2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Age Distribution Children Aged < 5 Years (2)

Summary of Trends in Age Distribution (3)									
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %					
1983	2,009	59.6	15.6	24.8					
1982	3,232	59.8	13.6	26.6					
1981	40	*	*	*					



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Records rejected if date of birth and date of visit unknown.

⁽⁴⁾ Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 1 Run Date: 8/20/2002

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

		Sumr	mary of ⁻	Trends i	n Grov	th and	l Anem	nia Indio	cators (3))			
	Bi	irthweight ((4)		Height and Weight							Anemia	
Year		Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Low Hb/Hct (7)		
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%	
2001	13,868	9.0	7.9	50,033	6.2	5.2	16.3	22,084	17.3	18.7	39,670	10.4	
2000	376	7.4	9.0	829	3.3	5.9	13.6	319	17.9	16.6	591	10.7	
1999	0	*	*	2	*	*	*	0	*	*	5	*	
1998	14,129	10.3	8.0	51,780	6.7	5.3	15.4	23,272	17.8	17.8	41,153	10.8	
1997	8,817	9.4	8.1	48,370	6.9	4.9	15.9	24,511	17.1	16.6	42,707	10.8	
1995	992	8.9	8.9	5,683	4.7	6.6	11.5	2,659	15.0	13.2	3,407	9.4	
1994	1,448	7.9	8.6	6,620	4.8	7.4	11.2	3,096	13.8	12.7	4,269	7.3	
1993	1,215	10.1	6.0	6,431	4.7	7.4	10.7	3,068	12.9	11.8	4,065	6.0	
1992	623	8.2	8.7	3,469	5.2	7.5	12.2	1,639	13.8	12.6	1,911	8.7	
1991	1,401	8.3	7.6	6,582	4.6	8.2	10.5	2,952	14.7	11.3	4,497	9.4	

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and height-for-age

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 2 Run Date: 8/20/2002

Summary of Trends in Growth and Anemia Indicators Children Aged < 5 Years (2)

	Summary of Trends in Growth and Anemia Indicators (3)											
	Ві	irthweight ((4)			Heiç	ght and W	/eight			Anemia Low Hb/Hct	
Year		Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			(7)	
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number	% 85th-<95th	% ≥95th	Number	%
1990	1,204	9.1	8.0	5,705	5.0	7.4	10.5	2,588	14.5	11.0	3,907	7.8
1989	1,237	10.3	7.0	5,570	4.8	6.8	10.4	2,582	13.7	11.0	3,909	8.4
1988	1,092	7.9	9.2	5,598	4.5	7.5	10.1	2,556	12.9	11.0	3,886	8.6
1987	1,039	8.3	9.9	4,983	4.7	7.2	9.3	2,416	13.6	9.9	3,209	7.5
1986	487	8.4	9.2	2,320	5.9	6.4	10.6	949	12.8	8.7	1,498	7.6
1985	615	9.1	7.8	2,268	6.7	8.1	8.8	852	11.0	6.7	1,506	11.0
1984	926	9.4	9.9	2,420	6.2	9.1	6.4	723	12.3	6.6	1,311	10.1
1983	844	8.9	8.8	1,928	6.6	12.6	7.6	488	8.8	7.2	978	9.6
1982	1,653	9.7	6.2	2,991	8.5	11.3	6.4	838	13.0	7.6	1,371	10.9
1981	30	*	*	26	*	*	*	4	*	*	6	*

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and height-for-age for age and height-for-age for children 2 years of age and height-for-age for age age and heig

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Breastfeeding Children Aged < 5 Years (2)

	Summary of Trends in Breastfeeding (3)									
				etfeeding (4)						
Year	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months				
2001	13,868	49.3	8,784	23.4	7,977	17.7				
2000	377	46.4	196	15.8	131	11.5				
1999	0	*	7	*	1	*				
1998	14,129	42.7	8,760	22.4	7,585	16.8				
1997	8,817	40.2	8,726	17.5	7,695	11.8				
1996	0	*	0	*	0	*				
1995	854	34.4	255	10.2	161	6.8				
1994	1,189	34.1	462	13.4	302	5.6				
1993	1,053	33.0	384	11.2	259	6.2				
1992	557	26.0	288	8.3	224	4.5				

⁽¹⁾ Reporting period is January 1 through December 31.



Page:

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Breastfeeding Children Aged < 5 Years (2)

	Summary of Trends in Breastfeeding (3)									
				etfeeding (4)						
Year	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months				
1991	1,251	28.3	535	7.7	446	4.9				
1990	1,066	30.8	493	9.1	391	3.1				
1989	1,108	27.6	540	8.7	349	4.9				
1988	1,008	27.6	618	6.6	404	3.7				
1987	968	26.0	624	7.7	490	4.5				
1986	486	37.4	284	10.2	202	4.0				
1985	616	36.4	334	8.7	182	7.7				
1984	925	38.9	372	16.9	194	10.3				
1983	846	40.3	252	19.0	132	14.4				
1982	1,653	30.7	334	13.8	221	5.4				

⁽¹⁾ Reporting period is January 1 through December 31.



Page:

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Breastfeeding Children Aged < 5 Years (2)

	Summary of Trends in Breastfeeding (3)									
			Breas	stfeeding (4)						
Year	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months				
1981	30	*	0	*	2	*				



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 14C 2001 Pediatric Nutrition Surveillance Children Aged < 5 Years

Page: 1 Run Date: 8 / 20 / 02

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 15C 2001 Pediatric Nutrition Surveillance Children Aged < 5 Years

Page: 1 Run Date: 8 / 20 / 02

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 16C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Growth Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Growth Indicators by Race/Ethnicity and Age (3) Underweight Overweight ≥ 2 Yrs Overweight Short Stature (4) (5) (5) (5) % % % % % Number Number Number Number <5th 85th - <95th <5th >95th >95th White, Not Hispanic 0 - 11 Months 5.423 6.4 5.423 7.9 5.423 10.4 0 0.0 0.0 12 - 23 Months 3,187 8.3 3,187 4.4 3,187 16.5 0 0.0 0.0 24 - 59 Months 6,259 5.6 6,259 4.1 6,259 15.2 6.259 17.5 15.2 Total 5.6 15.2 14,869 6.4 14,869 14,869 13.7 6,259 17.5 Black, Not Hispanic 4.734 4,734 4,734 0 0 - 11 Months 7.9 7.5 12.6 0.0 0.0 2,944 2,944 2,944 0 0.0 12 - 23 Months 9.0 4.7 19.0 0.0 24 - 59 Months 6.132 3.7 6.132 4.3 6,132 16.5 6.132 16.7 16.5 **Total** 13.810 6.3 13.810 5.5 13.810 15.7 6.132 16.7 16.5 Hispanic 0 - 11 Months 6,253 6,253 6.3 6,253 6.9 12.5 0 0.0 0.0 4.102 0 0.0 12 - 23 Months 4,102 7.8 3.2 4.102 21.3 0.0 8.850 3.3 8.850 23.1 8.850 23.1 24 - 59 Months 4.9 8.850 17.7 Total 19,205 6.0 19,205 4.5 19,205 19.2 8,850 17.7 23.1 American Indian/Alaskan Native 0 - 11 Months 33 33 33 0 21 21 21 0 12 - 23 Months 24 - 59 Months 36 36 36 36

90

36

Total

90

90



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 16C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Growth Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2) Page: 2 Run Date: 8/20/2002

	Gı	rowth Ind	icators by F	Race/Ethni	city and Ag	je ⁽³⁾			
	Short St	Short Stature (4)		veight)	Overweight (5)		≥ 2 Yrs Overweight (5)		
	Number	% <5th	Number	% <5th	Number	% <u>≥</u> 95th	Number	% 85th - <95th	% <u>></u> 95th
Asian/Pacific Islander									
0 - 11 Months	270	7.0	270	10.4	270	10.7	0	0.0	0.0
12 - 23 Months	116	5.2	116	10.3	116	9.5	0	0.0	0.0
24 - 59 Months	222	7.2	222	5.9	222	16.7	222	14.0	16.7
Total	608	6.7	608	8.7	608	12.7	222	14.0	16.7
All Other/Unknown									
0 - 11 Months	524	5.3	524	8.4	524	9.7	0	0.0	0.0
12 - 23 Months	342	7.0	342	6.4	342	14.6	0	0.0	0.0
24 - 59 Months	585	5.1	585	6.0	585	13.2	585	17.4	13.2
Total	1,451	5.7	1,451	7.0	1,451	12.3	585	17.4	13.2
Total All Race/Ethnic Groups									
0 - 11 Months	17,237	6.8	17,237	7.5	17,237	11.7	0	0.0	0.0
12 - 23 Months	10,712	8.2	10,712	4.2	10,712	18.9	0	0.0	0.0
24 - 59 Months	22,084	4.8	22,084	3.9	22,084	18.7	22,084	17.3	18.7
Total	50,033	6.2	50,033	5.2	50,033	16.3	22,084	17.3	18.7



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and height-for-age

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Connecticut	
Anemia Indicators by Race/Ethnicity	and Age
Children Aged < 5 Years	(2)

	Aneı	mia Indicat
		Anemia
		Low Hb/Hct
	Number	%
White, Not Hispanic		
6 - 11 Months	1,806	12.7
12 - 17 Months	1,905	7.7
18 - 23 Months	1,312	6.6
24 - 35 Months	2,416	8.2
36 - 59 Months	3,974	5.4
Total	11,413	7.7
Black, Not Hispanic		
6 - 11 Months	1,816	18.7
12 - 17 Months	1,698	13.3
18 - 23 Months	1,293	14.4
24 - 35 Months	2,357	15.0
36 - 59 Months	3,955	12.3
Total	11,119	14.3
Hispanic		
6 - 11 Months	2,206	16.2
12 - 17 Months	2,355	10.2
18 - 23 Months	1,800	8.7
24 - 35 Months	3,338	10.7
36 - 59 Months	5,771	6.7
Total	15,470	9.7
American Indian/Alaskan Native		_
6 - 11 Months	12	*
12 - 17 Months	13	*
18 - 23 Months	9	*
24 - 35 Months	14	*
36 - 59 Months	22	*
Total	70	*

tors by F	Race/Ethnicity and Age (3,	4)	
			Anemia
			Low Hb/Hct
		Number	%
1	Asian/Pacific Islander		
	6 - 11 Months	102	9.8
	12 - 17 Months	68	*
	18 - 23 Months	50	*
	24 - 35 Months	98	*
	36 - 59 Months	128	6.3
	Total	446	9.6
1	All Other/Unknown		
	6 - 11 Months	206	17.5
	12 - 17 Months	207	11.1
	18 - 23 Months	139	11.5
	24 - 35 Months	239	10.5
	36 - 59 Months	361	6.9
	Total	1,152	10.9
	Total All Race/Ethnic Groups		
	6 - 11 Months	6,148	15.8
	12 - 17 Months	6,246	10.4
	18 - 23 Months	4,603	9.8
	24 - 35 Months	8,462	11.2
	36 - 59 Months	14,211	7.9
	Total	39,670	10.4

Page:



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 8 / 20 / 02

	Sum	mary of	Trends i	n Growt	h and Aı	nemia Ir	dicato	rs by I	Race/Et	hnicity ⁽³⁾			
Race/Ethnicity and Year		Bir	thweight (4))			Hei	ght and \	Veight			Aner	mia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	2 Yrs Overweight (6)			Low HI	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number	% 85th - <95th	% <u>></u> 95th	Number	%
White, Not Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	4,325 150 0 4,918 3,392 0 397 653 510 243	8.4 7.3 * 9.2 8.1 * 7.6 6.4 6.9 5.8	9.9 11.3 * 10.4 10.4 * 10.6 11.6 6.9 10.3	14,869 295 0 16,507 15,980 0 2,091 2,592 2,563 1,338	6.4 4.1 * 6.5 7.1 * 4.9 4.7 4.6 4.9	5.6 5.4 * 6.1 5.4 * 7.6 9.1 8.5 9.6	13.7 11.9 * 12.8 13.3 * 8.9 8.1 7.3 7.8	6,259 98 0 6,861 7,397 0 913 1,126 1,164 607	17.5 * 16.4 16.8 * 16.3 9.9 10.6 10.9	15.2 * 14.8 13.9 * 9.7 9.3 8.1 9.2	11,413 195 0 12,573 13,452 0 1,210 1,652 1,599 728	7.7 6.7 * 7.4 7.8 * 9.8 7.0 5.3 6.3
Black, Not Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	3,815 105 0 4,194 2,635 0 279 371 316 188	11.5 7.6 * 12.6 11.1 * 10.0 10.5 12.3 12.8	6.2 6.7 * 6.6 5.7 * 6.8 5.7 4.4 8.0	13,810 236 0 15,322 14,119 0 1,448 1,711 1,658 1,001	6.3 2.1 * 6.4 6.5 * 4.3 5.7 4.9 4.8	5.5 7.2 * 5.6 5.4 * 7.0 7.1 7.9 7.2	15.7 14.4 * 14.8 15.4 * 12.6 13.0 12.5 14.4	6,132 91 0 6,904 7,166 0 671 807 815 488	16.7 * 17.2 16.1 * 13.4 14.6 14.5 15.6	16.5 * 16.0 15.3 * 14.6 15.0 12.0 13.3	11,119 174 1 12,333 12,584 0 869 1,022 1,029 540	14.3 13.8 * 14.8 14.6 * 10.7 7.5 7.4 10.4



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: 2 Run Date: 8 / 20 / 02

	Sum	mary of	Trends ii	n Growt	h and Aı	nemia In	dicato	rs by F	Race/Et	hnicity ⁽³⁾			
Race/Ethnicity and Year		Biı	rthweight (4))			Hei	ght and \	Veight			Anen	nia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Low Hi	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number	% 85th - <95th	% <u>></u> 95th	Number	%
Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	5,078 100 0 4,504 2,491 0 291 400 365 181	7.5 6.0 * 9.7 8.9 * 8.9 7.8 13.2 7.2	7.6 9.0 * 6.7 7.1 * 8.2 6.3 6.3 6.6	19,205 267 2 18,204 16,771 0 2,064 2,229 2,131 1,095	6.0 3.7 * 7.1 7.1 * 4.6 4.3 4.6 5.8	4.5 5.6 * 4.3 3.6 * 5.5 5.3 5.3	19.2 15.0 * 18.4 19.2 * 13.5 13.4 13.6 15.4	8,850 121 0 8,815 9,282 0 1,041 1,130 1,060 529	17.7 20.7 * 19.5 18.4 * 15.0 17.3 14.5 15.3	23.1 21.5 * 21.9 19.9 * 15.2 14.3 15.7 15.9	15,470 206 3 14,897 15,380 0 1,290 1,538 1,404 627	9.7 12.6 * 10.2 10.4 * 8.0 7.4 5.6 9.7
American Indian/Alaskan Native	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	29 0 0 33 14 0 19 21 17	* * * * * * * * * *	* * * * * * * * *	90 0 0 98 119 0 63 76 62 29	* * * * 3.4 * * * * * *	* * * * * 7.6 * * * * *	* * * 9.2 * * * * *	36 0 0 46 81 0 27 31 27 13	* * * * * * * * * *	* * * * * * * * *	70 0 0 74 111 0 29 50 28 14	* * * 9.9 * * * *



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date:

Run Date: 8 / 20 / 02

Race/Ethnicity and Year		Bir	rthweight (4))			Hei	ght and V	Veight			Aner	mia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Low H	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>></u> 95th	Number	% 85th - <95th	% n <u>></u> 95th	Number	%
Asian/Pacific Islander	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	217 5 0 173 127 0 6 3 7	10.6 * 8.1 11.8 *	6.0 * 4.6 6.3 * *	608 7 0 622 624 0 17 12 17 6	6.7 * 6.9 4.8 * *	8.7 * 7.9 7.1 * *	12.7 * 13.8 11.5 * * *	222 2 0 268 299 0 7 2 2	14.0 * 15.7 14.4 * *	16.7 * 16.4 13.4 * *	446 3 0 482 532 0 9 7 5	9.6 * 9.8 9.6 * *
All Other/Unknown	2001 2000 1999 1998 1997 1996 1995 1994 1993	404 16 0 307 158 0 0 0	9.4 * 7.2 14.6 * *	7.7 * 8.5 13.3 * * *	1,451 24 0 1,027 757 0 0 0 0	5.7 * 6.1 8.1 * *	7.0 * 5.5 8.3 * * *	12.3 * 13.6 14.3 * *	585 7 0 378 286 0 0 0	17.4 * 17.5 14.0 * *	13.2 * 15.1 12.9 * * *	1,152 13 1 794 648 0 0 0 0	10.9 * 12.5 9.7 * *



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: 4 Run Date: 8 / 20 / 02

Race/Ethnicity and Year		Bir	thweight (4)				Hei	ght and V	Veight			Anen	nia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Low Hb/Hct (7)	
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% <u>≥</u> 95th	Number	% 85th - <95th	% ı <u>≥</u> 95th	Number	%
Total All Race/Ethnic Groups	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	13,868 376 0 14,129 8,817 0 992 1,448 1,215 623	9.0 7.4 * 10.3 9.4 * 8.9 7.9 10.1 8.2	7.9 9.0 * 8.0 8.1 * 8.9 8.6 6.0 8.7	50,033 829 2 51,780 48,370 0 5,683 6,620 6,431 3,469	6.2 3.3 * 6.7 6.9 * 4.7 4.8 4.7 5.2	5.2 5.9 * 5.3 4.9 * 6.6 7.4 7.4 7.5	16.3 13.6 * 15.4 15.9 * 11.5 11.2 10.7 12.2	22,084 319 0 23,272 24,511 0 2,659 3,096 3,068 1,639	17.3 17.9 * 17.8 17.1 * 15.0 13.8 12.9 13.8	18.7 16.6 * 17.8 16.6 * 13.2 12.7 11.8 12.6	39,670 591 5 41,153 42,707 0 3,407 4,269 4,065 1,911	10.4 10.7 * 10.8 10.8 * 9.4 7.3 6.0 8.7



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page:

Race/Ethnicity and Year				Brea	astfeeding (4)		
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
White, Not Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	4,325 150 0 4,918 3,392 0 365 574 462 227	44.7 48.0 * 43.1 42.5 * 38.6 38.7 37.2 30.0	2,510 70 4 2,785 2,927 0 129 229 185 130	20.2 * 21.9 17.3 * 16.3 17.0 15.1 12.3	2,368 48 0 2,304 2,510 0 97 156 131 105	15.7 * 16.5 12.7 * 8.3 9.2 8.6
Black, Not Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	3,815 105 0 4,194 2,635 0 224 267 257 164	43.0 45.7 * 37.9 34.6 * 23.2 23.2 21.4 15.9	2,551 57 0 2,777 2,624 0 41 69 73 65	23.1 * 20.1 15.8 * * *	2,150 40 0 2,273 2,264 0 19 40 46 40	16.8 * 15.0 10.0 * * *

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 2 Run Date: 8 / 20 / 02

CDC

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Sumn	nary of Trend	ds in Breas	stfeeding Inc	licators by	Race/Ethnici	ty ⁽³⁾	
Race/Ethnicity and Year				Brea	astfeeding (4)		
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Hispanic	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	5,078 101 0 4,504 2,491 0 245 331 312 157	56.4 45.5 * 45.4 42.0 * 37.1 34.4 34.3 31.2	3,296 62 1 2,870 2,862 0 84 159 123 93	24.6 * 24.4 18.6 * * 9.4 7.3 *	3,084 42 1 2,752 2,665 0 45 103 81 78	18.3 * 17.6 11.9 * 2.9
American Indian/Alaskan Native	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	29 0 0 33 14 0 15 15 16 5	* * * * * * * * * *	13 0 0 20 9 0 1 4 3	* * * * * * * * * *	15 0 0 12 11 0 0 2 1	* * * * * * * * *

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Page: 3 Run Date: 8 / 20 / 02

CDC

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Sı	ummary of Trend	ds in Breas	stfeeding Inc	licators by	Race/Ethnici	ty ⁽³⁾	
Race/Ethnicity and Year				Brea	astfeeding (4)		
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Asian/Pacific Islander	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	217 5 0 173 127 0 5 2 6 4	48.8 * 44.5 38.6 * *	125 1 0 102 135 0 0 1 0	18.4 * 28.4 22.2 * *	86 0 0 89 93 0 0 1	* * * * * * * *
All Other/Unknown	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	404 16 0 307 158 0 0 0	69.1 * 58.3 57.0 * * *	289 6 2 206 169 0 0 0	41.2 * 30.6 21.9 * *	274 1 0 155 152 0 0 0	31.8 * 21.3 21.1 * *

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C

2001 Pediatric Nutrition Surveillance (1) Connecticut

Page:

Run Date: 8 / 20 / 02

CDC

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3) Breastfeeding Race/Ethnicity and Year (4) Number % Number % Number % Ever Breastfed Breastfed Breastfed At Least At Least 6 Months 12 Months 8.784 23.4 Total All Race/Ethnic Groups 2001 13.868 49.3 7.977 17.7 2000 377 46.4 196 15.8 131 11.5 1999 0 7 1 1998 14.129 42.7 8.760 22.4 7.585 16.8 1997 8,817 40.2 8,726 17.5 7,695 11.8 0 1996 0 0 1995 854 34.4 255 10.2 161 6.8 1994 1,189 34.1 462 13.4 302 5.6 1993 1,053 33.0 384 11.2 259 6.2 1992 26.0 288 224 557 8.3 4.5

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 20C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Run Date: 8 / 20 / 02

Page:

Summary of Trends in Growth Indicators by Age Children Aged < 5 Years (2)

				S	Summar	y of Trer	nds in (Frowth Indicate	ors by	Age	3)				
Age and Year *			He	ight and	Weight			Age and Year *			He	eight and	Weight		
		Short Stature (4)	Under- weight (5)	Over- weight (5)		≥ 2 Yrs Overweight (5)				Short Stature (4)	Under- weight (5)	Over- weight (5)		≥ 2 Yrs Overweight (5)	
	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th		Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th
0 - 11 Months								24 - 35 Months							
2001 2000	17,237 378	6.8 2.9	7.5 6.6	11.7 10.1	- -	- -	-	2001 2000	8,294 93	4.8	5.1 *	16.4	8,294 93	15.7 *	16.4 *
1999 1998	1 17,801	* 6.8	* 8.1	* 10.8	_	-	-	1999 1998	0 8,408	* 6.0	* 4.4	* 16.3	0 8,408	* 16.2	* 16.3
1997	12,888	7.9	7.1	12.8	_	-	-	1997	8,676	5.7	4.4	15.0	8,676	16.3	15.0
1996 1995	0 1,945	* 4.0	* 10.5	* 7.2	-	- -	-	1996 1995	0 988	* 4.9	* 5.4	11.2	0 988	* 16.0	* 11.2
1994	2,276 2,137	3.6	11.8	7.4	-	-	-	1994	1,076 1,095	5.1	4.4	11.5	1,076 1,095	13.9	11.5
1993 1992	1,117	4.9 4.5	12.9 12.4	6.6 9.1	-	- -	-	1993 1992	626	3.9 5.1	4.9 4.6	12.2 12.6	626	12.7 13.3	12.2 12.6
12 - 23 Months								36 - 47 Months							
2001 2000	10,712 132	8.2 4.5	4.2 7.6	18.9 16.7	-	- -	-	2001 2000	6,971 72	5.1 *	3.4	19.7	6,971 72	18.7	19.7 *
1999 1998	1 10,707	* 7.8	* 4.4	* 17.6	-	-	-	1999 1998	0 7,408	* 6.5	* 3.4	* 18.9	0 7,408	* 18.4	* 18.9
1997	10,971	8.6	4.4	18.2	-	-	-	1997	7,712	5.7	3.4	17.7	7,712	17.3	17.7
1996 1995	0 1,079	5.1	4.8	15.3	-	-	-	1996 1995	0 914	5.5	3.9	14.4	0 914	14.3	14.4
1994 1993	1,248 1,226	6.7 4.8	6.3 4.2	14.5 15.2	-	-	-	1994 1993	1,059 1,104	5.7 4.6	4.9 5.3	12.2 12.0	1,059 1,104	14.2 13.3	12.2 12.0
1992	713	6.0	5.5	15.8	-	-	-	1992	589	6.8	5.4	12.9	589	13.6	12.9

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 3 years of age and height-for-age

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 20C 2001 Pediatric Nutrition Surveillance (1) Connecticut

Connecticut	
Summary of Trends in Growth Indicate	ors by Age
Children Aged < 5 Years	(2)

					S	umma	ary of Tren	ds in
Age and	Year *			He	ight and	Weight		
			Short	Under-	Over-		≥ 2 Yrs	
			Stature (4)	weight (5)	weight (5)		Overweight (5)	
		Number	%	%	%	Number	%	%
			<5th	<5th	≥95th		85th-<95th	≥95th
48 - 59 N	1onths							
	2001	6,819	4.5	2.9	20.4	6,819	17.9	20.4
	2000	154	2.6	3.9	16.2	154	18.2	16.2
	1999	0	*	*	*	0	*	*
	1998	7,456	5.5	2.9	18.5	7,456	19.1	18.5
	1997	8,123	5.4	3.6	17.2	8,123	17.8	17.2
	1996	0	*	*	*	0	*	*
	1995	757	4.5	4.1	14.1	757	14.7	14.1
	1994	961	4.2	4.4	14.7	961	13.2	14.7
	1993	869	4.9	4.1	11.2	869	12.7	11.2
	1992	424	3.5	5.0	12.3	424	15.1	12.3
Total								
	2001	50,033	6.2	5.2	16.3	22,084	17.3	18.7
	2000	829	3.3	5.9	13.6	319	17.9	16.6
	1999	2	*	*	*	0	*	*
	1998	51,780	6.7	5.3	15.4	23,272	17.8	17.8
	1997	48,370	6.9	4.9	15.9	24,511	17.1	16.6
	1996	0	*	*	*	0	*	*
	1995	5,683	4.7	6.6	11.5	2,659	15.0	13.2
	1994	6,620	4.8	7.4	11.2	3,096	13.8	12.7
	1993	6,431	4.7	7.4	10.7	3,068	12.9	11.8
	1992	3,469	5.2	7.5	12.2	1,639	13.8	12.6

ı Gı	owth Indicat	ors by	Age (3)				
	Age and Year *			He	ight and	Weight		
			Short Stature (4)	Under- weight (5)	Over- weight (5)		≥ 2 Yrs Overweight (5)	
		Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th

Page:



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 3 years of age and height-for-age

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 21C

2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Anemia by Age Children Aged < 5 Years (2)

Summary of Trends in Anemia by Age (3) Age and Year Age and Year Anemia Anemia Low Hb/Hct (4) Low Hb/Hct (4) % % Number Number 6 - 11 Months 18 - 23 Months 2001 6,148 15.8 2001 4,603 9.8 2000 125 14.4 2000 48 1999 1999 0 1 1998 6,522 16.1 1998 4,810 9.4 1997 6,595 15.6 1997 4,997 10.6 1996 0 1996 0 1995 642 12.9 1995 422 9.2 1994 778 9.3 1994 555 8.1 702 578 1993 10.8 1993 4.5 1992 340 14.4 1992 252 9.1 12 - 17 Months 24 - 35 Months 2001 6,246 10.4 2001 8,462 11.2 2000 85 2000 96 1999 2 1999 1 1998 6,029 11.1 1998 8,546 11.8 1997 1997 6,102 10.8 8.819 11.6 1996 0 1996 1995 354 10.7 1995 775 11.7 1994 441 9.1 1994 893 6.7 1993 385 8.1 1993 889 5.4 1992 213 7.0 1992 417 8.2



Page:

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 21C

2001 Pediatric Nutrition Surveillance (1) Connecticut

Summary of Trends in Anemia by Age Children Aged < 5 Years (2)

Page: 2 Run Date: 8 / 20 / 02

Summary of Trends in Anemia by Age (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	9
36 - 59 Months	2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	14,211 237 1 15,246 16,194 0 1,214 1,602 1,511 689	7.9 6.3 * 8.2 8.5 * 5.8 5.8 4.1 6.7			1	
Total All Age Gro	ups 2001 2000 1999 1998 1997 1996 1995 1994 1993 1992	39,670 591 5 41,153 42,707 0 3,407 4,269 4,065 1,911	10.4 10.7 * 10.8 10.8 * 9.4 7.3 6.0 8.7				

⁽¹⁾ Reporting period is January 1 through December 31.



⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.