

Estimates

- Between 2000 and 2013, among adult current smokers, a significant increase occurred in the prevalence of having tried to quit smoking cigarettes in the past 12 months (52.0%-67.0%). [Figure 1]
- During 2000-2013, among youth in high school who are current smokers, a slight increase occurred in the prevalence of having tried to quit (53.0%-55.5%). [Figure 1]
- Between 2000 and 2013, among current smokers, significant increases occurred in the percentage of adult males (52.0%-66.4%) and females (52.0%-67.7%) who tried to quit smoking cigarettes. [Figure 2]
- During 2000-2013, among youth in high school who are current smokers, an increase occurred in the prevalence of having tried to quit smoking cigarettes among males (48.7%-54.0%) and was relatively unchanged among females (57.0%-57.2%). [Figure 2]

Figure 1





Figure 2



Data Sources: Connecticut Behavioral Risk Factor Surveillance System (BRFSS); 2000-2013.
Connecticut Youth Tobacco Survey (YTS); 2000-2013.

For Further Information

Connecticut Department of Public Health
Tobacco Use Prevention and Control Program
410 Capitol Avenue
PO Box 340308, MS #11HLS
Hartford, CT 06134-0308
Phone: 860-509-8251
Or Visit: www.ct.gov/dph/tobacco