

From: noreply-dphimmunizations@ct.gov <noreply@everbridge.net>

Sent: Monday, November 1, 2021 2:18 PM

Subject: COVID-19 Vaccine Program (CoVP) Provider Bulletin, Week of November 1



DEPARTMENT OF PUBLIC HEALTH

Dear Connecticut COVID-19 Vaccine Providers,

This communication is being sent to all key contacts at provider organizations administering COVID-19 vaccine— please read this message in its entirety. Please feel free to share it with others in your organization who may benefit from the update. Note that all of our communications are archived on our [web site](#).

UPDATES

ACTION NEEDED: Transition from CT Vaccine Finder to vaccines.gov

On Wednesday, November 10, the State of Connecticut will be transitioning from [CT Vaccine Finder](#) to [Vaccines.gov](#) (powered by VaccineFinder), the Centers for Disease Control and Prevention's (CDC) one-stop landing page for the public to find a COVID-19 vaccination clinic. This transition will reduce the reporting burden for providers and will increase the ability for Connecticut residents to easily find a clinic near them. The public can also look for flu clinics on Vaccines.gov if [providers choose to report flu inventory](#).

As soon as possible and no later than Wednesday, November 10, we ask that all providers complete the below steps:

Step 1) Ensure that each of your COVID-19 vaccination clinics is registered with VaccineFinder.

- *Tip: As a reminder, only clinics that have registered PINs with VaccineFinder will appear on Vaccines.gov. After CoVP enrollment your Primary Vaccine Coordinator received an email from VaccineFinder@auth.castlight.health.com that invited you to create a COVID Locating Health Account. In this email, you can select either to 1) Report on behalf of all of your locations, or 2) Report individually. You can enroll a new location by following the steps [here](#). For support, email CARS_HelpDesk@cdc.gov or call 833-748-1979.*

Step 2) Complete daily inventory reporting to VaccineFinder for each of your COVID-19 vaccination clinics.

- *Tip: Daily inventory reporting is required, per the [CDC COVID-19 Provider Agreement](#). Inaccurate inventory on VaccineFinder may impact your future orders with the state. For a step-by-step guide to submitting daily inventory, [click here](#).*

Step 3) (Optional) Make your clinics visible on vaccines.gov.

- *Tip: For detailed information on how to display your vaccination clinics on Vaccines.gov, [click here](#). If you do not choose to make your locations visible or do not report daily inventory (Step 2), your locations will not appear on the one-stop landing page where residents can find COVID-19 vaccination clinics. Providers may choose not to display their clinic if they are not open to the public (ex. small pediatric clinic conducting their own promotion and only vaccinating their patients.). If you are open to the public, we strongly suggest that you display your clinics.*

Pediatric Pfizer vaccine updates and orders

On Friday, the Food and Drug Administration (FDA) amended Pfizer-BioNTech's Emergency Use Authorization (EUA) to allow the vaccine to be administered to children ages 5 to 11 years. FDA resources are available at:

- [Press Release](#)
- [Updated Pfizer Factsheets](#)

As a reminder, pediatric Pfizer vaccine “pre-orders” placed during the week of October 18th being shipped and are anticipated to arrive this week. **No pediatric vaccine should be administered prior to an Advisory Committee on Immunization Practices (ACIP) recommendation and CDC director endorsement.** The ACIP is scheduled to discuss pediatric COVID-19 vaccine recommendations on **November 2 from 10am–5pm**. The meeting can be livestreamed [here](#).

Providers who did not place a pre-order for pediatric Pfizer vaccine can still request doses by reaching out to DPH.Immunizations@ct.gov. The minimum order for direct shipment prior to November 9th is 300 doses. Connecticut DPH will arrange for redistribution for orders less than 300 doses during this period.

Starting November 2nd, pediatric Pfizer vaccine orders can be placed in the ordering portal alongside other COVID-19 vaccines. Orders that are placed in the portal will be filled after November 9th as part of our typical ordering cycle for delivery the week of November 15th. Pediatric Pfizer orders can be requested in multiples of 100 doses for direct shipment from the manufacturer. Connecticut DPH will arrange for redistribution for orders less than 100 doses, so if you cannot use 100 doses before they would expire, please enter any number of doses you would like in multiples of 10. Requests for all COVID-19 vaccine doses should be submitted through our [ordering portal](#) **by noon on Tuesdays**.

As a reminder, the current adolescent and adult product cannot be used for children under 12. Storage conditions for pediatric vaccine differ from the adult formulation. The pediatric Pfizer vaccine for 5–11 year olds can be stored in the refrigerator for up to 10 weeks and in ultra-cold freezers for up to 6 months. Please note that new Pfizer shippers will be used to support all direct ship pediatric orders placed on November 9th and onward. These **shippers should not be used for storage**. Vaccine should be removed and placed in 2°-8°C (for up to 10 weeks) or ultra-low temperature freezers at -80°C (for up to 6 months).

Supplemental Invitation to Bid

A Supplemental Invitation to Bid for COVID-19 Vaccination Services has been posted. To submit a bid, please visit the [CTsource Bid Board](#) and search for Solicitation #21PSX0155. Submissions are due on Wednesday, 11/3 at 12pm. If any questions, email Arlene Watson-Paulin: Arlene.Watson-Paulin@CT.Gov.

New and updated resources

- [Vaccine Administration Resource Library | CDC](#) Includes information Vaccine administration e-learn with CE for pharmacists, infographics for identifying injection sites, short video demonstration injection and holding children.
- [Pinkbook: Vaccine Administration | CDC](#) includes strategies to decrease anxiety and procedural pain.
- Clinical Considerations: Vaccinating Adolescents at [COVID-19 Vaccine Webinar Series | CDC](#) There will be a similar one for younger children posted soon.
- COVID-19 clinical materials, including product-specific trainings, guidance and infographics, are here: [COVID-19 Vaccination Clinical and Professional Resources | CDC](#)
- [COVID-19 Pfizer Pediatric Tris-sucrose Vaccine EUA Fact Sheet for Recipients](#)
- [COVID-19 Vaccines for Children and Teens](#)
- [Moderna](#), [Janssen](#), and [Pfizer](#) updated standing orders for booster doses
- [Quick Reference Guide](#)

Most Moderna and Janssen vaccine preparation, administration, and storage and handling materials have been updated by CDC and posted as they become available. All documents should be completed and posted by today.

- Centers for Disease Control and Prevention (CDC) [booster shot webpage](#) for the public
- Updated Advisory Committee on Immunization Practices (ACIP) [Clinical Considerations](#)
- COCA Call: [What Clinicians Need to Know about the Recent Updates to CDC's Recommendations for COVID-19 Boosters](#) (recording now available)
- [Different COVID-19 Vaccines](#)
- [COVID-19 Vaccines While Pregnant or Breastfeeding](#)

Information from Moderna about Booster Doses

- **Syringe Availability**

To assure providers have an adequate supply of 1 mL syringes to support extraction of booster doses (0.25 mL) from a Moderna 14 vial, CDC will begin shipping an additional Moderna 10 ancillary kit with all orders placed as of October 30. **When possible, please use 3 mL syringes for extraction of primary series doses to ensure you have an adequate supply of 1 mL syringes to support extraction of booster doses from a Moderna vial.** Do not puncture the vial stopper more than 20 times.

- **Training – Moderna COVID-19 Vaccine Booster Dose**

Join Moderna for a webinar for vaccination providers to learn more about the Moderna COVID-19 Vaccine booster dose. **Webinar:** Important updates on the mRNA-1273 50 µg Booster Dose

- Thursday, November 4th at 3pm ET – [Register here for Nov 4](#)
- Thursday, November 11th at 12pm ET – [Register here for Nov 11](#)

- **Moderna Letter to Providers**

Moderna mailed a letter and fact sheet to alert vaccination providers that the volume of the booster dose is **0.25mL**, half the dose provided in the primary series administered.

The letter and fact sheet that were sent to providers can also be found at [Moderna COVID-19 Vaccine DHCP Letter-Fact Sheet 10.20.21.pdf](#).

Moderna Inventory and Adjustments in CT WiZ

- All Moderna orders approved after 10/22 will receive double ancillary kit shipments with each order to support administration of booster doses at a “half dose” strength
- **Terminology** – Moderna products should continue to be referenced as “Moderna 10” and “Moderna 14”.
 - Moderna 10 – *is not currently available for ordering.*
 - Moderna 14 – most inventory will be this product.
- Primary (0.5mL) and booster (0.25 mL) doses of Moderna are administered from the same multi-dose vial for 10 or 14 doses.
 - Therefore, you will need to adjust your inventory if you administer extra doses per a 10 or 14 dose vial.
 - Your inventory on-hand will need to be adjusted per vial based on how many additional doses were administered. Vaccine wastage will also need to be made per vial. Wastage is reported in whole doses.
- **How to make an inventory adjustment in CT WiZ:**
 - *Tip:* When administering vaccines, do not forget to track which type of dose (primary or booster) was extracted from each vial.
 - *Example:* You have a 14-dose vial and administered 20 doses (*because you can only puncture the cap a maximum of 20 times*), you make an inventory adjustment of 10 extra doses using the Reason code below and enter the Doses Adjustment of 4 whole doses wasted.
 - In CT WiZ, select Inventory – Vaccines – On-Hand – select Actions – select Adjustment:
 - select the Reason code: **Dose Count Variance Multi-Dose Vial** and

- enter the **Doses Adjusted** (*as whole doses.*)
- **For User Interface: How to document correct booster dosage administered in CT WiZ:**
 - In CT WiZ, find the patient – select Action – Add Vaccine – Create And Administer – Change **Dosage field to 25**
- **For EHR vendors and senders: please be sure to include the correct dosage volume** (primary 0.5mL or add in the booster 0.25 mL) for Moderna in the HL7 message to be reported from your Electronic Health Record (EHR) system to CT WiZ.

REMINDERS

We are keeping information on many important reminders from past communications in the space below. To prevent this section from becoming too lengthy, what remains is selected carefully. All past communications are available [here](#).

Booster Doses

There are now booster recommendations for all three COVID vaccines. All providers should actively reach out to patients to whom they gave first doses and encourage them to receive booster doses if eligible. All individuals 65 and older should receive a booster; Everyone who received J&J should receive a booster; and many individuals 18-64 who received an mRNA vaccine are likewise eligible for a booster. Please read below for details.

For individuals who received an mRNA vaccine (Pfizer-BioNTech or Moderna)

For individuals who received a Pfizer-BioNTech or Moderna COVID-19 vaccine, the following groups are eligible for a booster shot at 6 months or more after their initial series:

- 65 years and older
- Age 18+ who live in [long-term care settings](#)
- Age 18+ who have [underlying medical conditions](#)
- Age 18+ who work or live in [high-risk settings](#)

For individuals who received a J&J vaccine

For individuals who received a J&J vaccine, booster shots are recommended for those who are 18 and older and who were vaccinated two or more months ago. This means that all individuals who received a J&J vaccine should receive a booster.

Mixing & Matching (heterologous series)

Both the Food and Drug Administration (FDA) and CDC support individuals to receive a booster dose that is a different vaccine type than they originally received for their primary series if they

choose. CDC's recommendations now allow for this type of mix and match dosing for booster shots.

Pfizer-BioNTech 5 through 11 year old product introduction

What is currently known about COVID-19 vaccines for children <12 years old:

- The Pfizer-BioNTech COVID-19 Vaccine for 5–11-year-olds will be a new product with new packaging (orange cap, see attached PRELIMINARY reference document for more details) and a new national drug code (NDC). ***The current product for adults and adolescents should not be used in children.***
- The new product configuration will be 10-dose vials, in packages of 10 vials (100 dose total) pending FDA authorization. The product can be stored for 10 weeks at 2 to 8°C in the refrigerator. There will also be changes to the product shipper.
- The Immunization Program will support the transfer of vaccines in smaller quantities than is available for order if shipped directly from the manufacturer.
- COVID-19 pediatric vaccines will require diluent, and this will be provided with ancillary supplies which are configured specific to new vaccine packaging and appropriate for use in children.
- The Public Readiness and Emergency Preparedness (PREP) Act has expanded scope of practice for pharmacists and pharmacy technicians nationwide to allow for provision of vaccinations to children ≥3 years old (please see: <https://www.hhs.gov/sites/default/files/prep-act-guidance.pdf>).
 - Pharmacists, pharmacy interns, and qualified pharmacy technicians (if the pharmacy intern or technician acts under the supervision of such pharmacist and the pharmacy intern or technician is licensed or registered by his or her State board of pharmacy), may administer vaccines that the (ACIP) recommends to persons ages three through 18 according to ACIP's standard immunization schedule. This includes influenza and FDA authorized or FDA licensed COVID -19 vaccines.
 - The PREP Act Fact Sheet explaining the COVID-19 vaccination workforce can be found [here](#). The Fact Sheet described qualified persons covered to administer **COVID-19 vaccines** under the PREP Act Declaration and its amendments.
- The months of November and December have multiple holidays and potential for winter weather. Providers should factor this into their inventory management and clinical operations.

Assumptions regarding the COVID-19 vaccine program for children <12 years old:

- We should be ready to vaccinate children 5 through 11 years old shortly after the VRBPAC meeting on Oct 26th pending FDA authorization and ACIP recommendations.
- Pharmacies participating in the Federal Retail Pharmacy Program (FRPP) can order vaccine to select pharmacy locations increasing the number of locations children may go to get vaccinated.

- The CT Immunization Program continues to promote enrollment of CT Vaccine Program providers (CT’s pediatric vaccine program) to be COVID-19 Vaccine Providers.

Moderna Inventory Shelf Life Extension

- **Shelf-life extension:** Moderna continues to update data regarding extension of expired doses. A total of 78 lots have been approved for extension, and more may follow. These batches of Moderna vaccine have been processed for expiry extension and provide an additional 2 months. The product **must be maintained in the frozen state for the extension to be** applicable (extension does not apply to thawed vials). Moderna’s [look up tool](#) remains the official indicator of product expiry.

Recent COVID-19 and vaccine related publications

- [COVID-19 Vaccination and Non-COVID-19 Mortality Risk — Seven Integrated Health Care Organizations, United States, December 14, 2020–July 31, 2021](#)
- [Effectiveness of Pfizer-BioNTech mRNA Vaccination Against COVID-19 Hospitalization Among Persons Aged 12–18 Years — United States, June–September 2021](#)

Pfizer Vaccines Host “Medical Updates”

- These sessions are hosted by Pfizer and will be continuously updated to reflect new information and changes that evolve.
- Session topics, subject to change, may include: FDA indication & authorizations; CDC / ACIP recommendations; Packaging / presentation updates; Storage, handling & administration; Test your knowledge (Q&A scenarios for various storage & expiry conditions)
- Please click on the links below to join the sessions at the designated times. These sessions will be **updated** to reflect new information and changes that evolve. Such updates will be identified at the start of each session and further explained during each presentation.

Please click on the links below to join the sessions at the designated times. *

Date & Time	Password
Attendee link – November 1 – 5 PM ET	g9ZmgHaip32
Attendee link – November 2 – 5 PM ET	sJDZQERp325
Attendee link – November 3 – 12 PM ET	82qdN3PppPp
Attendee link – November 4 – 12 PM ET	Y4ZkXdh2bz7
Attendee link – November 5 – 12 PM ET	rJSpNpts332

[Johnson & Johnson’s Janssen COVID-19 Vaccine Overview and Safety | CDC](#)

Johnson & Johnson's Janssen (J&J/Janssen) COVID-19 Vaccine: Women younger than 50 years old should especially be aware of the rare risk of blood clots with low platelets after vaccination. There are other COVID-19 vaccines available for which this risk has not been seen. If you received a J&J/Janssen COVID-19 Vaccine, [here is what you need to know](#). Read the [CDC/FDA statement](#).

Guidance regarding eligibility verification for booster doses of COVID-19 vaccines

As has always been the case during the COVID-19 vaccine roll out, the State of Connecticut wants to ensure that vaccine access is simple and straightforward for vaccine recipients. Providers should continue to ensure that COVID vaccines are readily available for individuals who seek them and that administration of a COVID-19 vaccine is medically appropriate and conducted in line with the ACIP guidelines.

Ensuring patients receive a COVID vaccine at the appropriate interval:

Connecticut DPH recommends that providers have a procedure for ensuring that individuals receive their booster no sooner than the minimum recommended interval. This can be accomplished in one of several ways, in order of preference:

- Verify the vaccine information, including the brand and date(s) a patient received their previous COVID-19 vaccine doses using CT WiZ; or
- Review the Centers for Disease Control and Prevention (CDC) Vaccination Card, VAMS Certificate, CT WiZ Certificate, or other documentation of the recipient's vaccination record; or
- If neither of the above are possible, ask the patient for the vaccine brand(s) and dates of previous COVID-19 doses received during the scheduling process (and including logic in scheduling systems to ensure appropriate interval).

Ensuring recommended individuals receive a COVID booster:

Connecticut DPH recognizes that the categories of individuals for whom boosters may be confusing or not immediately clear to all patients. Connecticut DPH continues to support patient self-identification and attestation as a sufficient basis for confirming that an individual is within one of the categories for whom boosters are recommended.

Additional considerations:

DPH reminds providers that there is no residency requirement to receive COVID-19 vaccines and that identification (ID) is likewise not required. Although ID may be requested from the patient for purposes of billing, no individual should be turned away due to lack of ID and providers should endeavor to create a welcoming environment for people to seek COVID-19 vaccines.

Vaccine Lot Management and Expiration

In order to minimize the number of unused expired doses and manage expired doses correctly, we encourage providers to:

- Monitor expiration dates weekly, rotate stock as needed, and follow a "first in, first out" strategy to manage inventory.

- If nearing expiration, check posted manufacturer information for the most up to date expiration/extension information for vaccine lots.
- Based on the latest expiration information and unless instructed to do otherwise, REMOVE expired vaccine from the storage unit. Do not give staff opportunity to administer expired vaccine.
 - If expired vaccine is inadvertently is administered, it is considered a vaccine administration error and requires remediation including a VAERS report, contacting the recipient to inform them of the error, and may or may not require revaccination based on the manufacturers' guidance. Guidance on vaccine administration errors can be found in Appendix A of the [Interim Clinical Considerations for Use of COVID-19 Vaccines Currently Authorized in the United States](#).
- Vaccine disposal: dispose of the vaccine vial (with any remaining vaccine) and packaging as medical waste. Do NOT return vaccine in the thermal shipping container.
- Check your vaccine stock for lots expiring using the [CDC's Vaccine Lot Number and Expiration Date webpage](#).
- Any vial of J&J that has a date prior to September 23, 2021 has now expired. There will be no more extension. J&J has an expiration date look up [website](#).
- Request access to a new COVID-19 Vaccine Lot Number report via CDC's Vaccine Code Set Management Service (VCSMS). This report includes COVID-19 vaccine lot numbers and expiration dates provided to CDC by the vaccine manufacturers. This report is updated daily and can be used to support vaccine administration, inventory management, and jurisdiction IISs. Complete the registration form on [CDC's Vaccine Lot Number and Expiration Date webpage](#) to request access to the report.

Providers should dispose of expired vaccine appropriately and report the wastage in the DPH ticketing system. When reporting expired doses to the [Helpdesk](#), please select the "Report Vaccine Wastage" ticket option, then select "Other" as the wastage reason and type "Expiration" in the text box when prompted. Please make sure that inventory is updated in Vaccine Finder, VAMS, or CT WiZ as appropriate.

Communicating the Benefits of Influenza Vaccine during COVID-19 Please see new handout which includes updated recommendations on co-administration of flu and COVID-19 vaccines and tips for discussing flu vaccine with patients. Check it out [here](#).

Influenza Corner

The Advisory Committee on Immunization Practices *Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices, United States, 2021–22 Influenza Season* is available [here](#).

If a patient is eligible, both the flu and COVID-19 vaccines can be administered at the same visit, [as recommended by CDC and its Advisory Committee on Immunizations Practices \(ACIP\)](#).

In addition to flu vaccine, the COVID-19 vaccine can be given with other vaccines as well. Even though both vaccines can be given at the same visit, providers should follow the recommended schedule for either vaccine: If your patients haven't gotten the currently recommended doses of COVID-19 vaccine, they should get a COVID-19 vaccine as soon as possible, and ideally they should get a flu vaccine by the end of October.

Giving all vaccines for which a person is eligible at the same visit is considered a [best practice](#) as it increases the probability people will be up to date on recommended vaccines. It also is an important part of immunization practice, especially if a health care provider is uncertain that a patient will return for additional doses of vaccine.

Coadministration of COVID-19 and Influenza vaccines

You may administer COVID-19 and influenza vaccines without regard to timing (both live, attenuated and non-live influenza vaccines). This includes administration of COVID-19 and influenza vaccines on the same day, as well as coadministration at any time interval. ([CDC | Interim Clinical Considerations: Coadministration of COVID-19 and Influenza vaccines](#))

With influenza season approaching, there may be compelling logistical advantages to offering patients COVID-19 and influenza vaccines on the same day, and you may encourage patients to receive these on the same day. There are no safety concerns for coadministration.

Best practices for [administering more than one vaccine](#), including COVID-19 vaccines and influenza vaccines, include:

- When preparing more than one vaccine, label each with the name and dosage (amount) of vaccine, lot number, the initials of the preparer, and the exact beyond-use time, if applicable.
- Always inject vaccines into different injection sites.
 - Separate injection sites by 1 inch or more, if possible so that any local reactions can be differentiated.
 - Each muscle (deltoid, vastus lateralis) has multiple injection sites ([Vaccine Administration Route and Site](#)).
- If administered at the same time, COVID-19 vaccines and vaccines that might be more likely to cause a local injection site reaction (for example, [high-dose and adjuvanted inactivated influenza vaccines](#)) should be administered in different limbs, if possible ([Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices \(ACIP\)–United States, 2021-22](#)).
- Inject vaccines rapidly without aspiration since aspiration is not recommended before administering a vaccine ([Vaccine Administration Route and Site](#)).
- There are many existing resources on administration and co-administration of vaccines relevant for healthcare providers, including:
 - [Pink Book: Vaccine Administration | CDC](#)
 - [You Call The Shots: Vaccinating Adolescents](#) (cdc.gov)
 - [Vaccine Administration Training Module](#) (cdc.gov)
 - [Intramuscular \(IM\) Injection: Sites](#) - YouTube
 - CDC also is developing a job aid and video for health care professionals on coadministration of vaccines.

Ensure Best Practice by Checking CT WiZ Before Vaccinating

Prior to administering any vaccine, including the COVID-19 vaccine, it is important that a recipient's prior vaccine history be referenced in the Immunization Information System (CT WiZ); best clinical practice is to verify that the recommended interval has elapsed since a previous dose of vaccine and that the preferred brand of vaccine be administered.

- If your clinic established bi-directional data exchange, you can query CT WiZ for the patient immunization record [electronically from your EHR](#).
- Every CoVP enrolled clinic can [look up the patient immunization record](#) in the [CT WiZ user interface](#). Those who signed the CoVP Provider Agreement and the primary and back-up vaccine coordinator already have CT WiZ access.
- If your staff need access to CT WiZ, [request a username](#) and select the 'Clinic Access' role. Access to CT WiZ is defined in [EXECUTIVE ORDER NO. 13C](#).
- If you need support, please submit a [Helpdesk ticket](#).

Report vaccine temperature excursions and vaccines wasted to the [Helpdesk](#) by following the prompts and submitting a ticket.

- Immediately report all temperature excursions in storage units containing COVID-19 vaccines. Staff will be notified in real time of the excursion and will assist you. Please store vaccines in the storage unit in a paper bag marked "do not use" until a determination about the viability of the vaccine has been made, in consultation with the manufacturer and the Immunization Program.
- Also report to the Helpdesk all COVID-19 vaccines deemed wasted or expired. This includes Pfizer vials from which a sixth dose cannot be extracted.

Post Notice to Patients About Reporting to CT WiZ at CoVP Clinics

All clinics enrolled in the CoVP must post the [Notice to Patients About Reporting to CT WiZ](#), to let vaccine recipients know their records are being reported to CT WiZ. If you are reporting through VAMS, this data is transmitted to CT WiZ.

Replacement CDC COVID Vaccination Cards

- *Recommendation:* Post your procedure for recipients to request replacement cards, in a location visible to recipients at your clinic as well as on your website. If your clinic needs additional CDC COVID-19 Vaccination Cards, contact immunizations@ct.gov and we will email you a pdf of the card for your clinic to print (*note: these should not be posted on your website*). Large type print cards are also available via pdf.
- *Reminder:* You should issue CDC COVID-19 Vaccination Cards to your vaccine recipients. DPH cannot issue a CDC card to recipients. DPH can provide the immunization record from CT WiZ. DPH's procedure is on our COVID-19 Vaccine and DPH Immunization webpages at: [I lost my vaccine card, how do I get another one? \(ct.gov\)](#) and [Request an Official Immunization Record](#).

Upcoming Clinic Trainings and Office Hours

- **CoVP Update -- next meeting November 3rd**
Provides CoVP Vaccinating Providers updates on the CoVP Program and Q&A with DPH staff. **Every other Wednesday** 9:00am-10:00am; Join: [CoVP Office Hours](#)
- The **CoVP Enrollment Office Hours** Provides information about the CoVP enrollment process in CT WiZ and Q&A with DPH staff. **Mondays** 12:00pm-12:30pm; Join: [CoVP CT WiZ Enrollment Office Hours](#)

- **Vaccine Storage and Handling Assessment** For clinics who completed CoVP enrollment and received an invite to attend. **Tuesdays** 10:00am-11:00am & **Thursdays** 1:00pm-2:00pm; **By invitation Only**
- **VAMS & CT WiZ Live Helpdesk Office Hours** Clinics can ask questions about VAMS and CT WiZ functionality to DPH staff. **Tuesdays** 9:00am 10:00am;Join: [VAMS Live Helpdesk Office Hours](#)

Visit: [VAMS Training](#) and [CT WiZ Training](#) for enhancement release notes and training.

Thank you for all of your ongoing work and support of our COVID-19 vaccine roll-out in Connecticut.

CT DPH Immunization Program Main P: (860) 509-7929 | Fax: (860) 706-5429 | Email: dph.immunizations@ct.gov Website: [CT Immunization Program](#) | [COVID-19 Vaccine Program](#)

If you would like to unsubscribe from these communications, please send an email to Dph.immunizations@ct.gov with the subject line "Unsubscribe from COVID-19 Program communications".