

**Chlamydia, Gonorrhea, and Primary and Secondary Syphilis Cases by Year of Diagnosis  
Connecticut, 2007 - 2014**

Year	Disease					
	Chlamydia		Gonorrhea		Syphilis	
	No.	Rate*	No.	Rate*	No.	Rate*
2007	11,510	326	2,332	66	39	1
2008	12,448	351	2,793	79	33	1
2009	12,144	341	2,558	72	65	2
2010	12,692	355	2,568	72	99	3
2011	13,662	381	2,450	68	67	2
2012	13,521	376	2,197	61	81	2
2013	12,962	360	2,935	82	53	1
2014	13,134	365	2,223	62	87	2

Source: Connecticut STD Control Program, STD\*MIS Data

Source of population data: CDC Bridged-Race Population Estimates, United States July 1st resident population by state, county, age, sex, bridged-race, and Hispanic origin. Compiled from revised bridged-race 2000-2009 intercensal population estimates (released by NCHS on 7/26/2012); and bridged-race Vintage 2014 (2010-2014) postcensal population estimates (released by NCHS on 10/26/2012); and bridged-race vintage (2010-2014) postcensal population estimates (released by NCHS on 6/30/2015). Available on CDC WONDER Online Database. Accessed at <http://wonder.cdc.gov/bridged-race-v2014.html> on Aug 17, 2015

\*Rates = per 100,000 population. Rates based on counts <20 may be unstable and should be interpreted with caution.