

# STATE OF CONNECTICUT

## DEPARTMENT OF PUBLIC HEALTH

Renée D. Coleman-Mitchell, MPH  
Commissioner



Ned Lamont  
Governor  
Susan Bysiewicz  
Lt. Governor

### HEALTHCARE QUALITY AND SAFETY BRANCH

### BLAST FAX 2019-09

**TO:** Health Care Facilities

**FROM:** Donna Ortelle, R.N., M.S.N.,  
Section Chief  
Facility Licensing and Investigations Section

**DATE:** June 28, 2019

**SUBJECT:** Recommendations for Management of Health Facilities during Hot Weather

---

In anticipation for significant hot weather, please review your emergency plans, check your generators and its fuel supply, ensure the completeness of emergency food and other supplies, and as always, check with your local weather channel via television and radio.

These weather conditions can impact cooling systems and overall patient/resident health. Please see the attached recommendations for management of patients/residents during hot weather. While the attachment is a guidance document that was issued in July of 2010, it still remains very relevant.

Please call (860) 509-7407 should you have any additional questions.



Phone: (860) 509-7400 • Fax: (860) 509-7543  
Telecommunications Relay Service 7-1-1  
410 Capitol Avenue, P.O. Box 340308  
Hartford, Connecticut 06134-0308  
[www.ct.gov/dph](http://www.ct.gov/dph)

*Affirmative Action/Equal Opportunity Employer*



1997-1998

## REVENUE

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998

1997-1998


1997-1998

1997-1998

1997-1998

1997-1998

# DPH UPDATE



Wednesday, July 07, 2010

Connecticut Department of Public Health  
(860) 509-7270

## Recommendations for Management of Nursing Home Residents During Hot Weather

- Baseline assessment of all residents (some residents tolerate the heat better than others). Make particular note of residents with ongoing febrile illnesses, as well as those subject to excessive fluid loss (e.g., diarrhea, vomiting, open wounds).
- Regular, more frequent assessment of residents at risk (e.g. cardiovascular or respiratory disease, neurologic conditions that affect the temperature regulating mechanism, those who cannot communicate their thirst).
- Notify facility medical director. Maintain a roster of residents "at risk" and report on status regularly to medical director.
- Monitor and document air temperatures in various parts of the building at regular intervals.
- Ensure adequate fluids for each individual resident, as well as make fluids available for staff. Increase frequency of "rounds" to encourage resident consumption of fluids; set up "water stations" throughout the facility; offer various forms of fluids (e.g. popsicles, watermelon).
- Initiate and monitor Intake and Output on patients with risk factors/diagnoses and those whose intake is poor. Daily weights may also be appropriate.
- Ensure a sufficient and safe supply of fans to circulate air.
- Evaluate resident's clothing needs, especially those cognitively unable to evaluate own needs.
- Monitor residents' temperatures and provide cool sponge baths.
- If residents choose to go outside, monitor carefully for heat-related symptoms and identify those residents who may be prone to heat-related problems or photosensitivity due to medications. Encourage residents to sit in shaded outdoor areas; apply sunscreen as needed, unless resident is allergic to the product.
- Encourage residents to sit in areas of the facility that may be air-conditioned.

**DPH Update**

**Wednesday, July 07, 2010**

**Page 2 of 3**

- Prior to predicted heat waves, check air conditioning systems and supplies. Ensure that facility equipment maintenance contracts are current, as well as emergency call list for rental companies (e.g. portable air conditioning units).
- If air conditioning is available, provide for regular maintenance. If air conditioning problems develop, alert corporate office and/or local building authorities as appropriate. Communicate status of repairs to residents and families by posting signs in the facility. Incorporate heat-related events in the facility's Disaster Plan.
- Maintenance staff should make regular rounds and monitor building systems throughout the period of hot weather (e.g. overloaded electrical circuits, open windows). Documentation of monitoring efforts, findings and interventions should be maintained.
- Notify DPH regarding issues of mechanical failures and the measures implemented by the facility. DPH may call the nurse in charge to ascertain implementation of appropriate interventions, status of residents, and ambient temperatures.

**Comparison of Heatstroke and Heat Exhaustion**

<b>Heatstroke</b>	<b>Heat Exhaustion</b>
<b>Definition</b> A condition or derangement of the thermo-regulatory center due to exposure to the rays of the sun or very high temperatures. Loss of body heat is inadequate or absent.	<b>Definition</b> A state of definite weakness produced by the excess loss of normal fluids and sodium chloride in the form of sweat.
<b>History</b> Exposure to high environmental temperature; use of medications that increase heat production or inhibit perspiration.	<b>History</b> Exposure to heat, usually indoors
<b>Differential Symptoms</b> Face: Red, dry, and hot Skin: Hot, dry, and no sweating Temperature: High, 106° to 110°F (41.1° to 43.3°C) Pulse: Full, rapid, strong, bounding Respirations: Dyspneic, fast, sonorous Muscles: Tense and possible convulsions Eyes: Pupils are dilated but equal	<b>Differential Symptoms</b> Face: Pale, cool, and moist Skin: Cool, clammy, with profuse diaphoresis Temperature: Usually not above 100°F (37.8°C) Pulse: Weak, thready, and rapid Respirations: Shallow and quiet Muscles: Tense and contracted Eyes: Pupils are normal; eyeballs may be soft

**DPH Update**

**Wednesday, July 07, 2010**

**Page 3 of 3**

Treatment	Treatment
Absolute rest with head elevated; keep body cool by any means available until hospitalized, but do not use alcohol applied to skin. Take temperature every 10 minutes, and do not allow it to fall below 101°F (38.5°C) to prevent hypothermia. Drugs: Allow no stimulants; give infusions of normal saline (to force fluids).	Keep patient quiet; head should be lowered to prevent orthostatic hypotension; keep body warm to prevent onset of shock.  Drugs: Salty fluids and fruit juices should be given frequently in small amounts. Intravenous isotonic saline will be required if patient is unconscious.

**Source:** Taber's Cyclopedic Medical Dictionary,  
18<sup>th</sup> Edition

- If the above conditions are noted, monitor Intake and Output and administer oxygen if ordered by the physician.

###

