




Keeping Connecticut Healthy

Hospital Performance Comparisons, 2006

A REPORT ON QUALITY OF CARE IN CONNECTICUT HOSPITALS

Joan Foland, M.Phil, M.H.S.

Lloyd Mueller, Ph.D.



State of Connecticut
Department of Public Health
Planning Branch
Health Information, Surveillance, and Reporting Section
Health Care Quality, Statistics, Analysis, and Reporting Unit

April 2008

Hospital Performance Comparisons, 2006

A REPORT ON QUALITY OF CARE IN CONNECTICUT HOSPITALS

Table of Contents

	<i>Page</i>
Executive Summary.....	i
Introduction.....	1
Background	1
Hospital Quality of Care	3
Hospital Performance Comparisons [with Figures].....	7
Quality of Care Measures for Heart Attack Patients.....	19
Quality of Care Measures for Heart Failure Patients	22
Quality of Care Measures for Pneumonia Patients	24
Discussion.....	26
References.....	31
Resources.....	32
Acknowledgments.....	33
Connecticut Hospitals	34
Appendices	36
A. Definitions of Measures	
B. Tables	
2005 Heart Attack, Heart Failure, and Pneumonia Performance Rates	
2006 Heart Attack, Heart Failure, and Pneumonia Performance Rates	
Performance Rates with Significant Differences between 2005 and 2006	

Executive Summary

Increasing attention is being focused on evaluating and improving health care quality at both the state and national levels. Efforts are being made to provide standardized, useful and valid information to the public about hospital quality of care and also to promote quality improvement efforts within hospitals. The incentives are clear-- high quality care leads to fewer repeat hospitalizations, medical procedures, and medical errors, thereby reducing costs. Results presented in this report constitute an initial step in this ongoing process to evaluate and report on health care quality in Connecticut hospitals.

Connecticut's initiative began with the passage of legislation (Sections 19a-127 l-n of the Connecticut General Statutes) during the spring of 2002 that created a quality of care program within the Department of Public Health (DPH). Under that program, hospitals are required to collect and report quality of care information to the DPH in order to produce a public report that compares all licensed hospitals in the state. Connecticut has aligned its efforts with national quality initiatives aimed at collecting similar information.

Included in this report are comparisons among adult general acute-care hospitals in Connecticut about how often they provide the recommended care to patients who have been diagnosed with a heart attack, heart failure, or pneumonia, which are three common and costly medical conditions for which people go to the hospital. Hospital performance rates are provided for ten clinical measures that focus on treatments that are well established and generally accepted recommended care based on medical evidence.

Based upon 2006 hospitalization data, Connecticut hospitals' average performance rates exceed national averages on all ten of the clinical measures, and are significantly better on nine of the ten measures. Hospitals can achieve high levels of performance (Figure A). However, most hospitals still fall short of the goal of 100% on most of the measures. That is, performance gaps still exist between the care that could be given and the care that is being delivered.

Performance rates are improving, however. Between 2005 and 2006, Connecticut hospitals' performances rates improved statistically significantly for all ten measures.

Connecticut's Performance Compared to the U.S. Performance, 2006

Condition	Measure	Range in Connecticut 2006	Average	Average
			Connecticut Rate 2006	National Rate** 2006
Heart Attack	Aspirin at Arrival	89% - 100%	98%*	93%
	Aspirin at Discharge	88% - 100%	98%*	90%
	ACEI or ARB for LVSD at Discharge	68% - 100%	85%	83%
	Beta-Blocker at Discharge	91% - 100%	97%*	90%
	Beta-Blocker at Arrival	85% - 100%	95%*	87%
Heart Failure	LVF Assessment	84% - 100%	96%*	84%
	ACEI or ARB for LVSD at Discharge	74% - 100%	86%*	82%
Pneumonia	Oxygenation Assessment	99% - 100%	100%*	99%
	Pneumococcal Vaccination	10% - 96%	79%*	71%
	Timely Antibiotic	68% - 94%	81%*	80%

** Source: www.hospitalcompare.hhs.gov for hospitals participating in the Hospital Quality Alliance initiative.

Data are based upon patients hospitalized from 1/1/06 – 12/31/06.

* Difference is statistically significant ($p < 0.05$).

Figure A



Hospitals were excluded from a measure if they had fewer than 20 cases.

Connecticut's Performance between 2005 and 2006

Condition	Measure	2005	2006
Heart Attack	Aspirin at Arrival	96%	98%*
	Aspirin at Discharge	97%	98%*
	ACEI or ARB for LVSD at Discharge	82%	85%*
	Beta-Blocker at Discharge	96%	97%*
	Beta-Blocker at Arrival	94%	95%*
Heart Failure	LVF Assessment	95%	96%*
	ACEI or ARB for LVSD at Discharge	83%	86%*
Pneumonia	Oxygenation Assessment	100%	100%*#
	Pneumococcal Vaccination	67%	79%*
	Timely Antibiotic	79%	81%*

* Difference is statistically significant ($p < 0.05$).

Oxygenation Assessment for Pneumonia increased from 99.8% in 2005 to 99.9% in 2006 ($p = 0.02$).

Consumers should view this information as a starting point for educating themselves about hospital quality, for talking to their doctors about choosing a hospital for medical care, and for asking questions while receiving care in the hospital. This information should also be used by the medical community to heighten their awareness of the opportunity that exists to improve the care that they currently deliver.

Hospital Performance Comparisons: A Report on Quality of Care in Connecticut Hospitals

INTRODUCTION

Increasing attention is being focused on evaluating and improving health care quality at both the state and national levels. Efforts are being made to provide standardized, useful and valid information to the public about hospital quality of care and also to promote quality improvement efforts within hospitals. The incentives are clear-- high quality care leads to fewer repeat hospitalizations, medical procedures, and medical errors, thereby reducing costs. Results presented in this report constitute an initial step in this ongoing process to evaluate and report on health care quality in Connecticut hospitals. Included in this report are comparisons among adult general acute-care hospitals in Connecticut about how often they provide the recommended care to patients who have been diagnosed with a heart attack, heart failure, or pneumonia, which are three common and costly medical conditions for which people go to the hospital. Consumers should view this information as a starting point for educating themselves about hospital quality, for talking to their doctors about choosing a hospital for medical care, and for asking questions while receiving care in the hospital. This information should also be used by the medical community to heighten their awareness of the opportunity that exists to improve the care that they currently deliver.

The hospital quality measures in this report come from information collected on patients who were discharged from Connecticut's hospitals during the time period January 1, 2006 through December 31, 2006.

BACKGROUND

What is the impetus to improve quality?

Three landmark reports issued by the Institute of Medicine (IOM), a congressionally chartered advisory group to the federal government, have brought much attention to the problems regarding the quality and safety of health care. The first report, *To Err is Human: Building a Safer Health System*¹ revealed the extent to which medical errors cause harm to patients in hospitals, and it set forth a national agenda for improving patient safety. The second report, *Crossing the Quality Chasm: A New Health System for the 21st Century*² found that problems in the health care delivery system are the source of many errors and recommended that the Department of Health and Human Services identify a few areas for focused quality measurement and improvement. The latest report, *Leadership by Example: Coordinating Government*

*Roles in Improving Health Care Quality*³ explored how the federal government can leverage its unique position as regulator, purchaser, provider, and research sponsor to improve care. In the report, the IOM proposed a national quality enhancement strategy focused on performance measurement of clinical quality and patient perceptions of care, as well as a proposed research agenda to support quality enhancement.

Federal and state governments, employers, and the medical community realize that action is needed to improve health care quality and patient safety. Pressure is coming from many directions to make information available to the public about the quality of hospital care. As a result, different types of quality information have been offered to the public from several sources, including insurers, the business community, consumer organizations, and commercial enterprises that compile and sell “report cards.” The potential thus exists for confusing the public with conflicting and possibly misleading information.

In an effort to alleviate some of this confusion, Connecticut is aligning itself with national efforts to begin collecting standardized data from hospitals to provide comparable information across hospitals based on valid and reliable data.

Connecticut’s Quality-in-Health-Care Initiative

During the spring of 2002, the Connecticut General Assembly passed a law creating a quality of care program within the Department of Public Health (DPH) (Sections 19a-127 l-n of the Connecticut General Statutes). The purpose of the program is to measure the quality of care provided by health care facilities in Connecticut. The intent of the legislation is twofold – to increase public accountability for the health care delivery systems of the State’s hospitals and to foster improvement in the care provided by the hospitals. Hospitals are required to collect and report quality of care information to the DPH so that it can produce a public report that compares all licensed hospitals in the state on selected quality performance measures.

National Hospital Quality Alliance

A parallel quality initiative is occurring at the national level, and Connecticut has aligned its efforts to be consistent with the national initiative. The Hospital Quality Alliance (HQA) is a national public-private collaboration to encourage hospitals to collect and report hospital quality performance information. This effort is intended to make important information about hospital performance accessible to the public and

to inform and invigorate efforts to improve quality. All of Connecticut's adult acute care hospitals are participating in this national effort.

The HQA was initiated in December 2002 by the American Hospital Association, the Federation of American Hospitals and the Association of American Medical Colleges. The effort is also supported by the Centers for Medicare and Medicaid, the Joint Commission on Accreditation of Healthcare Organizations, the American Medical Association, the American Nurses Association, the National Association of Children's Hospitals and Related Organizations, the Agency for Healthcare Research and Quality, the National Quality Forum, AARP, and the AFL-CIO.

HOSPITAL QUALITY OF CARE

What is meant by "quality" of hospital care?

Quality of hospital care can take on many meanings. It may mean that there was a successful outcome (e.g., a patient survived a heart attack or was cured of pneumonia) or it may mean that a patient was satisfied with their stay in the hospital and that they thought they were treated well. Quality care can also mean that a patient was given a needed medicine, treatment, or diagnostic test at the right time. The last definition is the one that is used in this report.

Hospitals vary in terms of their quality of care. Gaps exist between the care that could be delivered and the care that is delivered. One way to measure quality hospital care is to determine whether or not a patient got the medicine, test, or treatment that is known to be effective for his or her condition. Through extensive research, national guidelines have been established for the recommended care of patients with various medical conditions. Three common medical conditions that have been broadly studied are heart attacks, heart failure, and pneumonia. For each condition, there are a number of recommended actions, which a hospital ought to be providing to a patient.

Examples of quality care include:

- Prescribing a medication, such as aspirin, to a patient who should get it and who does not have an allergy or other medical condition making it dangerous for them to receive the medication.
- Providing an important medication or diagnostic test within the recommended time frame, for example within 24 hours of a patient having a heart attack.

What are hospital quality measures?

A hospital quality measure is an indicator that represents one aspect of the care that scientific evidence has shown to provide the best results to most people with an illness or condition. A hospital's measure of performance, also referred to as a performance rate, shows the percentage of patients who are given the right care at the right time for a specific medical condition. For example, if a hospital gives an aspirin to 80 out of 100 patients upon admission to a hospital after a heart attack, then the hospital performance rate for that particular measure is 80%.

However, standard treatment may not be the best treatment for everyone. There may be specific reasons why a patient should not get a certain treatment. For instance, a patient who is allergic to aspirin should not be given aspirin. This patient would not be counted in the measure.

This report focuses on ten hospital performance measures as follows:

Medical Condition	Performance Measure
Heart Attack	Giving an aspirin within 24 hours of arrival at a hospital if it is appropriate for the patient.
	Giving a drug called a beta-blocker within 24 hours of arrival at the hospital if it is appropriate for the patient.
	Giving a medication called an ACE inhibitor or an ARB to reduce the workload of the heart, if the function of the heart has been impaired.
	Giving a prescription for aspirin when the patient leaves the hospital, if it is appropriate for the patient.
	Giving a prescription for a drug called a beta-blocker when the patient leaves the hospital if it is appropriate for the patient.
Heart Failure	Performing a diagnostic test to determine if the heart's function has been impaired, if the test has not been done previously.
	Giving a medication called an ACE inhibitor or an ARB to reduce the workload of the heart if the function of the heart has been impaired.
Pneumonia	Giving the patient an antibiotic within 4 hours of arrival at the hospital.
	Performing a diagnostic test to determine if the patient is receiving enough oxygen.
	Screening a patient to determine if they had previously received a pneumonia vaccine, and providing the vaccine if it is appropriate for the patient.

How were the 10 hospital quality measures selected?

The ten measures included in this report focus on treatments that are considered basic recommended care for heart attack, heart failure, and pneumonia. These conditions were chosen because they represent serious medical conditions that are common reasons why patients go to hospitals. The measures for each of these conditions are considered to be a starter set for public reporting that have been extensively tested for validity and reliability and are considered best practices of care. They have been endorsed by the National Quality Forum, a national standards setting body, and have been adopted by the Centers for Medicare and Medicaid Services (CMS) as part of the Hospital Quality Alliance. In addition to these reasons, the Connecticut Department of Public Health decided to align their state reporting efforts with that of CMS in an effort to standardize the data collection process and to reduce hospitals' reporting burden.

How were the data collected and is the information accurate?

Data used to measure hospitals' performance are gathered from medical records at each hospital for patients who have been diagnosed with heart attack, heart failure, or pneumonia. Such data collection involves a combination of data obtained from existing hospital information systems and abstraction of medical records performed by trained individuals. They are the same data used by the Centers for Medicare and Medicaid and the Joint Commission in their review of hospital quality of care. Processes are in place to standardize the collection and reporting of hospital data to ensure that hospitals collect the data consistently. In addition, audits are performed to validate the accuracy of the data.

How can you use hospital quality information?

Looking at hospital quality information can be used to see how quality of care differs among hospitals. It can also be used to see how often hospitals provide the type of care considered to be recommend for several common medical conditions. It shows what treatments are usually given and how well hospitals give these treatments. This information can be used when talking to your doctor or other health care professional about the care you might need or are getting in a hospital. It can also be used when thinking about what hospital you or a family member would go to if you needed to be hospitalized.

Although this report provides information about the quality of care provided for heart attack, heart failure, and pneumonia patients, it does not include information about care provided by hospitals for other medical conditions. The care provided for the three specified conditions may or may not be reflective of the care provided for other medical conditions.

What can you do to help with your medical care?

It is important that consumers get more involved in their health care. You should contact your personal physician, if you have questions about recommended care or any exceptions that may apply to you. In addition to learning about the type of care and treatment that you might expect to receive if you need to go to a hospital, you might also consider other factors when choosing a hospital such as:

- Travel time to a hospital for you and your family
- Insurance coverage
- Cost
- Whether your family doctor is associated with a particular hospital
- Satisfaction with hospital stays experienced by others

Using this report together with other information from other sources can help you make an informed decision about your medical care.

HOSPITAL PERFORMANCE COMPARISONS

What performance rates are presented?

For each of the 10 measures, hospital performance rates are displayed for all thirty of the non-federal adult acute-care hospitals in Connecticut.

In order to provide valid comparisons, only those patients who were eligible for the recommended treatments are counted. Patients who do not meet the criteria for inclusion as described in the appendix are excluded from the analysis. As long as a hospital provides, and documents that it provided, the recommended care to the identified eligible patients, then its performance rate should approach 100%.

Data for this report were collected on patients who had been in the hospital during the calendar year from January 1, 2006 through December 31, 2006. During this period of time, some hospitals treated only a small number of patients for some of the measures. When a hospital treats a very small number of patients, its performance rate is considered to be too unreliable for public reporting. Therefore, rates are shown only for those hospitals that treated a minimum of 20 eligible patients for each measure. No inferences about hospital performance should be made when results are not presented. The actual number of cases eligible for inclusion for each hospital can be found in the appendix.

Although hospitals should strive to achieve performance rates approaching 100%, the graphs for each of the measures include an additional reference score, the statewide average rate, to be used when looking at a hospital's performance. The average performance rate for Connecticut indicates the number of times Connecticut's hospitals, as a group, provided the recommended treatment to eligible patients in the state.


Although not presented in the graphs, a second reference score is presented in Tables B4 – B6 in the appendix. It is the national average performance rate. It is based upon data reported to CMS by hospitals that are participating in the Hospital Quality Alliance. The national scores are based upon patients hospitalized from January 1, 2006 through December 31, 2006.

The performance rates displayed are estimates of a hospital's true performance. Uncertainty exists in any estimate and this should be taken into consideration when looking at the results. For each measure, small differences in the rates may not be a sign of significant differences in care. Hospitals whose performance rates differ significantly from the statewide average are designated by black circles in the graphs. Higher values are better and lower values are worse. Hospitals whose performance does not differ significantly from the statewide average are designated by grey circles.

The following figures display the hospital performance comparison results of ten performance measures for the three medical conditions of heart attack, heart failure, and pneumonia.

Figure - 1

Performance Rates* for Connecticut Hospitals
Heart Attack -- Giving an Aspirin Within 24 Hours of Hospital Arrival
January 1 - December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 2

Performance Rates* for Connecticut Hospitals
Heart Attack -- Prescribing Aspirin Upon Patient's Discharge
January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 3

Performance Rates* for Connecticut Hospitals
Heart Attack -- Giving an ACE Inhibitor or ARB if Heart is Impaired
January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 4



Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.


Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 5

Performance Rates* for Connecticut Hospitals
Heart Attack -- Giving a Beta-Blocker Within 24 Hours of Hospital Arrival
January 1- December 31, 2006

----- Hospitals -----
(sorted by Hospital name)




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 6

Performance Rates* for Connecticut Hospitals
Heart Failure -- Testing the Function of the Heart
January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 7

Performance Rates* for Connecticut Hospitals
 Heart Failure -- Giving an ACE Inhibitor or ARB if Heart is Impaired
 January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences (p<0.05). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 8

Performance Rates* for Connecticut Hospitals
 Pneumonia -- Measuring the Oxygen Levels in the Blood
 January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 9

Performance Rates* for Connecticut Hospitals
Pneumonia -- Screening and/or Providing Pneumonia Vaccine
January 1- December 31, 2006




Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences (p<0.05). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

Figure - 10

Performance Rates* for Connecticut Hospitals
 Pneumonia -- Giving Antibiotics Within 4 Hours of Hospital Arrival
 January 1- December 31, 2006



Key: The black-shaded circles identify those hospitals whose rates differ from the statewide score, based on a statistical test for significant differences ($p < 0.05$). The grey-shaded circles identify values that are not significantly different from the statewide values.

Source: Connecticut Department of Public Health, Planning Branch, Healthcare Quality, Statistics, Analysis & Reporting. April 2008.

* Performance rates are not displayed if the number of eligible patients was less than 20.

QUALITY OF CARE MEASURES FOR HEART ATTACK PATIENTS

Why is this information important?

Heart disease is the leading cause of death in the United States and Connecticut. Heart attacks, also called acute myocardial infarctions (AMI), kill more than 1,600 Connecticut residents each year. Appropriate medical care following a heart attack can greatly increase a patient's chances for recovery. Appropriate medications in the weeks following a heart attack, together with rehabilitation and changes in lifestyle, can help to prevent another heart attack from occurring.

How is quality of care determined for heart attack patients?

Research studies show that there are several steps in treating a heart attack that can make a significant difference in a patient's recovery. This report identifies five types of recommended care following a heart attack and how often Connecticut hospitals implement these recommended treatments. The recommended types of care include:

- Giving aspirin within 24 hours of the patient's arrival at the hospital, if appropriate for the patient
- Giving a prescription for aspirin when the patient leaves the hospital, if appropriate for the patient
- Giving a medication, such as an ACE inhibitor or an ARB, to reduce the pressure in the heart, if heart function has been impaired
- Giving a prescription for a beta-blocker when the patient leaves the hospital, if appropriate for the patient
- Giving a drug called a beta-blocker within 24 hours of the patient's arrival at the hospital, if appropriate for the patient

Connecticut hospital medical records for heart attack patients (January 1, 2006 through December 31, 2006) were examined to find out how often patients were given each of these recommended treatments (see Figures 1-5). Higher percentages are better.

Measure 1. Percentage of heart attack patients who are given aspirin within 24 hours of arrival at the hospital (Figure 1)

Why is this information important?

Chewing or swallowing an aspirin as soon as symptoms of a heart attack begin may help reduce the severity of the attack. Aspirin can help prevent blood clots from forming or help dissolve blood clots that have formed. Following a heart attack, continued use of aspirin may help reduce the risk of another heart

attack. Aspirin can have side effects like stomach inflammation, bleeding, or allergic reactions. Talk to your doctor before using aspirin on a regular basis.

What can you do if your hospital does not do this?

If your hospital tells you that they believe you have had a heart attack (AMI) but you have not taken an aspirin at home or in the ambulance and have not been given an aspirin on arrival to the hospital, ask your doctor or nurse if this treatment would be appropriate for you.

Measure 2. Percentage of heart attack patients who are given an aspirin at discharge (Figure 2)

Why is this information important?

Aspirin can help prevent blood clots from forming or help dissolve blood clots that have formed. Following a heart attack, continued use of aspirin may help reduce the risk of another heart attack. Aspirin can have side effects like stomach inflammation, bleeding, or allergic reactions. Talk to your doctor before using aspirin on a regular basis.

What can you do if your hospital does not do this?

If you do not already take a daily dose of aspirin and your doctor does not prescribe one at the time of discharge, ask your doctor or nurse about taking a daily aspirin.

Measure 3. Percentage of heart attack patients who are given an ACE inhibitor or an ARB at discharge (Figure 3)

Why is this information important?

Angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs) are types of medicines used to treat heart attacks, heart failure, or a decreased function of the left heart chamber (left ventricular systolic dysfunction). ACE inhibitors or ARBs can help reduce the risk of death from a heart attack if taken within 24 hours of the first symptoms of a heart attack. Continued use may help prevent heart failure. ACE inhibitors or ARBs work by limiting the effects of a hormone (angiotensin II) that narrows blood vessels and increases blood pressure. They are intended to lower blood pressure and lessen the workload of the heart.

Not all patients can take ACE inhibitors due to allergies or other side effects, in which case physicians may prescribe ARBs. ARBs act on a more specific site to block the angiotensin II hormone. This decreases potential side effects for some patients thus making the ARB more tolerable.

What can you do if your hospital does not do this?

If you have not been given a prescription for an ACE inhibitor or an ARB upon discharge, you should ask your doctor or nurse if you should be prescribed one of the medications.

Measure 4. Percentage of heart attack patients who are given a beta blocker at discharge (Figure 4)

Why is this information important?

Beta blockers are a type of medicine that is used to lower blood pressure, treat chest pain (angina) and heart failure, and to help prevent a heart attack. Beta blockers relieve the stress on the heart by slowing the heart rate and reducing the force with which the heart muscles contract to pump blood. They also help keep blood vessels from constricting in the heart, brain, and body.

What can you do if your hospital does not do this?

Not everyone can take a beta blocker. If you are unsure if you can take a beta blocker and your doctor does not give you one at the time of discharge, ask your doctor whether or not it is appropriate for you.

Measure 5. Percentage of heart attack patients who are given a beta-blocker within 24 hours of arrival at the hospital (Figure 5)

Why is this information important?

Beta blockers are a type of medicine that is used to lower blood pressure, treat chest pain (angina) and heart failure, and to help prevent a heart attack. Beta blockers relieve the stress on the heart by slowing the heart rate and reducing the force with which the heart muscles contract to pump blood. They also help keep blood vessels from constricting in the heart, brain, and body.

What can you do if your hospital does not do this?

Not everyone can take a beta blocker. However, if you have not received a beta blocker on arrival to the hospital, ask your doctor or nurse if you should receive a beta blocker.

QUALITY OF CARE MEASURES FOR HEART FAILURE PATIENTS

Why is this information important?

Heart failure, also called “congestive heart failure,” kills more than 500 Connecticut residents each year. Congestive heart failure patients are frequently hospitalized and proper hospital care is important to improve their quality of life and to prevent additional hospitalizations. Heart failure can result from a heart attack, coronary artery disease, cardiomyopathy (heart muscle damage), or an overworked heart due to long-term conditions such as high blood pressure, diabetes, or a defect from birth. The recommended treatments for someone who is getting hospital care for heart failure include:

- Giving a diagnostic test, called a left ventricular function (LVF) assessment, to determine if heart function is impaired
- Giving a medication that reduces the workload of the heart such as an ACE inhibitor or an ARB

Connecticut hospital medical records were reviewed for heart failure patients (January 1, 2006 through December 31, 2006) to find out how often patients were given each of these recommended treatments (see Figures 6 and 7). Higher percentages are better.

Measure 1. Percentage of heart failure patients given a left ventricular function (LVF) assessment before, during, or after their hospitalization (Figure 6)

Why is this information important?

The proper treatment for heart failure depends on what area of the heart is affected. An important test to check how the left chamber of the heart is pumping is the left ventricular function (LVF) assessment. It can tell the doctor whether the left side of the patient’s heart is pumping properly or not. Other evaluations include getting the patient’s medical history, examining the patient, listening to the heart sounds, and other tests as ordered by a physician. These tests may include ECG (electrocardiogram), chest x-ray, blood work, and an echocardiogram.

What should you do if you don’t receive a left ventricular function assessment?

Anyone admitted to the hospital for heart failure should be assessed for left ventricular function before or during admission, or scheduled for this assessment after discharge. If you have not received an LVF assessment, ask your doctor to schedule one.

Measure 2. Percentage of heart failure patients who are given an ACE inhibitor or an ARB at discharge (Figure 7)

Why is this information important?

Angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARBs) are types of medicines used to treat heart attacks, heart failure, or a decreased function of the left heart chamber (left ventricular systolic dysfunction). ACE inhibitors or ARBs can help reduce the risk of death from a heart attack if taken within 24 hours of the first symptoms of a heart attack. Continued use may help prevent heart failure. ACE inhibitors or ARBs work by limiting the effects of a hormone (angiotensin II) that narrows blood vessels and increases blood pressure. They are intended to lower blood pressure and lessen the workload of the heart.

Not all patients can take ACE inhibitors due to allergies or other side effects, in which case physicians may prescribe ARBs. ARBs act on a more specific site to block the angiotensin II hormone. This decreases potential side effects for some patients thus making the ARB more tolerable.

What can you do if your hospital does not do this?

If you have not been given a prescription for an ACE inhibitor or an ARB upon discharge, you should ask your doctor or nurse if you should be prescribed one of the medications.

QUALITY OF CARE MEASURES FOR PNEUMONIA PATIENTS

Why is this information important?

Pneumonia kills more than 800 Connecticut residents each year. Patients who receive the appropriate care for pneumonia are less likely to be hospitalized again for the illness. The following quality information shows the care that is the recommended treatment for persons getting hospital care for pneumonia:

- A diagnostic test to determine whether the patient is receiving enough oxygen
- A screening test to determine whether the patient has received a pneumonia vaccine and, if not, providing the vaccine if appropriate
- Giving an antibiotic to the patient within four hours of arrival at the hospital

Connecticut hospital medical records for pneumonia patients (January 1, 2006 through December 31, 2006) were examined to find out how often patients were given each of these recommended treatments (see Figures 8-10). Higher percentages are better.

Measure 1. Percentage of patients with pneumonia who are given an oxygenation assessment within 24 hours of arrival at the hospital (Figure 8)

Why is this information important?

It is important to measure the amount of oxygen in your blood to see if you need oxygen therapy. Pneumonia can lower the oxygen in your blood because the air spaces in your lungs fill with fluid. The oxygen you breathe does not get into your bloodstream. The assessment may include an arterial blood gas (ABG) or pulse oximetry (electrodes attached to a part of your body like a finger, earlobe, or skin fold).

What can you do if your hospital does not do this?

If you do not have an assessment of your oxygen level through pulse oximetry or an ABG on arrival to the hospital, ask your doctor or nurse if you should have the test.

Measure 2. Percentage of patients with pneumonia who are screened for and/or given a pneumonia vaccination before discharge from the hospital (Figure 9)

Why is this information important?

The pneumococcal vaccine may help prevent, or lower the risk of complications of pneumonia caused by bacteria. It may also help prevent future infections.

What can you do if your hospital does not do this?

Patients who have previously received a pneumonia vaccination may not need to be vaccinated again. You should keep a record of your vaccinations that can be shown to hospital staff at the time of admission. If, during your hospital stay, you do not have a fever and have not received a pneumonia vaccination, ask your doctor or nurse about vaccination.

Measure 3. Percentage of patients with pneumonia who got antibiotics within 4 hours of arrival to the hospital (Figure 10)

Why is this information important?

Antibiotics are used to treat pneumonia caused by bacteria. Early treatment with antibiotics can cure bacterial pneumonia and reduce the possibility of complications.

What can you do if your hospital does not do this?

You may have received antibiotics from your physician before admission to the hospital; therefore you may not receive antibiotics within 4 hours of arrival at the hospital. If you have not received antibiotics before your admission to the hospital, ask your doctor or nurse if you will be receiving an antibiotic.

DISCUSSION

Hospital Mergers

On October 1, 2006, New Britain General Hospital and Bradley Memorial Hospital merged to become The Hospital of Central Connecticut with campuses in New Britain and Southington. Performance rates for The Hospital of Central Connecticut at Bradley Memorial include data for only the first three quarters of 2006. Fourth quarter data for Bradley Memorial are added to that for The Hospital of Central Connecticut at New Britain General.

Small Numbers of Cases on Individual Performance Measures

During 2006, some hospitals reported a small number of cases for one or more of the performance measures. In such cases, rates are considered to be too unreliable for public reporting. Therefore, rates are shown only for those measures for which hospitals treated a minimum of 20 eligible patients. No inferences can be made for those hospitals whose results are not presented. There were 33 out of a possible 300 hospital performance rates (11%) that could not be displayed due to the small number of eligible cases being reported.

Performance Rates for 2006

Table 1 shows the range of performance rates in Connecticut as well as a comparison of Connecticut hospitals' average performance rates to the average performance rates of hospitals in the United States. Connecticut hospitals performance rates average exceed national averages on all ten of the clinical measures, and are significantly better on nine of the ten measures. Hospitals can achieve high levels of performance as seen by the high end of the ranges, but there is still considerable variability in the performance of hospitals on most measures, suggesting inconsistencies in the care provided.

Using a performance benchmark of 90%¹, variability in attainment can be seen in Figure 11. For the pneumonia measure “Antibiotic within 4 hours of arrival at the hospital,” only 13% of the hospitals achieved a performance rate greater than 90%. At the other extreme, “Oxygenation assessment” for Pneumonia and “Beta-blocker at discharge” for Heart Attack, all of the hospitals (i.e., 100%) have rates in excess of 90%. However, most hospitals still fall short of the goal of 100% on most of the measures. That is, performance gaps still exist between the care that could be given and the care that is being delivered.

¹ The 90% benchmark was chosen to reflect the benchmark being considered by CMS for Medicare Hospital Value-Based Purchasing, particularly for measures with hospital performance concentrated around high values.

Table 1


Connecticut's Performance Compared to the U.S. Performance, 2006				
Condition	Measure	Range in Connecticut 2006	Average	Average
			Connecticut Rate 2006	National Rate** 2006
Heart Attack	Aspirin at Arrival	89% - 100%	98%*	93%
	Aspirin at Discharge	88% - 100%	98%*	90%
	ACEI or ARB for LVSD at Discharge	68% - 100%	85%	83%
	Beta-Blocker at Discharge	91% - 100%	97%*	90%
	Beta-Blocker at Arrival	85% - 100%	95%*	87%
Heart Failure	LVF Assessment	84% - 100%	96%*	84%
	ACEI or ARB for LVSD at Discharge	74% - 100%	86%*	82%
Pneumonia	Oxygenation Assessment	99% - 100%	100%*	99%
	Pneumococcal Vaccination	10% - 96%	79%*	71%
	Timely Antibiotic	68% - 94%	81%*	80%

** Source: www.hospitalcompare.hhs.gov for hospitals participating in the Hospital Quality Alliance initiative.

Data are based upon patients hospitalized from 1/1/06 – 12/31/06.

* Difference is statistically significant (p < 0.05).

Figure 11



Hospitals were excluded from a measure if they had fewer than 20 cases.

At both the state and national levels, average performance rates are low for the two measures related to the administration of an angiotensin converting enzyme inhibitor (ACEI) or an angiotensin receptor blocker (ARB) for either heart attack or heart failure patients. Rates are also low for the pneumonia measures related to the administration of timely antibiotics and pneumococcal vaccinations. These four rates, which are discussed further below, also have the widest inter-hospital range, suggesting that significant differences in practice patterns exist and that better performing hospitals may have developed practices that might ultimately benefit other hospitals that choose to adopt similar methods.

As previously noted, average Connecticut performance rates are low for the measures related to the administration of an ACEI or ARB for either heart attack (85%) or heart failure (86%) patients (Table 1). Only a third of the hospitals achieve a rate of 90% or better for the heart failure measure and half of the hospitals achieve a high rate for the heart attack measure (Figure 11). It had been anticipated that these rates would improve with the change in the definition of the performance measures to reflect the acceptable use of either ACEI *or* ARB for treatment of patients with LVSD at discharge, effective with January 1, 2005 discharges. Progress has been made. The statewide average rate related to the administration of an ACEI or ARB for heart attack increased from 82% in 2005 to 85% in 2006. The rate for heart failure patients also increased from 83% in 2005 to 86% in 2006 (Table 2). Both increases were statistically significant ($p < 0.05$).

Connecticut hospitals have had varying success in vaccinating all eligible patients aged 65 and over for pneumonia. In 2006, pneumococcal vaccination rates in Connecticut ranged from 10% to 96% with an average rate of 79%, which was a significant increase over the 67% average rate in 2005. Nearly two thirds of the hospitals (19 out of 30; Table 3) have seen a significant increase in their performance rates from 2005 to 2006 for this measure. This improvement is most likely due to a legislative change in Connecticut that allows hospitals to administer influenza and pneumococcal polysaccharide vaccines to patients without requiring an individual physician's order. Although the law was passed in 2004, implementation regulations were not approved until the fall of 2005.

Seven out of 30 hospitals showed significant improvement between 2005 and 2006 in their performance rates for the pneumonia measure of giving antibiotics within four hours of arrival at the hospital (Table 3). Despite this improvement, less than 14% of the hospitals achieved a 90% or greater performance rate in 2006 (Figure 11). In 2006, inter-hospital performance rates vary widely from 68% to 94% with a statewide average rate of 81% (Table 1). One reason for delayed administration of antibiotics may be due

to delays arising in diagnosing and treating patients arriving at crowded emergency departments. Those patients who are less sick or who have a nonclassic presentation may not receive the same priority as patients with suspected pneumonia.⁴

Another possible reason for the delayed administration of antibiotics is that many patients are treated by their primary care physician for pneumonia before they need hospitalization, in which case patients may be placed on antibiotics while still at home. Many of the new antibiotics are longer acting and may be given only once or twice a day. Therefore, a patient already taking an antibiotic who is then admitted to a hospital may not receive the next dose until the next time that it is due to be given, which may exceed the four-hour target being measured. If this information is not documented in the patient's medical record, the case will be incorrectly counted as a failure to give timely antibiotics.

The time to delivery of the first antibiotic dose for pneumonia patients has received much attention. On April 20, 2007, the National Quality Forum Pulmonary Consensus Standards Maintenance Committee endorsed the new pneumonia measure “Initial Antibiotic Received within Six Hours of Hospital Arrival” (PN-5c). The Centers for Medicare and Medicaid Services (CMS) will begin public reporting of PN-5c on the *Hospital Compare* website beginning in March 2008. CMS will suppress reporting of the measure “Initial Antibiotic Received within Four Hours of Hospital Arrival” (PN-5b).

Changes in the Quality of Hospital Care

Although some performance rates are still relatively low, most are high, and all are improving. Between 2005 and 2006, Connecticut hospitals' average performances rates increased significantly for all ten measures (Table 2). Those measures with the lowest rates in 2005 tended to increase the most. On a hospital-measure-specific level, 51 out of 262 hospital performance measures showed statistically significant rate increases from 2005 to 2006 (Table 3). Thirty-eight hospital measures were excluded due to small sample sizes in either year 2005 or 2006. Nearly two-thirds of the hospitals showed significant improvement on one of the three pneumonia measures, but there is still much room for improvement here. Details by hospital can be found in the appendices.

Table 2

Connecticut's Performance between 2005 and 2006			
Condition	Measure	2005	2006
Heart Attack	Aspirin at Arrival	96%	98%*
	Aspirin at Discharge	97%	98%*
	ACEI or ARB for LVSD at Discharge	82%	85%*
	Beta-Blocker at Discharge	96%	97%*
	Beta-Blocker at Arrival	94%	95%*
Heart Failure	LVF Assessment	95%	96%*
	ACEI or ARB for LVSD at Discharge	83%	86%*
Pneumonia	Oxygenation Assessment	100%	100%*#
	Pneumococcal Vaccination	67%	79%*
	Timely Antibiotic	79%	81%*

* Difference is statistically significant ($p < 0.05$).

Oxygenation Assessment for Pneumonia increased from 99.8% in 2005 to 99.9% in 2006 ($p = 0.02$).

Table 3

Change in Hospital Performance from 2005 to 2006*				
Measure	No. of Hospitals (out of 30) with 20+ Cases	No. of Hospitals with Significant Rate Increases	No. of Hospitals with Significant Rate Decreases	No. of Hospitals with no Significant Rate Changes
Heart Attack				
Aspirin at Arrival	27	4	1	22
Aspirin at Discharge	24	4	0	20
ACEI for LVSD at Discharge	13	2	0	11
Beta-Blocker at Discharge	24	3	0	21
Beta-Blocker at Arrival	26	2	1	23
Heart Failure				
LVF Assessment	30	8	1	21
ACEI for LVSD at Discharge	28	2	2	24
Pneumonia				
Oxygenation Assessment	30	0	0	30
Pneumococcal Vaccination	30	19	2	9
Timely Antibiotic	30	7	2	21
	262	51	9	202

* Hospitals were excluded from a measure if they had fewer than 20 cases for either 2005 or 2006. Differences are significant if $p < 0.05$.

REFERENCES

1. Institute of Medicine. *To err is human: building a safer health system*. Washington DC: *National Academy Press*, 2000.
2. Institute of Medicine. *Crossing the quality chasm: a new health system for the 21st century*. *National Academy Press*, 2001.
3. Institute of Medicine. *Leadership by example: coordinating government roles in improving health care quality*. *National Academy Press*, 2003.
4. Pines JM, Localio AR, Hollander JE, et al. The impact of emergency department crowding measures on time to antibiotics for patients with community-acquired pneumonia. *Ann Emerg Med* 2007;50:510-516.

RESOURCES

Below are some useful resources if you would like more information about hospital quality of care.

The **Connecticut Department of Public Health** is the state agency responsible for developing the Hospital Performance Comparisons Report. It is also the agency responsible for the licensing and regulatory oversight of Connecticut hospitals. For more information about the activities in the Connecticut Department of Public Health, visit their website at www.dph.state.ct.us or call them at 860-509-8000.

The **Connecticut Hospital Association** represents and serves Connecticut's hospitals. For more information about the hospitals in Connecticut, contact the Connecticut Hospital Association at www.cthosp.org or 203-294-7213.

Qualidigm® is the Quality Improvement Organization for Connecticut under the direction of the Centers for Medicare and Medicaid. They implement quality improvement programs with hospitals and serve as advocates for Medicare beneficiaries. Contact them at www.qualidigm.org or 860-632-2008.

For more information about the **Hospital Quality Alliance** initiative of the **Centers for Medicare and Medicaid**, visit the web site www.hospitalcompare.hhs.gov/.

The **Joint Commission** evaluates the quality and safety of care of health care organizations and accredits them. They have prepared information to help consumers select a hospital. Go to their website at www.jointcommission.org or call their Customer Service Department at 630-792-5800.

The **Agency for Health Research and Quality** (AHRQ) is the lead federal agency responsible for research on quality, cost, access, utilization, and health care outcomes and patient safety. AHRQ has a variety of resources for consumers including *Your Guide to Choosing Quality Health Care*. Visit their website at www.ahrq.gov.

ACKNOWLEDGEMENTS

We would like to thank Qualidigm® for their assistance in the data collection process and also the Connecticut Hospital Association for their valuable insight concerning the ongoing reporting process.

CONNECTICUT HOSPITALS

Only licensed hospitals that regularly care for adults with heart attacks, heart failure, and pneumonia are included in this report. This report does not contain information from pediatric, psychiatric, or rehabilitation hospitals.

For more information about the quality of care provided by hospitals in Connecticut, contact the quality improvement department of any of the hospitals listed below or visit the hospital's web site.

<p>The William W. Backus Hospital 326 Washington Street Norwich, CT 06360-2733 860-889-8331 www.backushospital.org</p>	<p>John Dempsey Hospital 263 Farmington Avenue Farmington, CT 06032-1941 860-679-2000 http://health.uhc.edu</p>
<p>Bradley Memorial Campus, The Hospital of Central Connecticut 81 Meriden Avenue Southington, CT 06489-3297 860-276-5000 www.thocc.org</p>	<p>Greenwich Hospital 5 Perryridge Road Greenwich, CT 06830-4697 203-863-3000 www.greenhosp.org</p>
<p>Bridgeport Hospital 267 Grant Street Bridgeport, CT 06610-0120 203-384-3000 www.bpthosp.org</p>	<p>Griffin Hospital 130 Division Street Derby, CT 06418-1326 203-735-7421 www.griffinhealth.org</p>
<p>Bristol Hospital Brewster Road Bristol, CT 06011-0977 860-585-3000 www.bristolhospital.org</p>	<p>Hartford Hospital 80 Seymour Street Hartford, CT 06102-5037 860-545-5000 www.harthosp.org</p>
<p>Danbury Hospital 24 Hospital Avenue Danbury, CT 06810-6099 203-739-7000 www.danhosp.org</p>	<p>The Charlotte Hungerford Hospital 540 Litchfield Street Torrington, CT 06790-0988 860-496-6666 www.charlottesweb.hungerford.org</p>
<p>Day Kimball Hospital 320 Pomfret Street Putnam, CT 06260-0901 860-928-6541 www.daykimball.org</p>	<p>Johnson Memorial Hospital 201 Chestnut Hill Road Stafford Springs, CT 06076-0860 860-684-4251 www.johnsonhealthnetwork.com/jmhinc.htm</p>

<p>Lawrence & Memorial Hospital 365 Montauk Avenue New London, CT 06320-4769 860-442-0711, ext. 2356 www.lmhosp.org</p>	<p>Saint Francis Hospital and Medical Center 114 Woodland Street Hartford, CT 06105-1200 860-714-4000 www.stfranciscare.org</p>
<p>Manchester Memorial Hospital 71 Haynes Street Manchester, CT 06040-4188 860-646-1222 www.echn.org</p>	<p>Saint Mary's Hospital 56 Franklin Street Waterbury, CT 06706-1281 203-709-6000 www.stmh.org</p>
<p>Middlesex Hospital 28 Crescent Street Middletown, CT 06457-3650 860-344-6000 www.midhosp.org</p>	<p>Hospital of Saint Raphael 1450 Chapel Street New Haven, CT 06511-1450 203-789-3000 www.srhs.org</p>
<p>MidState Medical Center 435 Lewis Avenue Meriden, CT 06451-2101 203-694-8200 www.midstatemedical.org</p>	<p>St. Vincent's Medical Center 2800 Main Street Bridgeport, CT 06606-4292 203-576-6000 www.svhs-ct.org</p>
<p>Milford Hospital 300 Seaside Avenue Milford, CT 06460-4603 203-876-4000 www.milfordhospital.org</p>	<p>Sharon Hospital 50 Hospital Hill Road Sharon, CT 06069-0789 860-364-4141 www.sharonhospital.com</p>
<p>New Britain General Campus, The Hospital of Central Connecticut 100 Grand Street New Britain, CT 06052-2017 860-224-5011 www.thocc.org</p>	<p>The Stamford Hospital 30 Shelburne Road Stamford, CT 06904-9317 203-276-1000 www.stamhealth.org</p>
<p>New Milford Hospital 21 Elm Street New Milford, CT 06776-2993 860-355-2611 www.newmilfordhospital.org</p>	<p>Waterbury Hospital 64 Robbins Street Waterbury, CT 06721 203-573-6000 www.waterburyhosp.org</p>
<p>Norwalk Hospital 34 Maple Street Norwalk, CT 06850-3894 203-852-2000 www.norwalkhosp.org</p>	<p>Windham Community Memorial Hospital 112 Mansfield Avenue Willimantic, CT 06226-2040 860-456-9116 www.wcmh.org</p>
<p>Rockville General Hospital 31 Union Street Vernon, CT 06066-3160 860-872-0501 www.echn.org</p>	<p>Yale-New Haven Hospital 20 York Street New Haven, CT 06510-3202 203-688-4242 www.ynhh.org</p>

Appendices A - B

Appendix A

DEFINITIONS OF MEASURES

Acute Myocardial Infarction (AMI or Heart Attack)

For the purposes of this report the AMI population consists of those patients over the age of 18 with a discharge ICD-9-CM code indicating an initial AMI episode (410.x1).

Each measure within the AMI measure set is calculated individually based on the inclusion/exclusion criteria for that particular measure; therefore, the denominators for each measure may be different.

Aspirin at arrival

Exclusion criteria:

- Less than 18 years of age
- Patients transferred from another acute care hospital on the day of arrival
- Patients received in transfer from another hospital, including another emergency department
- Patients discharged on day of arrival
- Patients expired on day of arrival
- Patients who left against medical advice on day of arrival
- Patients with contraindication to aspirin including:
 - active bleeding on arrival or within 24 hours of arrival,
 - aspirin allergy,
 - on warfarin/Coumadin prior to arrival
- Other explicitly linked reason documented by a physician, nurse practitioner or physician assistant for not giving aspirin on arrival

Aspirin at discharge

Exclusion criteria:

- Less than 18 years of age
- Patients transferred to another acute care hospital
- Patients who expired
- Patients who left against medical advice
- Patients discharged to hospice

- Patients with contraindication to aspirin including:
 - active bleeding on arrival or within 24 hours of arrival,
 - aspirin allergy,
 - on warfarin/Coumadin prior to arrival
- Other explicitly linked reason documented by a physician, nurse practitioner or physician assistant for not prescribing aspirin at discharge

ACEI or ARB for LVSD

Inclusion criteria:

- Chart documentation of Left Ventricular Ejection Fraction (LVEF) less than 40% or a narrative description of LVS consistent with moderate or severe systolic dysfunction

Exclusion criteria:

- Less than 18 years of age
- Patients transferred to another acute care hospital or federal hospital
- Patients who expired
- Patients who left against medical advice
- Patients discharged to hospice
- Patients with both a potential contraindication/reason for not prescribing an ACEI at discharge and a potential contraindication/reason for not prescribing an ARB at discharge, as evidenced by one or more of the following:
 - ACEI allergy and ARB allergy,
 - Moderate or severe aortic stenosis,
 - Physician, nurse practitioner, or physician assistant documentation of both a reason for not prescribing an ACEI at discharge and a reason for not prescribing an ARB at discharge,
 - Reason documented by physician, nurse practitioner, or physician assistant for not prescribing an ARB at discharge and an ACEI allergy,
 - Reason documented by physician, nurse practitioner, or physician assistant for not prescribing an ACEI at discharge and an ARB allergy.

Beta blocker prescribed at discharge

Exclusion criteria:

- Less than 18 years of age

- Patients transferred to another acute care hospital
- Patients who expired
- Patients who left against medical advice
- Patients discharged to hospice
- Patients with one or more of the following beta blocker contraindications/reasons for not prescribing:
 - Beta blocker allergy,
 - Bradycardia (heart rate less than 60 bpm) on day of discharge or day prior to discharge while not on beta blocker
 - Second or third degree heart block on ECG on arrival or during hospital stay and does not have a pacemaker
 - Systolic BP less than 90 mmHg on day of discharge or day prior to discharge while not on beta blocker
- Other explicitly linked reason documented by a physician, nurse practitioner or physician assistance

Beta blocker at arrival

Exclusion criteria:

- Less than 18 years of age
- Patients transferred from another acute care hospital on the day of arrival
- Patients received in transfer from another hospital, including another emergency department
- Patients discharged on day of arrival
- Patients expired on day of arrival
- Patients who left against medical advice on day of arrival
- Patients with one or more of the following beta blocker contraindications/reasons for not prescribing:
 - Beta blocker allergy,
 - Bradycardia (heart rate less than 60 bpm) on arrival or within 24 hours of arrival while not on beta blocker
 - Heart failure on arrival or within 24 hours after arrival
 - Second or third degree heart block on ECG on arrival or within 24 hours after arrival and does not have a pacemaker
 - Shock on arrival or within 24 hours after arrival

- Systolic BP less than 90 mmHg on arrival or within 24 hours after arrival
- Other explicitly linked reason documented by a physician, nurse practitioner or physician assistant for not giving a beta blocker within 24 hours after hospital arrival

Heart Failure

For the purposes of this report the Heart Failure population consists of those patients over the age of 18 with a discharge ICD-9-CM code indicating a Heart Failure episode (402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 404.93, 428.0, 428.1, 428.20, 428.21, 428.22, 428.23, 428.30, 428.31, 428.32, 428.33, 428.40, 428.41, 429.42, 428.43, 428.9).

Each measure within the Heart Failure measure set is calculated individually based on the inclusion/exclusion criteria for that particular measure; therefore, the denominators for each measure may be different.

Left Ventricular Function (LVF) Assessment

Exclusion criteria:

- Patients less than 18 years of age
- Patients transferred to another acute care hospital
- Patients who expired
- Patients who left against medical advice
- Patients discharged to hospice
- Other explicitly linked reason documented by a physician, nurse practitioner or physician assistant for no LVF assessment

ACEI or ARB for LVSD

Inclusion criteria:

- Chart documentation of Left Ventricular Ejection Fraction (LVEF) less than 40% or a narrative description of LVS consistent with moderate or severe systolic dysfunction

Exclusion criteria:

- Less than 18 years of age
- Patients transferred to another acute care hospital or federal hospital
- Patients who expired

- Patients who left against medical advice
- Patients discharged to hospice
- Patients with both a potential contraindication/reason for not prescribing an ACEI at discharge and a potential contraindication/reason for not prescribing an ARB at discharge, as evidenced by one or more of the following:
 - ACEI allergy and ARB allergy,
 - Moderate or severe aortic stenosis,
 - Physician, nurse practitioner, or physician assistant documentation of both a reason for not prescribing an ACEI at discharge and a reason for not prescribing an ARB at discharge,
 - Reason documented by physician, nurse practitioner, or physician assistant for not prescribing an ARB at discharge and an ACEI allergy,
 - Reason documented by physician, nurse practitioner, or physician assistant for not prescribing an ACEI at discharge and an ARB allergy.

Pneumonia

For the purposes of this report the pneumonia population consists of those patients over the age of 29 days with a discharge ICD-9-CM code indicating a principal diagnosis of pneumonia or a principal diagnosis of septicemia or respiratory failure with a secondary diagnosis of pneumonia.

Patients without a working diagnosis of pneumonia on admission or those for whom “comfort measures only” are prescribed during their hospitalization are immediately excluded from the population.

Each measure within the pneumonia measure set is calculated individually based on the inclusion/exclusion criteria for that particular measure; therefore, the denominators for each measure may be different.

Oxygenation assessment

Exclusion criteria:

- Patients received in transfer from another acute care hospital
- Patients who have no working diagnosis of pneumonia at the time of admission
- Patients receiving “comfort measures only”
- Patients less than 29 days of age

Pneumococcal screening and/or vaccination

Inclusion criteria:

- Patients over 65 years of age with a principal diagnosis of pneumonia or a principal diagnosis of septicemia or respiratory failure with a secondary diagnosis of pneumonia

Exclusion criteria:

- Patients received in transfer from another acute care hospital
- Patients who left against medical advice
- Patients who have no working diagnosis of pneumonia at the time of admission
- Patients receiving “comfort measures only”
- Patients less than 65 years of age
- Patient expired

Antibiotic timing

Exclusion criteria:

- Patients received in transfer from another acute care hospital
- Patients who have no working diagnosis of pneumonia at the time of admission
- Patients who do not receive antibiotics during hospitalization
- Patients receiving “comfort measures only”
- Patients less than 29 days of age
- Patients whose initial antibiotic was administered more than 36 hours from the time of arrival
- Does not include antibiotics received prior to hospitalization but this may cause an extended time to initial antibiotic in the hospital

Appendix B

HEART ATTACK, HEART FAILURE, AND PNEUMONIA PERFORMANCE RATES

Tables B1 – B3 display 2004 performance rates and the number of eligible patients for each hospital and individual measure for each of the three medical conditions -- heart attack, heart failure, and pneumonia. Comparison scores include the statewide average rate and the national average rate.

Tables B4 – B6 display 2005 performance rates and the number of eligible patients for each hospital and individual measure for each of the three medical conditions -- heart attack, heart failure, and pneumonia. Comparison scores include the statewide average rate and the national average rate.

Table B7 shows the change in hospital performance rates from 2004 to 2005 for those hospital-specific measures with statistically significant differences. These data are sorted by hospital name.

Table B1

Heart Attack Performance Rates for Connecticut Hospitals					
<i>January 1, 2005 to December 31, 2005</i>					
	Aspirin at Arrival	Aspirin at Discharge	ACEI or ARB for LVSD	Beta Blocker at Discharge	Beta Blocker at Arrival
National Average Rate*	92%	89%	80%	88%	86%
Connecticut Average Rate	96%	97%	82%	96%	94%
Bradley Memorial Hospital & Health Center	95% of 39 patients	**	**	**	92% of 26 patients
Bridgeport Hospital	96% of 170 patients	97% of 272 patients	91% of 44 patients	98% of 297 patients	96% of 140 patients
Bristol Hospital	93% of 73 patients	89% of 35 patients	61% of 23 patients	98% of 40 patients	98% of 55 patients
Charlotte Hungerford Hospital	95% of 59 patients	90% of 40 patients	**	92% of 39 patients	96% of 52 patients
Danbury Hospital	98% of 186 patients	99% of 134 patients	85% of 41 patients	95% of 139 patients	95% of 169 patients
Day Kimball Hospital	91% of 54 patients	96% of 25 patients	**	89% of 28 patients	74% of 46 patients
Greenwich Hospital Association	98% of 84 patients	100% of 38 patients	**	98% of 45 patients	98% of 63 patients
Griffin Hospital	99% of 116 patients	100% of 49 patients	**	98% of 65 patients	99% of 106 patients
Hartford Hospital	92% of 415 patients	94% of 729 patients	81% of 225 patients	95% of 881 patients	89% of 320 patients
Hospital Of St Raphael	93% of 275 patients	97% of 385 patients	77% of 93 patients	93% of 363 patients	81% of 284 patients
John Dempsey Hospital	99% of 113 patients	100% of 240 patients	100% of 46 patients	100% of 241 patients	100% of 112 patients
Johnson Memorial Hospital	95% of 20 patients	**	**	**	**
Lawrence & Memorial Hospital	97% of 171 patients	99% of 72 patients	**	100% of 80 patients	97% of 157 patients
Manchester Memorial Hospital	91% of 88 patients	97% of 33 patients	75% of 20 patients	100% of 42 patients	93% of 73 patients
Middlesex Hospital	96% of 169 patients	95% of 86 patients	86% of 44 patients	98% of 104 patients	96% of 108 patients
MidState Medical Center	94% of 125 patients	95% of 58 patients	**	94% of 63 patients	96% of 92 patients
Milford Hospital	95% of 58 patients	96% of 26 patients	**	100% of 24 patients	94% of 49 patients
New Britain General Hospital	97% of 225 patients	95% of 117 patients	78% of 41 patients	99% of 150 patients	99% of 152 patients
New Milford Hospital	**	**	**	**	**
Norwalk Hospital	96% of 127 patients	94% of 52 patients	83% of 23 patients	99% of 77 patients	97% of 73 patients
Rockville General Hospital	98% of 46 patients	100% of 27 patients	**	100% of 28 patients	**
Sharon Hospital	100% of 22 patients	**	**	**	95% of 21 patients
St Francis Hospital & Medical Center	94% of 186 patients	97% of 364 patients	77% of 170 patients	96% of 455 patients	91% of 101 patients
St Mary's Hospital	99% of 155 patients	98% of 107 patients	87% of 31 patients	99% of 113 patients	98% of 146 patients
St Vincent's Medical Center	97% of 346 patients	97% of 433 patients	88% of 144 patients	94% of 454 patients	94% of 324 patients
Stamford Hospital	99% of 109 patients	98% of 52 patients	**	98% of 63 patients	97% of 76 patients
Waterbury Hospital	95% of 144 patients	90% of 83 patients	70% of 23 patients	84% of 95 patients	96% of 103 patients
William W Backus Hospital	98% of 119 patients	100% of 46 patients	**	100% of 66 patients	100% of 78 patients
Windham Community Memorial Hospital	93% of 43 patients	95% of 21 patients	**	95% of 22 patients	86% of 29 patients
Yale-New Haven Hospital	98% of 132 patients	98% of 197 patients	85% of 65 patients	97% of 272 patients	97% of 91 patients

* Source: CMS Hospital Compare based on data 1/1/05 – 12/31/05.

** Performance rates are not displayed if denominators were less than 20 during the reporting period.

Table B2

Heart Failure Performance Rates for Connecticut Hospitals		
<i>January 1, 2005 to December 31, 2005</i>		
	LVF Assessment	ACEI or ARB for LVSD
National Average Rate*	81%	81%
Connecticut Average Rate	95%	83%
Bradley Memorial Hospital & Health Center	97% of 132 patients	86% of 36 patients
Bridgeport Hospital	98% of 335 patients	90% of 147 patients
Bristol Hospital	90% of 218 patients	73% of 90 patients
Charlotte Hungerford Hospital	96% of 135 patients	77% of 57 patients
Danbury Hospital	96% of 475 patients	81% of 204 patients
Day Kimball Hospital	94% of 145 patients	87% of 46 patients
Greenwich Hospital Association	91% of 291 patients	84% of 94 patients
Griffin Hospital	100% of 248 patients	89% of 73 patients
Hartford Hospital	92% of 797 patients	76% of 380 patients
Hospital Of St Raphael	96% of 571 patients	68% of 223 patients
John Dempsey Hospital	91% of 183 patients	100% of 77 patients
Johnson Memorial Hospital	89% of 94 patients	92% of 24 patients
Lawrence & Memorial Hospital	98% of 360 patients	85% of 122 patients
Manchester Memorial Hospital	97% of 225 patients	79% of 78 patients
Middlesex Hospital	95% of 266 patients	89% of 102 patients
MidState Medical Center	96% of 359 patients	77% of 114 patients
Milford Hospital	94% of 217 patients	78% of 64 patients
New Britain General Hospital	90% of 584 patients	78% of 160 patients
New Milford Hospital	92% of 60 patients	93% of 27 patients
Norwalk Hospital	99% of 424 patients	92% of 149 patients
Rockville General Hospital	99% of 134 patients	79% of 42 patients
Sharon Hospital	93% of 59 patients	**
St Francis Hospital & Medical Center	95% of 571 patients	83% of 291 patients
St Mary's Hospital	90% of 344 patients	79% of 121 patients
St Vincent's Medical Center	95% of 621 patients	92% of 247 patients
Stamford Hospital	98% of 371 patients	95% of 152 patients
Waterbury Hospital	87% of 381 patients	72% of 138 patients
William W Backus Hospital	100% of 320 patients	100% of 88 patients
Windham Community Memorial Hospital	97% of 146 patients	74% of 72 patients
Yale-New Haven Hospital	99% of 341 patients	78% of 120 patients

* Source: CMS Hospital Compare based on data 1/1/05 – 12/31/05.

** Performance rates are not displayed if the number of eligible patients was less than 20 during the reporting period.

Table B3

Pneumonia Performance Rates for Connecticut Hospitals <i>January 1, 2005 to December 31, 2005</i>			
	Oxygenation Assessment	Pneumococcal Vaccination	Timely Antibiotic
National Average Rate*	99%	59%	77%
Connecticut Average Rate	100%	67%	79%
Bradley Memorial Hospital & Health Center	100% of 169 patients	56% of 122 patients	85% of 138 patients
Bridgeport Hospital	100% of 251 patients	65% of 161 patients	74% of 214 patients
Bristol Hospital	98% of 426 patients	85% of 296 patients	84% of 368 patients
Charlotte Hungerford Hospital	100% of 251 patients	94% of 166 patients	83% of 189 patients
Danbury Hospital	100% of 528 patients	79% of 383 patients	83% of 401 patients
Day Kimball Hospital	100% of 216 patients	69% of 131 patients	79% of 201 patients
Greenwich Hospital Association	100% of 242 patients	88% of 173 patients	88% of 205 patients
Griffin Hospital	100% of 215 patients	92% of 156 patients	86% of 170 patients
Hartford Hospital	100% of 824 patients	25% of 484 patients	57% of 616 patients
Hospital Of St Raphael	100% of 344 patients	58% of 259 patients	82% of 270 patients
John Dempsey Hospital	97% of 195 patients	1% of 137 patients	74% of 170 patients
Johnson Memorial Hospital	100% of 152 patients	70% of 117 patients	85% of 107 patients
Lawrence & Memorial Hospital	100% of 422 patients	94% of 294 patients	86% of 325 patients
Manchester Memorial Hospital	100% of 278 patients	48% of 193 patients	66% of 235 patients
Middlesex Hospital	100% of 459 patients	63% of 347 patients	83% of 420 patients
MidState Medical Center	100% of 269 patients	79% of 190 patients	86% of 228 patients
Milford Hospital	100% of 293 patients	79% of 201 patients	79% of 225 patients
New Britain General Hospital	100% of 440 patients	79% of 315 patients	88% of 383 patients
New Milford Hospital	100% of 159 patients	51% of 126 patients	80% of 148 patients
Norwalk Hospital	100% of 453 patients	74% of 310 patients	78% of 362 patients
Rockville General Hospital	100% of 199 patients	61% of 117 patients	67% of 169 patients
Sharon Hospital	100% of 108 patients	34% of 71 patients	98% of 91 patients
St Francis Hospital & Medical Center	100% of 280 patients	73% of 185 patients	72% of 216 patients
St Mary's Hospital	100% of 299 patients	27% of 178 patients	75% of 243 patients
St Vincent's Medical Center	100% of 515 patients	83% of 362 patients	85% of 452 patients
Stamford Hospital	100% of 380 patients	87% of 282 patients	74% of 342 patients
Waterbury Hospital	99% of 328 patients	47% of 228 patients	76% of 259 patients
William W Backus Hospital	100% of 386 patients	62% of 256 patients	79% of 368 patients
Windham Community Memorial Hospital	100% of 305 patients	51% of 191 patients	79% of 246 patients
Yale-New Haven Hospital	100% of 221 patients	88% of 113 patients	81% of 197 patients

* Source: CMS Hospital Compare based on data 1/1/05 – 12/31/05.

** Performance rates are not displayed if the number of eligible patients was less than 20 during the reporting period.

Table B4

Heart Attack Performance Rates for Connecticut Hospitals					
January 1, 2006 to December 31, 2006					
	Aspirin at Arrival	Aspirin at Discharge	ACEI or ARB for LVSD	Beta Blocker at Discharge	Beta Blocker at Arrival
National Average Rate*	93%	90%	83%	90%	87%
Connecticut Average Rate	98%	98%	85%	97%	95%
Bridgeport Hospital	99% of 225 patients	100% of 257 patients	87% of 70 patients	97% of 270 patients	97% of 174 patients
Bristol Hospital	91% of 45 patients	96% of 24 patients	**	97% of 38 patients	92% of 37 patients
Charlotte Hungerford Hospital	94% of 48 patients	92% of 38 patients	**	91% of 44 patients	85% of 55 patients
Danbury Hospital	100% of 242 patients	99% of 297 patients	82% of 60 patients	95% of 287 patients	95% of 217 patients
Day Kimball Hospital	94% of 32 patients	96% of 24 patients	**	100% of 21 patients	90% of 20 patients
Greenwich Hospital Association	100% of 76 patients	98% of 53 patients	**	100% of 45 patients	100% of 51 patients
Griffin Hospital	100% of 87 patients	98% of 50 patients	91% of 23 patients	97% of 63 patients	99% of 69 patients
Hartford Hospital	96% of 349 patients	98% of 628 patients	85% of 216 patients	96% of 746 patients	88% of 282 patients
HCC at Bradley Memorial	100% of 29 patients	**	**	95% of 20 patients	96% of 26 patients
HCC at New Britain General	98% of 182 patients	89% of 121 patients	79% of 43 patients	97% of 143 patients	96% of 153 patients
Hospital of St Raphael	99% of 248 patients	100% of 347 patients	100% of 57 patients	100% of 352 patients	99% of 234 patients
John Dempsey Hospital	100% of 116 patients	100% of 159 patients	100% of 34 patients	100% of 157 patients	100% of 113 patients
Johnson Memorial Hospital	**	**	**	**	**
Lawrence & Memorial Hospital	98% of 128 patients	99% of 85 patients	90% of 21 patients	98% of 90 patients	99% of 99 patients
Manchester Memorial Hospital	95% of 42 patients	100% of 26 patients	**	98% of 42 patients	87% of 45 patients
Middlesex Hospital	98% of 89 patients	97% of 73 patients	97% of 32 patients	99% of 105 patients	97% of 62 patients
MidState Medical Center	96% of 57 patients	91% of 34 patients	**	100% of 44 patients	87% of 46 patients
Milford Hospital	100% of 37 patients	88% of 24 patients	**	100% of 26 patients	100% of 32 patients
New Milford Hospital	**	**	**	**	**
Norwalk Hospital	99% of 85 patients	94% of 36 patients	75% of 20 patients	98% of 55 patients	95% of 44 patients
Rockville General Hospital	94% of 31 patients	**	**	**	100% of 20 patients
Sharon Hospital	**	**	**	**	**
St Francis Hospital & Medical Center	97% of 199 patients	96% of 338 patients	68% of 111 patients	95% of 408 patients	94% of 156 patients
St Mary's Hospital	99% of 192 patients	99% of 188 patients	76% of 45 patients	96% of 199 patients	98% of 165 patients
St Vincent's Medical Center	100% of 284 patients	100% of 371 patients	97% of 111 patients	99% of 387 patients	99% of 243 patients
Stamford Hospital	98% of 132 patients	96% of 89 patients	90% of 31 patients	96% of 96 patients	98% of 84 patients
Waterbury Hospital	99% of 131 patients	98% of 133 patients	76% of 37 patients	93% of 158 patients	91% of 99 patients
William W Backus Hospital	89% of 64 patients	97% of 33 patients	**	98% of 56 patients	93% of 59 patients
Windham Community Memorial Hospital	94% of 35 patients	**	**	**	91% of 22 patients
Yale-New Haven Hospital	99% of 138 patients	99% of 256 patients	92% of 48 patients	98% of 283 patients	96% of 108 patients

* Source: CMS Hospital Compare based on data 1/1/06-12/31/06.

** Performance rates are not displayed if denominators were less than 20 during the reporting period.

Table B5

Heart Failure Performance Rates for Connecticut Hospitals		
<i>January 1, 2006 to December 31, 2006</i>		
	LVF Assessment	ACEI or ARB for LVSD
National Average Rate*	84%	82%
Connecticut Average Rate	96%	86%
Bridgeport Hospital	98% of 340 patients	87% of 140 patients
Bristol Hospital	88% of 203 patients	82% of 89 patients
Charlotte Hungerford Hospital	90% of 168 patients	78% of 49 patients
Danbury Hospital	95% of 435 patients	74% of 163 patients
Day Kimball Hospital	100% of 136 patients	90% of 41 patients
Greenwich Hospital Association	91% of 264 patients	90% of 81 patients
Griffin Hospital	100% of 231 patients	96% of 57 patients
Hartford Hospital	97% of 753 patients	80% of 341 patients
HCC at Bradley Memorial	95% of 130 patients	82% of 38 patients
HCC at New Britain General	94% of 642 patients	81% of 183 patients
Hospital of St Raphael	100% of 609 patients	99% of 170 patients
John Dempsey Hospital	93% of 196 patients	100% of 82 patients
Johnson Memorial Hospital	84% of 148 patients	89% of 27 patients
Lawrence & Memorial Hospital	99% of 400 patients	88% of 112 patients
Manchester Memorial Hospital	98% of 149 patients	92% of 48 patients
Middlesex Hospital	97% of 280 patients	90% of 90 patients
MidState Medical Center	99% of 343 patients	91% of 91 patients
Milford Hospital	99% of 193 patients	76% of 42 patients
New Milford Hospital	85% of 48 patients	**
Norwalk Hospital	98% of 384 patients	89% of 107 patients
Rockville General Hospital	97% of 116 patients	91% of 43 patients
Sharon Hospital	96% of 56 patients	**
St Francis Hospital & Medical Center	93% of 542 patients	77% of 254 patients
St Mary's Hospital	88% of 360 patients	89% of 96 patients
St Vincent's Medical Center	99% of 632 patients	97% of 207 patients
Stamford Hospital	96% of 382 patients	88% of 166 patients
Waterbury Hospital	96% of 353 patients	76% of 126 patients
William W Backus Hospital	97% of 310 patients	88% of 74 patients
Windham Community Memorial Hospital	95% of 129 patients	77% of 44 patients
Yale-New Haven Hospital	98% of 323 patients	86% of 101 patients

* Source: CMS Hospital Compare based on data 1/1/06-12/31/06.

** Performance rates are not displayed if the number of eligible patients was less than 20 during the reporting period.

Table B6

Pneumonia Performance Rates for Connecticut Hospitals			
<i>January 1, 2006 to December 31, 2006</i>			
	Oxygenation Assessment	Pneumococcal Vaccination	Timely Antibiotic
National Average Rate*	99%	71%	80%
Connecticut Average Rate	100%	79%	81%
Bridgeport Hospital	100% of 201 patients	89% of 149 patients	83% of 154 patients
Bristol Hospital	100% of 415 patients	92% of 288 patients	89% of 336 patients
Charlotte Hungerford Hospital	100% of 256 patients	96% of 192 patients	79% of 203 patients
Danbury Hospital	100% of 471 patients	68% of 364 patients	87% of 364 patients
Day Kimball Hospital	100% of 217 patients	84% of 166 patients	80% of 178 patients
Greenwich Hospital Association	100% of 245 patients	96% of 178 patients	78% of 201 patients
Griffin Hospital	100% of 226 patients	95% of 152 patients	84% of 184 patients
Hartford Hospital	100% of 720 patients	57% of 487 patients	68% of 565 patients
HCC at Bradley Memorial	100% of 127 patients	86% of 99 patients	91% of 107 patients
HCC at New Britain General	100% of 492 patients	82% of 346 patients	85% of 398 patients
Hospital of St Raphael	100% of 343 patients	94% of 259 patients	82% of 263 patients
John Dempsey Hospital	100% of 163 patients	10% of 122 patients	77% of 148 patients
Johnson Memorial Hospital	100% of 197 patients	53% of 160 patients	73% of 146 patients
Lawrence & Memorial Hospital	100% of 350 patients	94% of 231 patients	86% of 263 patients
Manchester Memorial Hospital	100% of 324 patients	81% of 234 patients	76% of 257 patients
Middlesex Hospital	100% of 460 patients	79% of 384 patients	90% of 376 patients
MidState Medical Center	100% of 267 patients	96% of 200 patients	90% of 194 patients
Milford Hospital	100% of 287 patients	93% of 212 patients	82% of 205 patients
New Milford Hospital	100% of 126 patients	67% of 94 patients	82% of 104 patients
Norwalk Hospital	100% of 299 patients	82% of 221 patients	76% of 238 patients
Rockville General Hospital	100% of 160 patients	79% of 118 patients	78% of 120 patients
Sharon Hospital	100% of 117 patients	64% of 81 patients	94% of 108 patients
St Francis Hospital & Medical Center	100% of 224 patients	82% of 152 patients	73% of 167 patients
St Mary's Hospital	99% of 328 patients	70% of 225 patients	72% of 274 patients
St Vincent's Medical Center	100% of 446 patients	86% of 293 patients	86% of 398 patients
Stamford Hospital	100% of 268 patients	86% of 196 patients	76% of 225 patients
Waterbury Hospital	99% of 254 patients	66% of 201 patients	74% of 205 patients
William W Backus Hospital	100% of 271 patients	74% of 184 patients	86% of 222 patients
Windham Community Memorial Hospital	100% of 231 patients	62% of 145 patients	89% of 187 patients
Yale-New Haven Hospital	100% of 187 patients	92% of 114 patients	87% of 150 patients

* Source: CMS Hospital Compare based on data 1/1/06-12/31/06.

** Performance rates are not displayed if the number of eligible patients was less than 20 during the reporting period.

Table B7**Hospital Performance Rates with Significant Differences between 2005 and 2006***

Hospital	Measure#	2005 Performance Rate	2006 Performance Rate	Difference	P-value
Bridgeport Hospital	PN-2	64.6	89.3	24.7	0.000
Bristol Hospital	PN-2	85.5	92.0	6.5	0.013
Bristol Hospital	PN-5b	83.7	89.3	5.6	0.036
Danbury Hospital	AMI-1	97.8	100.0	2.2	0.035
Danbury Hospital	PN-2	78.6	68.4	-10.2	0.002
Day Kimball Hospital	HF-2	94.5	100.0	5.5	0.007
Day Kimball Hospital	PN-2	68.7	83.7	15.0	0.003
Greenwich Hospital Association	PN-2	88.4	95.5	7.1	0.018
Greenwich Hospital Association	PN-5b	87.8	77.6	-10.2	0.008
Hartford Hospital	AMI-1	91.6	95.7	4.1	0.027
Hartford Hospital	AMI-2	94.1	98.4	4.3	0.000
Hartford Hospital	HF-2	92.2	97.1	4.9	0.000
Hartford Hospital	PN-2	25.4	57.5	32.1	0.000
Hartford Hospital	PN-5b	56.7	67.6	11.0	0.000
HCC at Bradley Memorial	PN-2	55.7	85.9	30.1	0.000
HCC at New Britain General	HF-2	89.9	93.8	3.9	0.016
Hospital of St Raphael	AMI-1	93.5	99.2	5.7	0.000
Hospital of St Raphael	AMI-2	97.1	100.0	2.9	0.001
Hospital of St Raphael	AMI-3	77.4	100.0	22.6	0.000
Hospital of St Raphael	AMI-5	93.4	100.0	6.6	0.000
Hospital of St Raphael	AMI-6	81.3	99.1	17.8	0.000
Hospital of St Raphael	HF-2	95.6	99.8	4.2	0.000
Hospital of St Raphael	HF-3	67.7	99.4	31.7	0.000
Hospital of St Raphael	PN-2	58.3	93.8	35.5	0.000
John Dempsey Hospital	PN-2	1.5	9.8	8.4	0.004
Johnson Memorial Hospital	PN-2	70.1	52.5	-17.6	0.004
Johnson Memorial Hospital	PN-5b	85.0	72.6	-12.4	0.021
Manchester Memorial Hospital	PN-2	47.7	81.2	33.5	0.000
Manchester Memorial Hospital	PN-5b	65.5	76.3	10.7	0.010
Middlesex Hospital	PN-2	62.5	79.2	16.6	0.000
Middlesex Hospital	PN-5b	82.6	90.4	7.8	0.001
Midstate Medical Center	HF-2	95.5	98.5	3.0	0.025
Midstate Medical Center	HF-3	77.2	91.2	14.0	0.008
Midstate Medical Center	PN-2	79.5	95.5	16.0	0.000

Hospital Performance Rates with Significant Differences from 2005 to 2006 (cont.)

Hospital	Measure#	2005 Performance Rate	2006 Performance Rate	Difference	P-value
Milford Hospital, Inc	HF-2	93.5	99.5	5.9	0.001
Milford Hospital, Inc	PN-2	78.6	93.4	14.8	0.000
New Milford Hospital	PN-2	50.8	67.0	16.2	0.019
Norwalk Hospital Association	PN-2	73.5	82.4	8.8	0.021
Rockville General Hospital	PN-2	60.7	78.8	18.1	0.003
Rockville General Hospital	PN-5b	67.5	78.3	10.9	0.047
Sharon Hospital	PN-2	33.8	64.2	30.4	0.000
St Mary's Hospital	PN-2	27.0	69.8	42.8	0.000
St Vincent's Medical Center	AMI-1	96.5	100.0	3.5	0.001
St Vincent's Medical Center	AMI-2	97.2	99.7	2.5	0.004
St Vincent's Medical Center	AMI-3	88.2	97.3	9.1	0.009
St Vincent's Medical Center	AMI-5	93.6	99.0	5.4	0.000
St Vincent's Medical Center	AMI-6	93.8	98.8	4.9	0.002
St Vincent's Medical Center	HF-2	95.5	98.9	3.4	0.000
Stamford Hospital	HF-3	94.7	88.0	-6.8	0.046
Waterbury Hospital Health Center	AMI-2	90.4	97.7	7.4	0.024
Waterbury Hospital Health Center	AMI-5	84.2	93.0	8.8	0.032
Waterbury Hospital Health Center	HF-2	87.4	96.0	8.6	0.000
Waterbury Hospital Health Center	PN-2	47.4	65.7	18.3	0.000
William W Backus Hospital	AMI-1	98.3	89.1	-9.3	0.009
William W Backus Hospital	AMI-6	100.0	93.2	-6.8	0.032
William W Backus Hospital	HF-2	100.0	97.1	-2.9	0.002
William W Backus Hospital	HF-3	100.0	87.8	-12.2	0.001
William W Backus Hospital	PN-2	62.1	74.5	12.3	0.007
William W Backus Hospital	PN-5b	78.5	86.0	7.5	0.029
Windham Comm Mem Hosp	PN-5b	78.9	89.3	10.4	0.004

*Comparisons were excluded if fewer than 20 cases per hospital-measure were eligible during 2005 or 2006.
Differences are significant if p<0.05.

# Measure	Description	# Measure	Description	# Measure	Description
AMI-1	Aspirin at Arrival	AMI-6	Beta Blocker at Arrival	PN-1	Oxygenation Assessment
AMI-2	Aspirin at Discharge	HF-2	LVF Assessment	PN-2	Pneumococcal Vaccination
AMI-3	ACEI or ARB for LVSD	HF-3	ACEI or ARB for LVSD	PN-5b	Antibiotic received within 4 hrs
AMI-5	Beta Blocker at Discharge				