

## Year 2012 Connecticut Hospitalizations

In 2012 there were 327,848 hospitalizations of Connecticut residents in Connecticut hospitals (Table H-1), not including those related to pregnancy and birth, with total charges of over 12.7 billion dollars. The age-adjusted hospitalization rate for all conditions besides pregnancy and birth decreased 2% from the previous year to 8,101 per 100,000 residents of all ages. The lowest rate was for ages 5-14 (1,637 per 100,000) and the highest for ages 65 and older (28,542 per 100,000). Overall, the median charge per hospital stay increased 8.3% to \$24,318 (unadjusted for inflation) and the median length of stay was three days. Among listed conditions and procedures, the longest median stay was for Alzheimer's disease (ten days), and the highest median charge was for non-fatal spinal injury (\$94,946).

The leading diagnosis for all ages combined, as well as for ages 45-64, was digestive diseases (Table H-5). Respiratory diseases was the leading cause of hospitalization for ages 0-4; while mental disorders was the leading cause for ages 5-14, 15-24, and 25-44. Heart disease was the leading cause of hospitalization for ages 65 and older, and for males of all ages. Digestive diseases was the leading cause of hospitalization in women. Among white residents, black residents, and Hispanic residents, the leading causes of hospitalization were, respectively, heart disease, respiratory diseases, and digestive diseases (Table H-2).

The most common type of injury among males and females, whites, blacks, and Hispanics, was an unintentional fall (Tables H-3, H-4). Hospitalization rate ratios between the sexes and race/ethnicity groups appear in Tables H-6 and H-7. High male to female, black to white, and Hispanic to white ratios occurred simultaneously within some conditions, such as diabetes, amputation with diabetes, and firearms injury. High female to male ratios marked a few conditions, such as cholelithiasis (gallstones), and high white to black ratios were present for some conditions, such as hip fracture.

Hospitalizations related to pregnancy and birth (Tables H-1 and H-2) were not used in rankings or rate ratios reported in Tables H-2, H-5, H-6, and H-7. In 2012 there were 73,431 pregnancy and birth-related discharges, with charges of more than 1.1 billion dollars. For the most accurate record of Connecticut resident births and related risk factors and outcomes, see the "Vital Statistics" section of the DPH website. For the most accurate record of Connecticut resident cancer risk factors and outcomes, see the "Tumor Registry" (CTR) section of the DPH website.