

# Connecticut Resident Age-Adjusted Rates for 62 Cause of Death Classifications by Sex Hispanic Residents, 1990

Page

## Connecticut Resident Age-Adjusted Mortality Rate (AAMR) Tables by Cause of Death and Sex, Hispanic Residents, 1990

Rates calculated using the 1940, 1970 and 2000 standard populations

1940 standard population

2

1970 standard population

4

2000 standard population

6

## Connecticut Resident Age-Adjusted Years of Potential Life Lost Rate (YPLL) Tables by Cause of Death and Sex, Hispanic Residents, 1990

Rates calculated using the 1940, 1970 and 2000 standard populations

1940 standard population

8

1970 standard population

14

2000 standard population

20

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR
*****Causes of Death*****															
All causes	558	262.4	332.1	0	326.0	340	327.0	445.0	0	441.0	218	200.6	234.7	0	230.1
Infectious diseases and parasitic disease (1-139)	61	28.7	30.5	0	25.2	45	43.3	47.2	0	37.6	16	14.7	15.5	0	.
..Septicemia (038)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
....Septicemia-related (038)	36	16.9	21.6	0	15.2	22	21.2	29.6	0	.	14	12.9	.	.	.
..HIV infection (042-044)	44	20.7	21.4	0	17.7	34	32.7	35.2	0	27.1	10	9.2	.	.	.
All cancer (140-208)	73	34.3	53.7	0	60.4	39	37.5	68.2	0	65.0	34	31.3	42.8	0	56.3
..Oral cavity or pharynx (140.0-149.9)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4.7	.	.	.	2	1.9	.	.	.	8	7.4	.	.	.
....Colon cancer (153)	8	3.8	.	.	.	2	1.9	.	.	.	6	5.5	.	.	.
....Rectal cancer (154)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Pancreatic cancer (157)	2	0.9	.	.	.	1	1.0	.	.	.	1	0.9	.	.	.
..Lung and other respiratory cancer (160-165)	18	8.5	13.8	0	.	12	11.5	.	.	.	6	5.5	.	.	.
....Lung cancer (162.2-.9)	18	8.5	13.8	0	.	12	11.5	.	.	.	6	5.5	.	.	.
..Malignant melanoma of skin (172)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Female breast cancer (174)	2	.	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Cervical cancer (180)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Ovarian cancer (183)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Prostate cancer (185)	2	.	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Bladder cancer (188)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	0.9	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Leukemia (204-208)	5	2.4	.	.	.	3	2.9	.	.	.	2	1.8	.	.	.
Diabetes mellitus (250)	15	7.1	10.8	0	.	4	3.8	.	.	.	11	10.1	.	.	.
..Diabetes-related (250)	42	19.7	31.7	0	51.2	19	18.3	32.7	0	58.4	23	21.2	30.9	0	46.0
Alzheimer's diseases (331.0)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	70.1	99.4	0	120.4	73	70.2	117.6	0	159.7	76	69.9	85.2	0	92.4
..Diseases of the heart (390-398,402,404-429)	109	51.3	73.0	0	91.2	54	51.9	86.3	0	122.3	55	50.6	62.5	0	69.0
....Heart disease-related (390-398,402,404-429)	208	97.8	132.1	0	161.7	106	101.9	159.3	0	205.6	102	93.8	108.7	0	128.8
....Coronary heart disease (CHD) (402,410-414,429.2)	72	33.9	49.8	0	63.5	40	38.5	66.4	0	85.5	32	29.4	37.3	0	48.3
....Congestive heart failure (CHF) (428.0)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
..Hypertension with or without renal disease (401,403)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
....Hypertension-related (401,403)	25	11.8	18.3	0	21.1	9	8.7	.	.	.	16	14.7	20.6	0	.
..Cerebrovascular disease (430-438)	27	12.7	17.5	0	21.1	11	10.6	.	.	.	16	14.7	16.9	0	19.0
..Atherosclerosis (440)	3	1.4	.	.	.	2	1.9	.	.	.	1	0.9	.	.	.
....Atherosclerosis-related (440)	10	4.7	.	.	11.0	4	3.8	.	.	.	6	5.5	.	.	.
..Aortic aneurysm (441)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
Pneumonia & influenza (480-487)	15	7.1	7.7	0	.	7	6.7	.	.	.	8	7.4	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	12	5.6	.	.	.	8	7.7	.	.	.	4	3.7	.	.	.
..COPD-related (490-496)	29	13.6	19.6	0	20.7	16	15.4	25.4	0	.	13	12.0	.	.	19.9

(CONTINUED)

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR
*****Causes of Death*****															
..Asthma (493)	6	2.8	.	.	.	4	3.8	.	.	.	2	1.8	.	.	.
Chronic liver disease and cirrhosis (571)	23	10.8	16.2	.	0	13.1	18	17.3	28.5	0	28.1	5	4.6	.	.
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
..Nephritis-related (580-589)	48	22.6	34.0	0	27.4	24	23.1	35.7	0	39.1	24	22.1	32.4	0	.
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
Unintentional injuries (E800-E949)	34	16.0	16.5	0	15.3	27	26.0	26.3	0	27.6	7	6.4	.	.	.
..Motor vehicle accidents (E810-E825)	21	9.9	9.2	0	7.7	18	17.3	16.3	0	.	3	2.8	.	.	.
..Falls and fall related injuries (E880-E888)	3	1.4	.	.	.	1	1.0	.	.	.	2	1.8	.	.	.
..Drowning (E830,E832,E910)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
Suicide (E950-E959)	26	12.2	13.1	0	.	22	21.2	24.6	0	.	4	3.7	.	.	.
Homicide and legal intervention (E960-E978)	32	15.0	13.7	0	9.2	27	26.0	24.2	0	.	5	4.6	.	.	.
..Homicide (E960-E969)	32	15.0	13.7	0	9.2	27	26.0	24.2	0	.	5	4.6	.	.	.
All injury (E800-E999)	95	44.7	44.8	0	31.1	79	76.0	78.1	0	56.1	16	14.7	14.4	0	.
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	9.4	8.8	0	7.7	17	16.3	15.5	0	.	3	2.8	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	4	1.9	.	.	.	2	1.9	.	.	.	2	1.8	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	18.8	18.4	1	.	32	30.8	31.7	1	.	8	7.4	.	.	.
....Suicide (E955.0-.4)	12	5.6	.	.	.	9	8.7	.	.	.	3	2.8	.	.	.
....Homicide (E965.0-.4)	26	12.2	11.4	0	.	21	20.2	19.4	0	.	5	4.6	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	2.8	.	.	.	5	4.8	.	.	.	1	0.9	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	8.0	11.6	0	13.0	14	13.5	.	.	28.0	3	2.8	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	1.4	.	.	.	3	2.9	.	.	.	.	.	.	.	.

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR
*****Causes of Death*****															
All causes	558	262.4	402.0	0	421.7	340	327.0	517.6	0	551.2	218	200.6	304.4	0	319.1
Infectious diseases and parasitic disease (1-139)	61	28.7	28.9	0	24.4	45	43.3	44.1	0	34.9	16	14.7	15.7	0	.
..Septicemia (038)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
....Septicemia-related (038)	36	16.9	27.0	0	21.7	22	21.2	35.7	0	.	14	12.9	.	.	.
..HIV infection (042-044)	44	20.7	18.3	0	14.9	34	32.7	30.3	0	22.6	10	9.2	.	.	.
All cancer (140-208)	73	34.3	68.5	0	76.5	39	37.5	87.7	0	80.5	34	31.3	54.4	0	72.7
..Oral cavity or pharynx (140.0-149.9)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4.7	.	.	.	2	1.9	.	.	.	8	7.4	.	.	.
....Colon cancer (153)	8	3.8	.	.	.	2	1.9	.	.	.	6	5.5	.	.	.
....Rectal cancer (154)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Pancreatic cancer (157)	2	0.9	.	.	.	1	1.0	.	.	.	1	0.9	.	.	.
..Lung and other respiratory cancer (160-165)	18	8.5	18.1	0	.	12	11.5	.	.	.	6	5.5	.	.	.
....Lung cancer (162.2-.9)	18	8.5	18.1	0	.	12	11.5	.	.	.	6	5.5	.	.	.
..Malignant melanoma of skin (172)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Female breast cancer (174)	2	.	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Cervical cancer (180)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Ovarian cancer (183)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Prostate cancer (185)	2	.	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Bladder cancer (188)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	0.9	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Leukemia (204-208)	5	2.4	.	.	.	3	2.9	.	.	.	2	1.8	.	.	.
Diabetes mellitus (250)	15	7.1	14.4	0	.	4	3.8	.	.	.	11	10.1	.	.	.
..Diabetes-related (250)	42	19.7	40.2	0	68.7	19	18.3	40.5	0	77.8	23	21.2	39.8	0	62.3
Alzheimer's diseases (331.0)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	70.1	138.5	0	176.9	73	70.2	159.8	0	231.2	76	69.9	122.1	0	139.9
..Diseases of the heart (390-398,402,404-429)	109	51.3	101.9	0	133.4	54	51.9	117.5	0	176.5	55	50.6	90.0	0	104.0
....Heart disease-related (390-398,402,404-429)	208	97.8	178.5	0	226.6	106	101.9	203.7	0	285.4	102	93.8	156.1	0	185.1
....Coronary heart disease (CHD) (402,410-414,429.2)	72	33.9	69.7	0	96.3	40	38.5	91.4	0	128.2	32	29.4	54.2	0	75.2
....Congestive heart failure (CHF) (428.0)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
..Hypertension with or without renal disease (401,403)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
....Hypertension-related (401,403)	25	11.8	23.1	0	28.9	9	8.7	.	.	.	16	14.7	26.3	0	.
..Cerebrovascular disease (430-438)	27	12.7	24.5	0	31.7	11	10.6	.	.	.	16	14.7	23.7	0	30.8
..Atherosclerosis (440)	3	1.4	.	.	.	2	1.9	.	.	.	1	0.9	.	.	.
....Atherosclerosis-related (440)	10	4.7	.	.	18.0	4	3.8	.	.	.	6	5.5	.	.	.
..Aortic aneurysm (441)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
Pneumonia & influenza (480-487)	15	7.1	14.7	0	.	7	6.7	.	.	.	8	7.4	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	12	5.6	.	.	.	8	7.7	.	.	.	4	3.7	.	.	.
..COPD-related (490-496)	29	13.6	26.9	0	31.0	16	15.4	34.2	0	.	13	12.0	.	.	28.2

(CONTINUED)

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR
*****Causes of Death*****															
..Asthma (493)	6	2.8	.	.	.	4	3.8	.	.	.	2	1.8	.	.	.
Chronic liver disease and cirrhosis (571)	23	10.8	17.5	0	13.6	18	17.3	29.8	0	29.4	5	4.6	.	.	.
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
..Nephritis-related (580-589)	48	22.6	39.6	0	36.9	24	23.1	40.3	0	52.9	24	22.1	39.0	0	.
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
Unintentional injuries (E800-E949)	34	16.0	16.0	0	15.6	27	26.0	24.1	0	26.7	7	6.4	.	.	.
..Motor vehicle accidents (E810-E825)	21	9.9	8.5	0	7.6	18	17.3	14.6	0	.	3	2.8	.	.	.
..Falls and fall related injuries (E880-E888)	3	1.4	.	.	.	1	1.0	.	.	.	2	1.8	.	.	.
..Drowning (E830,E832,E910)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
Suicide (E950-E959)	26	12.2	12.5	0	.	22	21.2	24.2	0	.	4	3.7	.	.	.
Homicide and legal intervention (E960-E978)	32	15.0	12.8	0	8.4	27	26.0	23.7	0	.	5	4.6	.	.	.
..Homicide (E960-E969)	32	15.0	12.8	0	8.4	27	26.0	23.7	0	.	5	4.6	.	.	.
All injury (E800-E999)	95	44.7	42.6	0	29.6	79	76.0	74.7	0	51.6	16	14.7	13.9	0	.
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	9.4	8.2	0	7.6	17	16.3	14.0	0	.	3	2.8	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	4	1.9	.	.	.	2	1.9	.	.	.	2	1.8	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	18.8	17.6	1	.	32	30.8	32.0	1	.	8	7.4	.	.	.
....Suicide (E955.0-.4)	12	5.6	.	.	.	9	8.7	.	.	.	3	2.8	.	.	.
....Homicide (E965.0-.4)	26	12.2	10.7	0	.	21	20.2	19.5	0	.	5	4.6	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	2.8	.	.	.	5	4.8	.	.	.	1	0.9	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	8.0	11.8	0	13.7	14	13.5	.	.	29.6	3	2.8	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	1.4	.	.	.	3	2.9	.	.	.	.	.	.	.	.

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR	No. of deaths	Crude rate	AAMR	Significance	1989 AAMR
*****Causes of Death*****															
All causes	558	262.4	509.0	0	556.4	340	327.0	637.8	0	716.0	218	200.6	399.7	0	433.6
Infectious diseases and parasitic disease (1-139)	61	28.7	36.7	0	30.7	45	43.3	56.4	0	42.6	16	14.7	19.8	0	
..Septicemia (038)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
....Septicemia-related (038)	36	16.9	36.4	0	31.6	22	21.2	46.3	0	.	14	12.9	.	.	.
..HIV infection (042-044)	44	20.7	22.0	0	17.9	34	32.7	37.0	0	27.3	10	9.2	.	.	.
All cancer (140-208)	73	34.3	82.5	0	94.7	39	37.5	106.0	0	96.6	34	31.3	65.6	0	92.4
..Oral cavity or pharynx (140.0-149.9)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4.7	.	.	.	2	1.9	.	.	.	8	7.4	.	.	.
....Colon cancer (153)	8	3.8	.	.	.	2	1.9	.	.	.	6	5.5	.	.	.
....Rectal cancer (154)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Pancreatic cancer (157)	2	0.9	.	.	.	1	1.0	.	.	.	1	0.9	.	.	.
..Lung and other respiratory cancer (160-165)	18	8.5	21.9	0	.	12	11.5	.	.	.	6	5.5	.	.	.
....Lung cancer (162.2-.9)	18	8.5	21.9	0	.	12	11.5	.	.	.	6	5.5	.	.	.
..Malignant melanoma of skin (172)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Female breast cancer (174)	2	.	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Cervical cancer (180)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Ovarian cancer (183)	1	.	.	.	.	.	.	.	.	.	1	0.9	.	.	.
..Prostate cancer (185)	2	.	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Bladder cancer (188)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	0.9	.	.	.	.	.	.	.	.	2	1.8	.	.	.
..Leukemia (204-208)	5	2.4	.	.	.	3	2.9	.	.	.	2	1.8	.	.	.
Diabetes mellitus (250)	15	7.1	17.8	0	.	4	3.8	.	.	.	11	10.1	.	.	.
..Diabetes-related (250)	42	19.7	47.6	0	84.5	19	18.3	47.3	0	95.0	23	21.2	47.3	0	77.1
Alzheimer's diseases (331.0)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	70.1	189.0	0	249.2	73	70.2	211.9	0	320.4	76	69.9	171.0	0	201.2
..Diseases of the heart (390-398,402,404-429)	109	51.3	139.4	0	185.1	54	51.9	157.6	0	241.7	55	50.6	125.5	0	147.0
....Heart disease-related (390-398,402,404-429)	208	97.8	246.4	0	307.2	106	101.9	269.0	0	383.0	102	93.8	223.4	0	255.5
....Coronary heart disease (CHD) (402,410-414,429.2)	72	33.9	94.5	0	135.9	40	38.5	121.9	0	179.6	32	29.4	75.2	0	107.5
....Congestive heart failure (CHF) (428.0)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
..Hypertension with or without renal disease (401,403)	1	0.5	.	.	.	1	1.0	.	.	.	.	.	.	.	.
....Hypertension-related (401,403)	25	11.8	29.0	0	37.4	9	8.7	.	.	.	16	14.7	32.8	0	.
..Cerebrovascular disease (430-438)	27	12.7	33.8	0	48.3	11	10.6	.	.	.	16	14.7	33.9	0	49.6
..Atherosclerosis (440)	3	1.4	.	.	.	2	1.9	.	.	.	1	0.9	.	.	.
....Atherosclerosis-related (440)	10	4.7	.	.	26.6	4	3.8	.	.	.	6	5.5	.	.	.
..Aortic aneurysm (441)	5	2.4	.	.	.	2	1.9	.	.	.	3	2.8	.	.	.
Pneumonia & influenza (480-487)	15	7.1	25.7	0	.	7	6.7	.	.	.	8	7.4	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	12	5.6	.	.	.	8	7.7	.	.	.	4	3.7	.	.	.
..COPD-related (490-496)	29	13.6	35.6	0	43.8	16	15.4	43.8	0	.	13	12.0	.	.	39.1

(CONTINUED)

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Number of deaths, crude mortality rate and age-adjusted mortality rate (AAMR) (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes				Male				Female						
	*****1990*****				*****1990*****				*****1990*****						
	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR	No. of deaths	Crude rate	AAMR	Signif- icance	1989 AAMR
*****Causes of Death*****															
..Asthma (493)	6	2.8	.	.	.	4	3.8	.	.	.	2	1.8	.	.	.
Chronic liver disease and cirrhosis (571)	23	10.8	20.5	0	14.7	18	17.3	34.0	0	31.6	5	4.6	.	.	.
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
..Nephritis-related (580-589)	48	22.6	44.9	0	48.2	24	23.1	46.1	0	69.9	24	22.1	43.6	0	.
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	5.6	.	.	.	7	6.7	.	.	.	5	4.6	.	.	.
Unintentional injuries (E800-E949)	34	16.0	16.6	0	16.9	27	26.0	25.1	0	28.2	7	6.4	.	.	.
..Motor vehicle accidents (E810-E825)	21	9.9	8.2	0	7.2	18	17.3	14.7	0	.	3	2.8	.	.	.
..Falls and fall related injuries (E880-E888)	3	1.4	.	.	.	1	1.0	.	.	.	2	1.8	.	.	.
..Drowning (E830,E832,E910)	2	0.9	.	.	.	2	1.9	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	0.5	.	.	.	.	.	.	.	.	1	0.9	.	.	.
Suicide (E950-E959)	26	12.2	14.5	0	.	22	21.2	28.8	0	.	4	3.7	.	.	.
Homicide and legal intervention (E960-E978)	32	15.0	12.8	0	9.0	27	26.0	23.8	0	.	5	4.6	.	.	.
..Homicide (E960-E969)	32	15.0	12.8	0	9.0	27	26.0	23.8	0	.	5	4.6	.	.	.
All injury (E800-E999)	95	44.7	45.3	0	32.8	79	76.0	80.7	0	57.2	16	14.7	14.5	0	.
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	9.4	7.8	0	7.2	17	16.3	14.0	0	.	3	2.8	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	4	1.9	.	.	.	2	1.9	.	.	.	2	1.8	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	18.8	17.8	1	.	32	30.8	32.7	1	.	8	7.4	.	.	.
....Suicide (E955.0-.4)	12	5.6	.	.	.	9	8.7	.	.	.	3	2.8	.	.	.
....Homicide (E965.0-.4)	26	12.2	10.7	0	.	21	20.2	19.6	0	.	5	4.6	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	2.8	.	.	.	5	4.8	.	.	.	1	0.9	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	8.0	13.7	0	14.5	14	13.5	.	.	31.3	3	2.8	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	1.4	.	.	.	3	2.9	.	.	.	.	.	.	.	.

\* - Crude and age-adjusted mortality rates are calculated per 100,000 population.  
 . - The AAMR is not reported due to a small number of deaths (<15).  
 1 - The 1990 AAMR is significantly different (prob<0.05) from the 1989 AAMR.  
 0 - The 1990 AAMR is not significantly different (prob>0.05) from the 1989 AAMR.

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	558	371	447	10,329	14,474	4968.3	3876.9	1	7240.3	5869.2	1
Infectious diseases and parasitic disease (1-139)	61	55	57	1,625	2,190	776.6	612.7	0	1024.4	818.5	0
..Septicemia (038)	12	8	8	215	295	.	.	0	.	.	.
...Septicemia-related (038)	36	22	27	512	765	259.2	107.7	0	408.3	187.0	0
..HIV infection (042-044)	44	43	44	1,285	1,722	605.9	497.7	0	792.2	654.3	0
All cancer (140-208)	73	36	53	505	968	316.7	487.0	0	662.9	889.2	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4	4	20	60	.	.	.	.	.	.
...Colon cancer (153)	8	4	4	20	60	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	2	1	2	3	20	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	18	6	13	75	173	53.7	.	0	130.9	.	0
...Lung cancer (162.2-.9)	18	6	13	75	173	53.7	.	0	130.9	.	0
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	5	3	4	68	100	.	.	.	.	.	.
Diabetes mellitus (250)	15	7	11	83	178	48.2	.	0	122.3	.	0
..Diabetes-related (250)	42	22	34	230	525	158.8	177.5	0	393.7	470.7	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	63	94	990	1,782	609.3	601.1	0	1181.9	1184.7	0
..Diseases of the heart (390-398,402,404-429)	109	46	69	670	1,248	422.9	398.2	0	847.0	840.4	0
...Heart disease-related (390-398,402,404-429)	208	105	138	1,969	3,196	1112.5	980.0	0	1921.9	1836.5	0
...Coronary heart disease (CHD) (402,410-414,429.2)	72	28	46	340	720	240.1	207.0	0	536.5	480.6	0
...Congestive heart failure (CHF) (428.0)	5	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	25	13	20	163	330	120.0	101.8	0	247.9	211.2	0
..Cerebrovascular disease (430-438)	27	11	16	240	377	135.6	160.2	0	226.8	260.0	0
..Atherosclerosis (440)	3	2	2	20	40	.	.	.	.	.	.
...Atherosclerosis-related (440)	10	3	4	23	55	.	11.8	.	.	58.2	.
..Aortic aneurysm (441)	5	.	2	0	10	.	.	.	.	.	.
Pneumonia & influenza (480-487)	15	2	3	25	53	19.3	.	0	40.8	.	0
Chronic obstructive pulmonary disease (COPD) (490-496)	12	8	9	197	285	.	.	.	.	.	.
..COPD-related (490-496)	29	12	18	237	387	130.7	72.6	0	234.0	154.3	0
..Asthma (493)	6	6	6	187	247	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	23	17	22	303	500	189.9	143.0	0	319.8	266.7	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	9	12	215	322	.	.	.	.	.	.
..Nephritis-related (580-589)	48	32	42	609	994	325.7	140.9	0	588.1	303.3	0
...End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	9	12	215	322	.	.	.	.	.	.
Unintentional injuries (E800-E949)	34	32	33	1,162	1,490	533.4	422.7	0	666.6	541.2	0
..Motor vehicle accidents (E810-E825)	21	21	21	835	1,045	379.0	252.3	0	454.7	316.1	0
..Falls and fall related injuries (E880-E888)	3	1	2	3	20	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	26	24	25	750	993	360.1	.	0	463.2	.	0
Homicide and legal intervention (E960-E978)	32	31	31	1,257	1,567	557.7	311.0	0	664.3	389.7	0
..Homicide (E960-E969)	32	31	31	1,257	1,567	557.7	311.0	0	664.3	389.7	0
All injury (E800-E999)	95	90	92	3,261	4,171	1495.7	927.7	0	1851.2	1183.4	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	20	20	802	1,002	364.9	252.3	0	437.3	316.1	0
..All falls (E880-E886,E888,E957,E968.1,E987)	4	2	3	25	53	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	38	39	1,435	1,818	658.3	.	1	801.2	.	1
...Suicide (E955.0-.4)	12	11	12	373	485	.	.	.	.	.	.
...Homicide (E965.0-.4)	26	25	25	983	1,233	445.8	.	0	533.6	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	5	5	148	198	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	13	17	268	423	161.8	136.7	0	258.9	260.0	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	340	257	295	7,251	10,041	7190.4	5806.0	0	10329.8	8619.6	0
Infectious diseases and parasitic disease (1-139)	45	42	43	1,237	1,665	1219.0	955.6	0	1607.1	1267.2	0
..Septicemia (038)	7	5	5	148	198	.	.	.	.	.	.
...Septicemia-related (038)	22	16	18	387	562	399.5	.	0	614.0	.	0
..HIV infection (042-044)	34	33	34	965	1,302	945.4	762.4	0	1252.9	1001.1	0
All cancer (140-208)	39	21	29	258	513	380.6	474.4	0	807.3	946.3	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	2	1	1	8	18	.	.	.	.	.	.
...Colon cancer (153)	2	1	1	8	18	.	.	.	.	.	.
...Rectal cancer (154)	.	.	.	.	.	.	.	.	.	.	.
..Pancreatic cancer (157)	1	.	1	0	8	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	12	5	8	58	120	.	.	.	.	.	.
...Lung cancer (162.2-.9)	12	5	8	58	120	.	.	.	.	.	.
..Malignant melanoma of skin (172)	.	.	.	.	.	.	.	.	.	.	.
..Female breast cancer (174)	.	.	.	.	.	.	.	.	.	.	.
..Cervical cancer (180)	.	.	.	.	.	.	.	.	.	.	.
..Ovarian cancer (183)	.	.	.	.	.	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	.	.	.	.	.	.	.	.	.	.	.
..Leukemia (204-208)	3	1	2	23	35	.	.	.	.	.	.
Diabetes mellitus (250)	4	2	4	40	70	.	.	.	.	.	.
..Diabetes-related (250)	19	11	17	143	288	192.0	210.2	0	434.6	549.0	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	73	37	51	593	1,043	753.3	805.6	0	1446.1	1543.6	0
..Diseases of the heart (390-398,402,404-429)	54	28	38	445	780	569.0	571.5	0	1082.2	1131.4	0
...Heart disease-related (390-398,402,404-429)	106	66	82	1,257	2,007	1492.6	1318.3	0	2530.7	2405.4	0
...Coronary heart disease (CHD) (402,410-414,429.2)	40	19	28	263	500	366.7	305.8	0	750.4	650.4	0
...Congestive heart failure (CHF) (428.0)	2	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	9	5	8	78	140	.	.	.	.	.	.
..Cerebrovascular disease (430-438)	11	4	7	85	143	.	.	.	.	.	.
..Atherosclerosis (440)	2	1	1	3	13	.	.	.	.	.	.
...Atherosclerosis-related (440)	4	1	2	3	15	.	.	.	.	.	.
..Aortic aneurysm (441)	2	.	1	0	8	.	.	.	.	.	.
Pneumonia & influenza (480-487)	7	2	3	25	53	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	8	6	7	157	225	.	.	.	.	.	.
..COPD-related (490-496)	16	8	11	177	275	182.3	.	0	324.8	.	0
..Asthma (493)	4	4	4	147	187	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	18	14	18	240	405	338.6	304.4	0	580.9	570.5	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	7	6	7	162	230	.	.	.	.	.	.
..Nephritis-related (580-589)	24	17	22	390	592	411.1	205.4	0	682.4	445.8	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	7	6	7	162	230	.	.	.	.	.	.
Unintentional injuries (E800-E949)	27	26	27	952	1,220	883.0	808.0	0	1102.1	1033.2	0
..Motor vehicle accidents (E810-E825)	18	18	18	687	867	632.8	.	0	768.3	.	0
..Falls and fall related injuries (E880-E888)	1	.	1	0	8	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	.	.	.	.	.	.	.	.	.	.	.
Suicide (E950-E959)	22	20	21	595	798	600.2	.	0	788.5	.	0
Homicide and legal intervention (E960-E978)	27	26	26	1,089	1,349	975.3	.	0	1156.5	.	0
..Homicide (E960-E969)	27	26	26	1,089	1,349	975.3	.	0	1156.5	.	0
All injury (E800-E999)	79	75	77	2,729	3,489	2550.2	1681.1	0	3165.2	2154.4	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	17	17	17	655	825	603.7	.	0	732.1	.	0
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	2	23	40	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	32	30	31	1,145	1,448	1071.7	.	1	1309.6	.	1
...Suicide (E955.0-.4)	9	8	9	250	333	.	.	.	.	.	.
...Homicide (E965.0-.4)	21	20	20	815	1,015	751.4	.	0	894.6	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	5	4	4	115	155	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	14	10	14	175	300	.	290.8	.	.	556.0	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	218	114	152	3,078	4,433	2874.5	2043.5	0	4369.9	3307.2	0
Infectious diseases and parasitic disease (1-139)	16	13	14	388	525	363.1	.	0	482.6	.	0
..Septicemia (038)	5	3	3	68	98	.	.	.	.	.	.
...Septicemia-related (038)	14	6	9	125	203	.	.	.	.	.	.
..HIV infection (042-044)	10	10	10	320	420	.	.	.	.	.	.
All cancer (140-208)	34	15	24	248	455	261.6	505.3	0	538.3	850.6	0
..Oral cavity or pharynx (140.0-149.9)	.	.	.	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	8	3	3	13	43	.	.	.	.	.	.
...Colon cancer (153)	6	3	3	13	43	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	1	1	1	3	13	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	6	1	5	18	53	.	.	.	.	.	.
...Lung cancer (162.2-.9)	6	1	5	18	53	.	.	.	.	.	.
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	.	.	.	.	.	.	.	.	.	.	.
..Bladder cancer (188)	.	.	.	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	2	2	2	45	65	.	.	.	.	.	.
Diabetes mellitus (250)	11	5	7	43	108	.	.	.	.	.	.
..Diabetes-related (250)	23	11	17	88	238	128.4	149.2	0	357.4	406.0	0
Alzheimer's diseases (331.0)	.	.	.	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	76	26	43	397	740	480.6	413.1	0	950.4	870.5	0
..Diseases of the heart (390-398,402,404-429)	55	18	31	225	468	291.1	234.5	0	637.7	580.6	0
...Heart disease-related (390-398,402,404-429)	102	39	56	712	1,189	770.4	658.7	0	1381.0	1321.5	0
...Coronary heart disease (CHD) (402,410-414,429.2)	32	9	18	78	220	126.2	116.6	0	347.2	331.7	0
...Congestive heart failure (CHF) (428.0)	3	.	.	0	0	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	.	.	.	.	.	.	.	.	.	.	.
...Hypertension-related (401,403)	16	8	12	85	190	121.8	.	0	275.4	.	0
..Cerebrovascular disease (430-438)	16	7	9	155	235	165.8	124.2	0	261.2	197.5	0
..Atherosclerosis (440)	1	1	1	18	28	.	.	.	.	.	.
...Atherosclerosis-related (440)	6	2	2	20	40	.	.	.	.	.	.
..Aortic aneurysm (441)	3	.	1	0	3	.	.	.	.	.	.
Pneumonia & influenza (480-487)	8	.	.	0	0	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	4	2	2	40	60	.	.	.	.	.	.
..COPD-related (490-496)	13	4	7	60	113	.	116.9	.	.	207.7	.
..Asthma (493)	2	2	2	40	60	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	5	3	4	63	95	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1940 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	5	3	5	53	93	.	.	.	.	.	.
..Nephritis-related (580-589)	24	15	20	220	402	244.9	.	0	499.2	.	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	5	3	5	53	93	.	.	.	.	.	.
Unintentional injuries (E800-E949)	7	6	6	210	270	.	.	.	.	.	.
..Motor vehicle accidents (E810-E825)	3	3	3	148	178	.	.	.	.	.	.
..Falls and fall related injuries (E880-E888)	2	1	1	3	13	.	.	.	.	.	.
..Drowning (E830,E832,E910)	.	.	.	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	4	4	4	155	195	.	.	.	.	.	.
Homicide and legal intervention (E960-E978)	5	5	5	168	218	.	.	.	.	.	.
..Homicide (E960-E969)	5	5	5	168	218	.	.	.	.	.	.
All injury (E800-E999)	16	15	15	533	683	472.7	.	0	585.7	.	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	3	3	3	148	178	.	.	.	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	1	3	13	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	8	8	8	290	370	.	.	.	.	.	.
....Suicide (E955.0-.4)	3	3	3	123	153	.	.	.	.	.	.
....Homicide (E965.0-.4)	5	5	5	168	218	.	.	.	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	1	1	1	33	43	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	3	3	3	93	123	.	.	.	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	.	.	.	.	.	.	.	.	.	.	.

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	558	371	447	10,329	14,474	4681.4	3681.3	1	6960.1	5715.6	0
Infectious diseases and parasitic disease (1-139)	61	55	57	1,625	2,190	661.0	527.2	0	871.0	704.4	0
..Septicemia (038)	12	8	8	215	295	.	.	.	.	.	.
...Septicemia-related (038)	36	22	27	512	765	232.2	96.3	0	382.9	179.1	0
..HIV infection (042-044)	44	43	44	1,285	1,722	510.6	415.2	0	665.0	544.0	0
All cancer (140-208)	73	36	53	505	968	309.0	476.3	0	691.6	910.1	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4	4	20	60	.	.	.	.	.	.
...Colon cancer (153)	8	4	4	20	60	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	2	1	2	3	20	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	18	6	13	75	173	54.6	.	0	142.7	.	0
...Lung cancer (162.2-.9)	18	6	13	75	173	54.6	.	0	142.7	.	0
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	5	3	4	68	100	.	.	.	.	.	.
Diabetes mellitus (250)	15	7	11	83	178	47.7	.	0	132.8	.	0
..Diabetes-related (250)	42	22	34	230	525	162.4	177.0	0	429.4	514.8	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	63	94	990	1,782	595.3	575.2	0	1219.8	1217.6	0
..Diseases of the heart (390-398,402,404-429)	109	46	69	670	1,248	409.7	384.2	0	873.7	877.5	0
...Heart disease-related (390-398,402,404-429)	208	105	138	1,969	3,196	1041.2	916.9	0	1890.3	1851.1	0
...Coronary heart disease (CHD) (402,410-414,429.2)	72	28	46	340	720	240.6	205.2	0	571.0	515.5	0
...Congestive heart failure (CHF) (428.0)	5	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	25	13	20	163	330	121.8	103.4	0	261.9	224.7	0
..Cerebrovascular disease (430-438)	27	11	16	240	377	137.3	147.6	0	234.4	248.1	0
..Atherosclerosis (440)	3	2	2	20	40	.	.	.	.	.	.
...Atherosclerosis-related (440)	10	3	4	23	55	.	13.8	.	.	69.9	.
..Aortic aneurysm (441)	5	.	2	0	10	.	.	.	.	.	.
Pneumonia & influenza (480-487)	15	2	3	25	53	20.0	.	0	43.6	.	0
Chronic obstructive pulmonary disease (COPD) (490-496)	12	8	9	197	285	.	.	.	.	.	.
..COPD-related (490-496)	29	12	18	237	387	129.1	69.5	0	240.8	160.1	0
..Asthma (493)	6	6	6	187	247	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	23	17	22	303	500	175.5	144.2	0	307.3	273.6	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	9	12	215	322	.	.	.	.	.	.
..Nephritis-related (580-589)	48	32	42	609	994	309.4	140.4	0	591.4	321.0	0
...End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	9	12	215	322	.	.	.	.	.	.
Unintentional injuries (E800-E949)	34	32	33	1,162	1,490	511.9	407.6	0	631.4	517.2	0
..Motor vehicle accidents (E810-E825)	21	21	21	835	1,045	366.9	248.7	0	431.5	309.3	0
..Falls and fall related injuries (E880-E888)	3	1	2	3	20	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	26	24	25	750	993	323.9	.	0	413.3	.	0
Homicide and legal intervention (E960-E978)	32	31	31	1,257	1,567	519.0	295.9	0	606.7	364.1	0
..Homicide (E960-E969)	32	31	31	1,257	1,567	519.0	295.9	0	606.7	364.1	0
All injury (E800-E999)	95	90	92	3,261	4,171	1394.6	871.2	0	1701.8	1097.1	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	20	20	802	1,002	356.4	248.7	0	418.7	309.3	0
..All falls (E880-E886,E888,E957,E968.1,E987)	4	2	3	25	53	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	38	39	1,435	1,818	610.9	.	1	733.6	.	1
...Suicide (E955.0-.4)	12	11	12	373	485	.	.	.	.	.	.
...Homicide (E965.0-.4)	26	25	25	983	1,233	411.6	.	0	484.3	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	5	5	148	198	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	13	17	268	423	146.8	138.8	0	242.3	269.5	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	340	257	295	7,251	10,041	6706.3	5437.6	0	9803.1	8260.0	0
Infectious diseases and parasitic disease (1-139)	45	42	43	1,237	1,665	1031.1	815.5	0	1357.2	1080.8	0
..Septicemia (038)	7	5	5	148	198	.	.	.	.	.	.
...Septicemia-related (038)	22	16	18	387	562	361.2	.	0	575.6	.	0
..HIV infection (042-044)	34	33	34	965	1,302	790.4	630.1	0	1047.9	824.8	0
All cancer (140-208)	39	21	29	258	513	388.3	454.1	0	859.8	967.5	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	2	1	1	8	18	.	.	.	.	.	.
...Colon cancer (153)	2	1	1	8	18	.	.	.	.	.	.
...Rectal cancer (154)	.	.	.	.	.	.	.	.	.	.	.
..Pancreatic cancer (157)	1	.	1	0	8	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	12	5	8	58	120	.	.	.	.	.	.
...Lung cancer (162.2-.9)	12	5	8	58	120	.	.	.	.	.	.
..Malignant melanoma of skin (172)	.	.	.	.	.	.	.	.	.	.	.
..Female breast cancer (174)	.	.	.	.	.	.	.	.	.	.	.
..Cervical cancer (180)	.	.	.	.	.	.	.	.	.	.	.
..Ovarian cancer (183)	.	.	.	.	.	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	.	.	.	.	.	.	.	.	.	.	.
..Leukemia (204-208)	3	1	2	23	35	.	.	.	.	.	.
Diabetes mellitus (250)	4	2	4	40	70	.	.	.	.	.	.
..Diabetes-related (250)	19	11	17	143	288	188.6	213.3	0	461.1	600.6	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	73	37	51	593	1,043	722.3	756.0	0	1474.2	1565.0	0
..Diseases of the heart (390-398,402,404-429)	54	28	38	445	780	543.4	534.4	0	1097.6	1153.7	0
...Heart disease-related (390-398,402,404-429)	106	66	82	1,257	2,007	1378.6	1218.4	0	2455.3	2397.1	0
...Coronary heart disease (CHD) (402,410-414,429.2)	40	19	28	263	500	357.8	291.9	0	779.1	677.1	0
...Congestive heart failure (CHF) (428.0)	2	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	9	5	8	78	140	.	.	.	.	.	.
..Cerebrovascular disease (430-438)	11	4	7	85	143	.	.	.	.	.	.
..Atherosclerosis (440)	2	1	1	3	13	.	.	.	.	.	.
...Atherosclerosis-related (440)	4	1	2	3	15	.	.	.	.	.	.
..Aortic aneurysm (441)	2	.	1	0	8	.	.	.	.	.	.
Pneumonia & influenza (480-487)	7	2	3	25	53	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	8	6	7	157	225	.	.	.	.	.	.
..COPD-related (490-496)	16	8	11	177	275	179.3	.	0	334.1	.	0
..Asthma (493)	4	4	4	147	187	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	18	14	18	240	405	321.3	307.3	0	568.2	586.2	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	7	6	7	162	230	.	.	.	.	.	.
..Nephritis-related (580-589)	24	17	22	390	592	370.9	206.4	0	652.7	472.3	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	7	6	7	162	230	.	.	.	.	.	.
Unintentional injuries (E800-E949)	27	26	27	952	1,220	826.0	772.3	0	1016.7	976.5	0
..Motor vehicle accidents (E810-E825)	18	18	18	687	867	594.7	.	0	708.5	.	0
..Falls and fall related injuries (E880-E888)	1	.	1	0	8	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	.	.	.	.	.	.	.	.	.	.	.
Suicide (E950-E959)	22	20	21	595	798	541.6	.	0	707.9	.	0
Homicide and legal intervention (E960-E978)	27	26	26	1,089	1,349	927.0	.	0	1079.4	.	0
..Homicide (E960-E969)	27	26	26	1,089	1,349	927.0	.	0	1079.4	.	0
All injury (E800-E999)	79	75	77	2,729	3,489	2376.5	1556.6	0	2908.5	1969.6	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	17	17	17	655	825	573.0	.	0	682.0	.	0
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	2	23	40	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	32	30	31	1,145	1,448	1015.2	.	1	1226.0	.	1
...Suicide (E955.0-.4)	9	8	9	250	333	.	.	.	.	.	.
...Homicide (E965.0-.4)	21	20	20	815	1,015	713.0	.	0	835.2	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	5	4	4	115	155	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	14	10	14	175	300	.	295.5	.	.	577.1	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	218	114	152	3,078	4,433	2768.8	2011.8	0	4322.3	3357.2	0
Infectious diseases and parasitic disease (1-139)	16	13	14	388	525	314.7	.	0	418.9	.	0
..Septicemia (038)	5	3	3	68	98	.	.	.	.	.	.
...Septicemia-related (038)	14	6	9	125	203	.	.	.	.	.	.
..HIV infection (042-044)	10	10	10	320	420	.	.	.	.	.	.
All cancer (140-208)	34	15	24	248	455	239.9	505.5	0	545.8	874.2	0
..Oral cavity or pharynx (140.0-149.9)	.	.	.	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	8	3	3	13	43	.	.	.	.	.	.
...Colon cancer (153)	6	3	3	13	43	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	1	1	1	3	13	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	6	1	5	18	53	.	.	.	.	.	.
...Lung cancer (162.2-.9)	6	1	5	18	53	.	.	.	.	.	.
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	.	.	.	.	.	.	.	.	.	.	.
..Bladder cancer (188)	.	.	.	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	2	2	2	45	65	.	.	.	.	.	.
Diabetes mellitus (250)	11	5	7	43	108	.	.	.	.	.	.
..Diabetes-related (250)	23	11	17	88	238	138.4	145.5	0	401.8	443.9	0
Alzheimer's diseases (331.0)	.	.	.	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	76	26	43	397	740	482.6	409.5	0	999.0	918.9	0
..Diseases of the heart (390-398,402,404-429)	55	18	31	225	468	289.2	241.8	0	675.2	635.3	0
...Heart disease-related (390-398,402,404-429)	102	39	56	712	1,189	738.7	631.4	0	1391.9	1364.0	0
...Coronary heart disease (CHD) (402,410-414,429.2)	32	9	18	78	220	135.3	126.0	0	388.1	376.1	0
...Congestive heart failure (CHF) (428.0)	3	.	.	0	0	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	.	.	.	.	.	.	.	.	.	.	.
...Hypertension-related (401,403)	16	8	12	85	190	125.6	.	0	296.4	.	0
..Cerebrovascular disease (430-438)	16	7	9	155	235	170.0	110.4	0	269.1	182.9	0
..Atherosclerosis (440)	1	1	1	18	28	.	.	.	.	.	.
...Atherosclerosis-related (440)	6	2	2	20	40	.	.	.	.	.	.
..Aortic aneurysm (441)	3	.	1	0	3	.	.	.	.	.	.
Pneumonia & influenza (480-487)	8	.	.	0	0	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	4	2	2	40	60	.	.	.	.	.	.
..COPD-related (490-496)	13	4	7	60	113	.	110.2	.	.	205.9	.
..Asthma (493)	2	2	2	40	60	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	5	3	4	63	95	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (&lt;15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 1970 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	5	3	5	53	93	.	.	.	.	.	.
..Nephritis-related (580-589)	24	15	20	220	402	250.8	.	0	533.2	.	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	5	3	5	53	93	.	.	.	.	.	.
Unintentional injuries (E800-E949)	7	6	6	210	270	.	.	.	.	.	.
..Motor vehicle accidents (E810-E825)	3	3	3	148	178	.	.	.	.	.	.
..Falls and fall related injuries (E880-E888)	2	1	1	3	13	.	.	.	.	.	.
..Drowning (E830,E832,E910)	.	.	.	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	4	4	4	155	195	.	.	.	.	.	.
Homicide and legal intervention (E960-E978)	5	5	5	168	218	.	.	.	.	.	.
..Homicide (E960-E969)	5	5	5	168	218	.	.	.	.	.	.
All injury (E800-E999)	16	15	15	533	683	437.2	.	0	535.3	.	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	3	3	3	148	178	.	.	.	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	1	3	13	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	8	8	8	290	370	.	.	.	.	.	.
...Suicide (E955.0-.4)	3	3	3	123	153	.	.	.	.	.	.
...Homicide (E965.0-.4)	5	5	5	168	218	.	.	.	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	1	1	1	33	43	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	3	3	3	93	123	.	.	.	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	.	.	.	.	.	.	.	.	.	.	.

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	558	371	447	10,329	14,474	5077.1	4049.7	0	7530.4	6256.2	0
Infectious diseases and parasitic disease (1-139)	61	55	57	1,625	2,190	809.7	632.9	0	1061.0	841.2	0
..Septicemia (038)	12	8	8	215	295	.	.	0	.	.	.
...Septicemia-related (038)	36	22	27	512	765	281.7	126.7	0	447.0	221.9	0
..HIV infection (042-044)	44	43	44	1,285	1,722	631.4	511.1	0	818.3	665.8	0
All cancer (140-208)	73	36	53	505	968	355.8	547.7	0	756.5	1010.4	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	10	4	4	20	60	.	.	.	.	.	.
...Colon cancer (153)	8	4	4	20	60	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	2	1	2	3	20	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	18	6	13	75	173	65.9	.	0	160.2	.	0
...Lung cancer (162.2-.9)	18	6	13	75	173	65.9	.	0	160.2	.	0
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	5	3	4	68	100	.	.	.	.	.	.
Diabetes mellitus (250)	15	7	11	83	178	52.6	.	0	138.5	.	0
..Diabetes-related (250)	42	22	34	230	525	187.3	212.9	0	462.4	565.6	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	149	63	94	990	1,782	694.4	685.1	0	1363.8	1375.6	0
..Diseases of the heart (390-398,402,404-429)	109	46	69	670	1,248	480.6	459.7	0	976.4	985.3	0
...Heart disease-related (390-398,402,404-429)	208	105	138	1,969	3,196	1260.5	1073.1	0	2187.8	2072.6	0
...Coronary heart disease (CHD) (402,410-414,429.2)	72	28	46	340	720	286.3	249.5	0	638.4	579.3	0
...Congestive heart failure (CHF) (428.0)	5	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	25	13	20	163	330	147.2	125.0	0	299.4	257.6	0
..Cerebrovascular disease (430-438)	27	11	16	240	377	151.8	182.1	0	258.0	296.7	0
..Atherosclerosis (440)	3	2	2	20	40	.	.	.	.	.	.
...Atherosclerosis-related (440)	10	3	4	23	55	.	13.6	.	.	70.1	.
..Aortic aneurysm (441)	5	.	2	0	10	.	.	.	.	.	.
Pneumonia & influenza (480-487)	15	2	3	25	53	23.2	.	0	48.3	.	0
Chronic obstructive pulmonary disease (COPD) (490-496)	12	8	9	197	285	.	.	.	.	.	.
..COPD-related (490-496)	29	12	18	237	387	143.0	86.2	0	263.2	186.0	0
..Asthma (493)	6	6	6	187	247	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	23	17	22	303	500	229.9	172.5	0	381.8	313.3	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Both Sexes										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	12	9	12	215	322	.	.	.	.	.	.
..Nephritis-related (580-589)	48	32	42	609	994	354.6	171.2	0	650.9	368.4	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	12	9	12	215	322	.	.	.	.	.	.
Unintentional injuries (E800-E949)	34	32	33	1,162	1,490	496.9	398.7	0	619.1	513.5	0
..Motor vehicle accidents (E810-E825)	21	21	21	835	1,045	341.8	221.4	0	405.0	278.8	0
..Falls and fall related injuries (E880-E888)	3	1	2	3	20	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	26	24	25	750	993	358.1	.	0	459.5	.	0
Homicide and legal intervention (E960-E978)	32	31	31	1,257	1,567	483.1	301.2	0	565.8	375.7	0
..Homicide (E960-E969)	32	31	31	1,257	1,567	483.1	301.2	0	565.8	375.7	0
All injury (E800-E999)	95	90	92	3,261	4,171	1381.5	905.3	0	1700.0	1154.1	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	20	20	20	802	1,002	328.2	221.4	0	388.4	278.8	0
..All falls (E880-E886,E888,E957,E968.1,E987)	4	2	3	25	53	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	40	38	39	1,435	1,818	572.9	.	1	691.1	.	1
...Suicide (E955.0-.4)	12	11	12	373	485	.	.	.	.	.	.
...Homicide (E965.0-.4)	26	25	25	983	1,233	379.6	.	0	447.1	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	6	5	5	148	198	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	17	13	17	268	423	191.9	164.3	0	304.6	304.1	0
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	340	257	295	7,251	10,041	7375.7	6052.1	0	10740.8	9141.5	0
Infectious diseases and parasitic disease (1-139)	45	42	43	1,237	1,665	1287.6	987.6	0	1682.3	1302.1	0
..Septicemia (038)	7	5	5	148	198	.	.	.	.	.	.
...Septicemia-related (038)	22	16	18	387	562	427.0	.	0	659.9	.	0
..HIV infection (042-044)	34	33	34	965	1,302	1009.0	787.9	0	1324.4	1023.6	0
All cancer (140-208)	39	21	29	258	513	445.3	526.7	0	942.9	1066.4	0
..Oral cavity or pharynx (140.0-149.9)	2	2	2	15	35	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	2	1	1	8	18	.	.	.	.	.	.
...Colon cancer (153)	2	1	1	8	18	.	.	.	.	.	.
...Rectal cancer (154)	.	.	.	.	.	.	.	.	.	.	.
..Pancreatic cancer (157)	1	.	1	0	8	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	12	5	8	58	120	.	.	.	.	.	.
...Lung cancer (162.2-.9)	12	5	8	58	120	.	.	.	.	.	.
..Malignant melanoma of skin (172)	.	.	.	.	.	.	.	.	.	.	.
..Female breast cancer (174)	.	.	.	.	.	.	.	.	.	.	.
..Cervical cancer (180)	.	.	.	.	.	.	.	.	.	.	.
..Ovarian cancer (183)	.	.	.	.	.	.	.	.	.	.	.
..Prostate cancer (185)	2	1	1	8	18	.	.	.	.	.	.
..Bladder cancer (188)	1	.	.	0	0	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	.	.	.	.	.	.	.	.	.	.	.
..Leukemia (204-208)	3	1	2	23	35	.	.	.	.	.	.
Diabetes mellitus (250)	4	2	4	40	70	.	.	.	.	.	.
..Diabetes-related (250)	19	11	17	143	288	224.0	253.2	0	510.1	659.6	0
Alzheimer's diseases (331.0)	1	.	.	0	0	.	.	.	.	.	.
Cardiovascular disease (390-459)	73	37	51	593	1,043	857.8	924.9	0	1661.5	1808.6	0
..Diseases of the heart (390-398,402,404-429)	54	28	38	445	780	648.4	649.6	0	1243.6	1318.9	0
...Heart disease-related (390-398,402,404-429)	106	66	82	1,257	2,007	1714.3	1411.1	0	2902.5	2671.9	0
...Coronary heart disease (CHD) (402,410-414,429.2)	40	19	28	263	500	437.4	369.4	0	891.5	787.8	0
...Congestive heart failure (CHF) (428.0)	2	1	2	3	20	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	1	1	1	8	18	.	.	.	.	.	.
...Hypertension-related (401,403)	9	5	8	78	140	.	.	.	.	.	.
..Cerebrovascular disease (430-438)	11	4	7	85	143	.	.	.	.	.	.
..Atherosclerosis (440)	2	1	1	3	13	.	.	.	.	.	.
...Atherosclerosis-related (440)	4	1	2	3	15	.	.	.	.	.	.
..Aortic aneurysm (441)	2	.	1	0	8	.	.	.	.	.	.
Pneumonia & influenza (480-487)	7	2	3	25	53	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	8	6	7	157	225	.	.	.	.	.	.
..COPD-related (490-496)	16	8	11	177	275	187.6	.	0	348.9	.	0
..Asthma (493)	4	4	4	147	187	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	18	14	18	240	405	417.8	367.2	0	703.5	670.4	0

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Male										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	7	6	7	162	230	.	.	.	.	.	.
..Nephritis-related (580-589)	24	17	22	390	592	446.9	251.9	0	752.0	542.5	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	7	6	7	162	230	.	.	.	.	.	.
Unintentional injuries (E800-E949)	27	26	27	952	1,220	830.1	766.8	0	1032.2	984.2	0
..Motor vehicle accidents (E810-E825)	18	18	18	687	867	577.6	.	0	693.9	.	0
..Falls and fall related injuries (E880-E888)	1	.	1	0	8	.	.	.	.	.	.
..Drowning (E830,E832,E910)	2	2	2	90	110	.	.	.	.	.	.
..Residential fires (E890-E899)	.	.	.	.	.	.	.	.	.	.	.
Suicide (E950-E959)	22	20	21	595	798	619.7	.	0	812.9	.	0
Homicide and legal intervention (E960-E978)	27	26	26	1,089	1,349	833.6	.	0	971.9	.	0
..Homicide (E960-E969)	27	26	26	1,089	1,349	833.6	.	0	971.9	.	0
All injury (E800-E999)	79	75	77	2,729	3,489	2373.5	1657.8	0	2932.6	2121.3	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	17	17	17	655	825	549.3	.	0	659.5	.	0
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	2	23	40	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	32	30	31	1,145	1,448	928.9	.	1	1128.9	.	1
...Suicide (E955.0-.4)	9	8	9	250	333	.	.	.	.	.	.
...Homicide (E965.0-.4)	21	20	20	815	1,015	626.1	.	0	733.6	.	0
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	5	4	4	115	155	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	14	10	14	175	300	.	349.4	.	.	650.2	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	3	2	2	50	70	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).

CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
All causes	218	114	152	3,078	4,433	2928.5	2163.0	0	4574.4	3597.2	0
Infectious diseases and parasitic disease (1-139)	16	13	14	388	525	365.1	.	0	485.6	.	0
..Septicemia (038)	5	3	3	68	98	.	.	.	.	.	.
...Septicemia-related (038)	14	6	9	125	203	.	.	.	.	.	.
..HIV infection (042-044)	10	10	10	320	420	.	.	.	.	.	.
All cancer (140-208)	34	15	24	248	455	278.3	573.9	0	595.5	973.6	0
..Oral cavity or pharynx (140.0-149.9)	.	.	.	.	.	.	.	.	.	.	.
..Colorectal cancer (153.0-154.3,154.8,159.0)	8	3	3	13	43	.	.	.	.	.	.
...Colon cancer (153)	6	3	3	13	43	.	.	.	.	.	.
...Rectal cancer (154)	1	.	.	0	0	.	.	.	.	.	.
..Pancreatic cancer (157)	1	1	1	3	13	.	.	.	.	.	.
..Lung and other respiratory cancer (160-165)	6	1	5	18	53	.	.	.	.	.	.
...Lung cancer (162.2-.9)	6	1	5	18	53	.	.	.	.	.	.
..Malignant melanoma of skin (172)	1	.	1	0	8	.	.	.	.	.	.
..Female breast cancer (174)	2	2	2	70	90	.	.	.	.	.	.
..Cervical cancer (180)	1	1	1	18	28	.	.	.	.	.	.
..Ovarian cancer (183)	1	.	.	0	0	.	.	.	.	.	.
..Prostate cancer (185)	.	.	.	.	.	.	.	.	.	.	.
..Bladder cancer (188)	.	.	.	.	.	.	.	.	.	.	.
..Brain & other CNS cancer (191-192)	2	2	2	20	40	.	.	.	.	.	.
..Leukemia (204-208)	2	2	2	45	65	.	.	.	.	.	.
Diabetes mellitus (250)	11	5	7	43	108	.	.	.	.	.	.
..Diabetes-related (250)	23	11	17	88	238	153.9	178.1	0	420.7	487.5	0
Alzheimer's diseases (331.0)	.	.	.	.	.	.	.	.	.	.	.
Cardiovascular disease (390-459)	76	26	43	397	740	548.7	465.0	0	1103.4	1000.2	0
..Diseases of the heart (390-398,402,404-429)	55	18	31	225	468	329.7	281.3	0	739.2	692.0	0
...Heart disease-related (390-398,402,404-429)	102	39	56	712	1,189	852.4	755.1	0	1553.6	1539.0	0
...Coronary heart disease (CHD) (402,410-414,429.2)	32	9	18	78	220	150.5	140.0	0	415.3	397.8	0
...Congestive heart failure (CHF) (428.0)	3	.	.	0	0	.	.	.	.	.	.
..Hypertension with or without renal disease (401,403)	.	.	.	.	.	.	.	.	.	.	.
...Hypertension-related (401,403)	16	8	12	85	190	148.5	.	0	328.9	.	0
..Cerebrovascular disease (430-438)	16	7	9	155	235	189.9	137.4	0	299.9	221.2	0
..Atherosclerosis (440)	1	1	1	18	28	.	.	.	.	.	.
...Atherosclerosis-related (440)	6	2	2	20	40	.	.	.	.	.	.
..Aortic aneurysm (441)	3	.	1	0	3	.	.	.	.	.	.
Pneumonia & influenza (480-487)	8	.	.	0	0	.	.	.	.	.	.
Chronic obstructive pulmonary disease (COPD) (490-496)	4	2	2	40	60	.	.	.	.	.	.
..COPD-related (490-496)	13	4	7	60	113	.	138.3	.	.	246.2	.
..Asthma (493)	2	2	2	40	60	.	.	.	.	.	.
Chronic liver disease and cirrhosis (571)	5	3	4	63	95	.	.	.	.	.	.

(CONTINUED)

\* - Age-adjusted YPLL are calculated per 100,000 population.  
 . - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).



CONNECTICUT Hispanic Resident DEATHS, 1990  
 Age-adjusted years of potential life lost (YPLL) under 65 and 75 (\*)  
 for selected causes of death by sex of decedent  
 \*\*\* Age-adjusted using 2000 U.S. standard population \*\*\*

	Female										
	Number of Deaths, 1990			Years of life lost, 1990		Adjusted YPLL < 65			Adjusted YPLL < 75		
	All	< 65	< 75	< 65	< 75	1990	1989	Signif- icance	1990	1989	Signif- icance
*****Causes of Death*****											
Nephritis, nephrotic syndrome/nephrosis (580-589)	5	3	5	53	93	.	.	.	.	.	.
..Nephritis-related (580-589)	24	15	20	220	402	267.7	.	0	556.4	.	0
..End stage renal disease (ESRD) (581-588,591,593.8,593.9)	5	3	5	53	93	.	.	.	.	.	.
Unintentional injuries (E800-E949)	7	6	6	210	270	.	.	.	.	.	.
..Motor vehicle accidents (E810-E825)	3	3	3	148	178	.	.	.	.	.	.
..Falls and fall related injuries (E880-E888)	2	1	1	3	13	.	.	.	.	.	.
..Drowning (E830,E832,E910)	.	.	.	.	.	.	.	.	.	.	.
..Residential fires (E890-E899)	1	1	1	18	28	.	.	.	.	.	.
Suicide (E950-E959)	4	4	4	155	195	.	.	.	.	.	.
Homicide and legal intervention (E960-E978)	5	5	5	168	218	.	.	.	.	.	.
..Homicide (E960-E969)	5	5	5	168	218	.	.	.	.	.	.
All injury (E800-E999)	16	15	15	533	683	426.4	.	0	523.7	.	0
..All motor vehicle accidents (E810-E819,E958.5,E988.5)	3	3	3	148	178	.	.	.	.	.	.
..All falls (E880-E886,E888,E957,E968.1,E987)	2	1	1	3	13	.	.	.	.	.	.
..All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)	8	8	8	290	370	.	.	.	.	.	.
...Suicide (E955.0-.4)	3	3	3	123	153	.	.	.	.	.	.
...Homicide (E965.0-.4)	5	5	5	168	218	.	.	.	.	.	.
..All poisoning (E850-E869,E950-E952,E962,E972,E980-E982)	1	1	1	33	43	.	.	.	.	.	.
Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)	3	3	3	93	123	.	.	.	.	.	.
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)	.	.	.	.	.	.	.	.	.	.	.

\* - Age-adjusted YPLL are calculated per 100,000 population.

. - Age-adjusted YPLL under 65 and 75 are not reported due to a small number of deaths (<15).