

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

## Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	4	2.0	0.99	2	1.9	1.37	2	2.0	1.45
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	0	0.0	0.00	1	1.2	1.19
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.0	0.71	1	1.0	0.99	1	1.0	1.03
25-29	2	0.9	0.62	1	0.9	0.87	1	0.9	0.88
30-34	3	1.1	0.66	3	2.3	1.32	0	0.0	0.00
35-39	3	1.2	0.70	2	1.6	1.14	1	0.8	0.80
40-44	2	0.9	0.64	2	1.8	1.30	0	0.0	0.00
45-49	6	3.0	1.24	4	4.1	2.06	2	2.0	1.41
50-54	4	2.7	1.36	4	5.5	2.77	0	0.0	0.00
55-59	9	7.1	2.35	4	6.5	3.23	5	7.6	3.41
60-64	20	15.2	3.41	13	20.8	5.76	7	10.2	3.86
65-69	16	12.6	3.15	5	8.8	3.92	11	15.8	4.75
70-74	46	40.4	5.95	23	47.3	9.86	23	35.2	7.34
75-79	60	67.2	8.67	33	93.0	16.19	27	50.1	9.65
80-84	59	103.9	13.53	21	107.4	23.44	38	102.1	16.56
85+	106	220.6	21.43	26	211.5	41.48	80	223.7	25.01
Total deaths	343			144			199		
Connecticut									
. Crude rate		11.7	-		10.2	-		13.2	-
. AAMR (1940 Std. mil.)		5.0	0.32		5.8	0.52		4.3	0.40
. AAMR (1970 Std. mil.)		7.5	0.42		8.5	0.71		6.6	0.52
. AAMR (2000 Std. mil.)		10.7	0.58		12.0	1.03		9.6	0.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.9	-		4.5	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is higher  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.0	0.70	1	1.0	0.97	1	1.0	1.02
5-9	2	1.1	0.76	2	2.1	1.49	0	0.0	0.00
10-14	5	2.9	1.30	3	3.4	1.97	2	2.4	1.68
15-19	7	4.2	1.59	4	4.7	2.34	3	3.7	2.15
20-24	10	5.0	1.60	8	7.9	2.80	2	2.1	1.45
25-29	10	4.4	1.39	5	4.4	1.95	5	4.4	1.97
30-34	33	12.6	2.20	20	15.3	3.42	13	9.9	2.76
35-39	83	33.4	3.66	35	28.3	4.78	48	38.4	5.54
40-44	118	53.1	4.89	60	55.0	7.10	58	51.3	6.74
45-49	179	90.7	6.78	91	93.6	9.81	88	87.8	9.36
50-54	255	172.8	10.82	121	167.7	15.24	134	177.8	15.36
55-59	398	312.1	15.64	214	345.4	23.61	184	280.6	20.69
60-64	613	467.4	18.88	336	536.8	29.28	277	404.0	24.27
65-69	963	758.9	24.46	542	949.6	40.79	421	603.0	29.39
70-74	1,081	948.7	28.86	563	1157.6	48.79	518	793.2	34.85
75-79	1,103	1234.4	37.17	596	1679.9	68.81	507	941.1	41.80
80-84	914	1609.7	53.24	464	2373.8	110.20	450	1208.6	56.97
85+	874	1818.9	61.52	347	2822.5	151.52	527	1473.8	64.20
Total deaths	6,650			3,412			3,238		
Connecticut									
. Crude rate		227.6	-		240.7	-		215.4	-
. AAMR (1940 Std. mil.)		123.3	1.70		149.5	2.72		105.2	2.18
. AAMR (1970 Std. mil.)		163.7	2.08		205.2	3.54		137.0	2.58
. AAMR (2000 Std. mil.)		206.3	2.55		265.1	4.68		171.4	3.07
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		133.1	-		162.7	-		111.7	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.81	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	7	3.5	1.34	5	5.1	2.30	2	2.0	1.41
50-54	7	4.7	1.79	7	9.7	3.67	0	0.0	0.00
55-59	10	7.8	2.48	7	11.3	4.27	3	4.6	2.64
60-64	3	2.3	1.32	3	4.8	2.77	0	0.0	0.00
65-69	17	13.4	3.25	6	10.5	4.29	11	15.8	4.75
70-74	10	8.8	2.78	5	10.3	4.60	5	7.7	3.42
75-79	17	19.0	4.61	6	16.9	6.90	11	20.4	6.16
80-84	10	17.6	5.57	6	30.7	12.53	4	10.7	5.37
85+	13	27.1	7.50	5	40.7	18.19	8	22.4	7.91
Total deaths	95			51			44		
Connecticut									
. Crude rate		3.3	-		3.6	-		2.9	-
. AAMR (1940 Std. mil.)		1.9	0.22		2.6	0.39		1.3	0.23
. AAMR (1970 Std. mil.)		2.4	0.26		3.2	0.46		1.8	0.28
. AAMR (2000 Std. mil.)		3.0	0.31		4.1	0.59		2.2	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.5	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Colon cancer (153)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	4	1.6	0.80	2	1.6	1.14	2	1.6	1.13
40-44	8	3.6	1.27	4	3.7	1.83	4	3.5	1.77
45-49	9	4.6	1.52	6	6.2	2.52	3	3.0	1.73
50-54	13	8.8	2.44	7	9.7	3.67	6	8.0	3.25
55-59	29	22.7	4.22	19	30.7	7.04	10	15.2	4.82
60-64	46	35.1	5.17	26	41.5	8.15	20	29.2	6.52
65-69	77	60.7	6.92	46	80.6	11.88	31	44.4	7.98
70-74	86	75.5	8.14	44	90.5	13.64	42	64.3	9.92
75-79	112	125.3	11.84	62	174.8	22.19	50	92.8	13.13
80-84	107	188.4	18.22	55	281.4	37.94	52	139.7	19.37
85+	123	256.0	23.08	43	349.8	53.34	80	223.7	25.01
Total deaths	616			316			300		
Connecticut									
. Crude rate		21.1	-		22.3	-		20.0	-
. AAMR (1940 Std. mil.)		9.9	0.46		13.0	0.78		7.7	0.54
. AAMR (1970 Std. mil.)		14.2	0.59		18.8	1.07		11.1	0.69
. AAMR (2000 Std. mil.)		19.0	0.77		25.5	1.48		15.0	0.88
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.0	-		13.5	-		9.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.5	0.45	1	0.9	0.92	0	0.0	0.00
45-49	7	3.5	1.34	5	5.1	2.30	2	2.0	1.41
50-54	6	4.1	1.66	2	2.8	1.96	4	5.3	2.65
55-59	6	4.7	1.92	4	6.5	3.23	2	3.0	2.16
60-64	10	7.6	2.41	7	11.2	4.23	3	4.4	2.53
65-69	13	10.2	2.84	10	17.5	5.54	3	4.3	2.48
70-74	19	16.7	3.83	10	20.6	6.50	9	13.8	4.59
75-79	28	31.3	5.92	17	47.9	11.62	11	20.4	6.16
80-84	18	31.7	7.47	9	46.0	15.35	9	24.2	8.06
85+	24	49.9	10.20	8	65.1	23.01	16	44.7	11.19
Total deaths	132			73			59		
Connecticut									
. Crude rate		4.5	-		5.1	-		3.9	-
. AAMR (1940 Std. mil.)		2.3	0.23		3.1	0.39		1.6	0.26
. AAMR (1970 Std. mil.)		3.1	0.28		4.4	0.51		2.3	0.32
. AAMR (2000 Std. mil.)		4.1	0.36		5.7	0.69		3.1	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.6	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.38	0	0.0	0.00	1	0.8	0.76
35-39	3	1.2	0.70	1	0.8	0.81	2	1.6	1.13
40-44	4	1.8	0.90	2	1.8	1.30	2	1.8	1.25
45-49	7	3.5	1.34	7	7.2	2.72	0	0.0	0.00
50-54	10	6.8	2.14	6	8.3	3.39	4	5.3	2.65
55-59	20	15.7	3.51	10	16.1	5.10	10	15.2	4.82
60-64	31	23.6	4.24	15	24.0	6.19	16	23.3	5.83
65-69	52	41.0	5.68	24	42.0	8.58	28	40.1	7.58
70-74	61	53.5	6.85	32	65.8	11.63	29	44.4	8.25
75-79	56	62.7	8.38	24	67.6	13.81	32	59.4	10.50
80-84	52	91.6	12.70	15	76.7	19.81	37	99.4	16.34
85+	61	126.9	16.25	9	73.2	24.40	52	145.4	20.17
Total deaths	358			145			213		
Connecticut									
. Crude rate		12.3	-		10.2	-		14.2	-
. AAMR (1940 Std. mil.)		6.2	0.37		6.6	0.58		5.8	0.48
. AAMR (1970 Std. mil.)		8.5	0.47		8.7	0.73		8.2	0.60
. AAMR (2000 Std. mil.)		11.0	0.59		10.8	0.92		10.7	0.74
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		7.6	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung and other respiratory cancer (160-165)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.44	0	0.0	0.00	1	0.9	0.88
30-34	3	1.1	0.66	3	2.3	1.32	0	0.0	0.00
35-39	7	2.8	1.06	4	3.2	1.61	3	2.4	1.39
40-44	23	10.4	2.16	15	13.7	3.55	8	7.1	2.50
45-49	50	25.3	3.58	24	24.7	5.04	26	25.9	5.09
50-54	75	50.8	5.87	35	48.5	8.20	40	53.1	8.39
55-59	136	106.6	9.14	85	137.2	14.88	51	77.8	10.89
60-64	216	164.7	11.21	124	198.1	17.79	92	134.2	13.99
65-69	327	257.7	14.25	207	362.7	25.21	120	171.9	15.69
70-74	338	296.6	16.14	194	398.9	28.64	144	220.5	18.38
75-79	311	348.1	19.74	181	510.2	37.92	130	241.3	21.16
80-84	203	357.5	25.09	127	649.7	57.65	76	204.1	23.41
85+	104	216.4	21.22	64	520.6	65.07	40	111.9	17.69
Total deaths	1,795			1,064			731		
Connecticut									
. Crude rate		61.4	-		75.1	-		48.6	-
. AAMR (1940 Std. mil.)		36.0	0.93		48.1	1.55		27.1	1.12
. AAMR (1970 Std. mil.)		46.4	1.13		64.5	1.99		34.1	1.32
. AAMR (2000 Std. mil.)		55.2	1.31		79.3	2.49		39.8	1.50
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		40.8	-		58.5	-		27.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.44	0	0.0	0.00	1	0.9	0.88
30-34	3	1.1	0.66	3	2.3	1.32	0	0.0	0.00
35-39	7	2.8	1.06	4	3.2	1.61	3	2.4	1.39
40-44	22	9.9	2.11	14	12.8	3.43	8	7.1	2.50
45-49	48	24.3	3.51	22	22.6	4.83	26	25.9	5.09
50-54	69	46.8	5.63	29	40.2	7.46	40	53.1	8.39
55-59	132	103.5	9.01	82	132.4	14.62	50	76.2	10.78
60-64	211	160.9	11.07	121	193.3	17.57	90	131.3	13.84
65-69	316	249.0	14.01	199	348.7	24.72	117	167.6	15.49
70-74	331	290.5	15.97	188	386.6	28.19	143	219.0	18.31
75-79	298	333.5	19.32	171	482.0	36.86	127	235.7	20.92
80-84	195	343.4	24.59	120	613.9	56.04	75	201.4	23.26
85+	103	214.4	21.12	63	512.4	64.56	40	111.9	17.69
Total deaths	1,736			1,016			720		
Connecticut									
. Crude rate		59.4	-		71.7	-		47.9	-
. AAMR (1940 Std. mil.)		34.7	0.91		45.8	1.51		26.7	1.11
. AAMR (1970 Std. mil.)		44.8	1.11		61.5	1.94		33.5	1.31
. AAMR (2000 Std. mil.)		53.3	1.29		75.7	2.43		39.2	1.49
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		39.3	-		56.0	-		26.3	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	1.5	0.87	2	2.0	1.40	1	1.0	1.03
25-29	1	0.4	0.44	0	0.0	0.00	1	0.9	0.88
30-34	2	0.8	0.54	1	0.8	0.76	1	0.8	0.76
35-39	8	3.2	1.14	5	4.0	1.81	3	2.4	1.39
40-44	4	1.8	0.90	2	1.8	1.30	2	1.8	1.25
45-49	2	1.0	0.72	2	2.1	1.45	0	0.0	0.00
50-54	4	2.7	1.36	1	1.4	1.39	3	4.0	2.30
55-59	9	7.1	2.35	5	8.1	3.61	4	6.1	3.05
60-64	8	6.1	2.16	2	3.2	2.26	6	8.8	3.57
65-69	9	7.1	2.36	4	7.0	3.50	5	7.2	3.20
70-74	20	17.6	3.92	14	28.8	7.69	6	9.2	3.75
75-79	8	9.0	3.17	6	16.9	6.90	2	3.7	2.63
80-84	11	19.4	5.84	6	30.7	12.53	5	13.4	6.01
85+	9	18.7	6.24	3	24.4	14.09	6	16.8	6.85
Total deaths	98			53			45		
Connecticut									
. Crude rate		3.4	-		3.7	-		3.0	-
. AAMR (1940 Std. mil.)		2.1	0.24		2.5	0.37		1.9	0.32
. AAMR (1970 Std. mil.)		2.5	0.26		3.1	0.43		2.1	0.34
. AAMR (2000 Std. mil.)		3.1	0.31		3.9	0.55		2.5	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.6	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	1	0.4	0.44	-	-	-	1	0.9	0.88
30-34	3	1.1	0.66	-	-	-	3	2.3	1.32
35-39	20	8.0	1.80	-	-	-	20	16.0	3.58
40-44	22	9.9	2.11	-	-	-	22	19.5	4.15
45-49	21	10.6	2.32	-	-	-	21	21.0	4.57
50-54	33	22.4	3.89	-	-	-	33	43.8	7.62
55-59	46	36.1	5.32	-	-	-	46	70.1	10.34
60-64	53	40.4	5.55	-	-	-	53	77.3	10.62
65-69	66	52.0	6.40	-	-	-	66	94.5	11.64
70-74	81	71.1	7.90	-	-	-	81	124.0	13.78
75-79	65	72.7	9.02	-	-	-	65	120.7	14.97
80-84	56	98.6	13.18	-	-	-	56	150.4	20.10
85+	71	147.8	17.54	-	-	-	71	198.6	23.56
Total deaths	538			-			538		
Connecticut									
. Crude rate		-	-		-	-		35.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		20.3	1.00
. AAMR (1970 Std. mil.)		-	-		-	-		24.6	1.13
. AAMR (2000 Std. mil.)		-	-		-	-		30.0	1.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		21.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	1	0.4	0.38	-	-	-	1	0.8	0.76
35-39	3	1.2	0.70	-	-	-	3	2.4	1.39
40-44	1	0.5	0.45	-	-	-	1	0.9	0.88
45-49	4	2.0	1.01	-	-	-	4	4.0	2.00
50-54	6	4.1	1.66	-	-	-	6	8.0	3.25
55-59	0	0.0	0.00	-	-	-	0	0.0	0.00
60-64	0	0.0	0.00	-	-	-	0	0.0	0.00
65-69	4	3.2	1.58	-	-	-	4	5.7	2.86
70-74	5	4.4	1.96	-	-	-	5	7.7	3.42
75-79	4	4.5	2.24	-	-	-	4	7.4	3.71
80-84	1	1.8	1.76	-	-	-	1	2.7	2.69
85+	3	6.2	3.60	-	-	-	3	8.4	4.84
Total deaths	32			-			32		
Connecticut									
. Crude rate		-	-		-	-		2.1	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.4	0.28
. AAMR (1970 Std. mil.)		-	-		-	-		1.5	0.29
. AAMR (2000 Std. mil.)		-	-		-	-		1.9	0.35
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	0	0.0	0.00	-	-	-	0	0.0	0.00
50-54	0	0.0	0.00	-	-	-	0	0.0	0.00
55-59	3	2.4	1.36	-	-	-	3	4.6	2.64
60-64	4	3.0	1.52	-	-	-	4	5.8	2.92
65-69	7	5.5	2.09	-	-	-	7	10.0	3.79
70-74	6	5.3	2.15	-	-	-	6	9.2	3.75
75-79	3	3.4	1.94	-	-	-	3	5.6	3.22
80-84	4	7.0	3.52	-	-	-	4	10.7	5.37
85+	6	12.5	5.10	-	-	-	6	16.8	6.85
Total deaths	33			-			33		
Connecticut									
. Crude rate		-	-		-	-		2.2	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.1	0.21
. AAMR (1970 Std. mil.)		-	-		-	-		1.4	0.26
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	4	1.8	0.90	-	-	-	4	3.5	1.77
45-49	5	2.5	1.13	-	-	-	5	5.0	2.23
50-54	8	5.4	1.92	-	-	-	8	10.6	3.75
55-59	11	8.6	2.60	-	-	-	11	16.8	5.06
60-64	21	16.0	3.49	-	-	-	21	30.6	6.68
65-69	23	18.1	3.78	-	-	-	23	32.9	6.87
70-74	29	25.5	4.73	-	-	-	29	44.4	8.25
75-79	26	29.1	5.71	-	-	-	26	48.3	9.46
80-84	23	40.5	8.45	-	-	-	23	61.8	12.88
85+	23	47.9	9.98	-	-	-	23	64.3	13.41
Total deaths	173			-			173		
Connecticut									
. Crude rate		-	-		-	-		11.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	0.52
. AAMR (1970 Std. mil.)		-	-		-	-		7.6	0.62
. AAMR (2000 Std. mil.)		-	-		-	-		9.3	0.72
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Prostate cancer (185)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	-	-	-
5-9	0	0.0	0.00	0	0.0	0.00	-	-	-
10-14	0	0.0	0.00	0	0.0	0.00	-	-	-
15-19	0	0.0	0.00	0	0.0	0.00	-	-	-
20-24	0	0.0	0.00	0	0.0	0.00	-	-	-
25-29	0	0.0	0.00	0	0.0	0.00	-	-	-
30-34	0	0.0	0.00	0	0.0	0.00	-	-	-
35-39	0	0.0	0.00	0	0.0	0.00	-	-	-
40-44	0	0.0	0.00	0	0.0	0.00	-	-	-
45-49	2	1.0	0.72	2	2.1	1.45	-	-	-
50-54	3	2.0	1.17	3	4.2	2.40	-	-	-
55-59	7	5.5	2.07	7	11.3	4.27	-	-	-
60-64	26	19.8	3.89	26	41.5	8.15	-	-	-
65-69	46	36.3	5.35	46	80.6	11.88	-	-	-
70-74	71	62.3	7.40	71	146.0	17.33	-	-	-
75-79	102	114.2	11.30	102	287.5	28.47	-	-	-
80-84	98	172.6	17.43	98	501.4	50.64	-	-	-
85+	93	193.5	20.07	93	756.5	78.44	-	-	-
Total deaths	448			448			-		
Connecticut									
. Crude rate		-	-		31.6	-		-	-
. AAMR (1940 Std. mil.)		-	-		15.8	0.79		-	-
. AAMR (1970 Std. mil.)		-	-		26.0	1.23		-	-
. AAMR (2000 Std. mil.)		-	-		38.4	1.88		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		16.6	-		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.81	0	0.0	0.00
40-44	1	0.5	0.45	0	0.0	0.00	1	0.9	0.88
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.68	0	0.0	0.00	1	1.3	1.33
55-59	4	3.1	1.57	4	6.5	3.23	0	0.0	0.00
60-64	11	8.4	2.53	10	16.0	5.05	1	1.5	1.46
65-69	18	14.2	3.34	13	22.8	6.32	5	7.2	3.20
70-74	19	16.7	3.83	14	28.8	7.69	5	7.7	3.42
75-79	33	36.9	6.43	24	67.6	13.81	9	16.7	5.57
80-84	44	77.5	11.68	24	122.8	25.06	20	53.7	12.01
85+	38	79.1	12.83	16	130.1	32.54	22	61.5	13.12
Total deaths	170			106			64		
Connecticut									
. Crude rate		5.8	-		7.5	-		4.3	-
. AAMR (1940 Std. mil.)		2.4	0.21		4.0	0.41		1.2	0.19
. AAMR (1970 Std. mil.)		3.7	0.29		6.2	0.61		2.0	0.27
. AAMR (2000 Std. mil.)		5.2	0.40		8.7	0.88		3.1	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.1	-		3.6	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	0	0.0	0.00	1	1.2	1.19
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.50	1	1.0	0.99	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.1	0.66	2	1.5	1.08	1	0.8	0.76
35-39	4	1.6	0.80	2	1.6	1.14	2	1.6	1.13
40-44	9	4.1	1.35	8	7.3	2.59	1	0.9	0.88
45-49	9	4.6	1.52	6	6.2	2.52	3	3.0	1.73
50-54	17	11.5	2.79	12	16.6	4.80	5	6.6	2.97
55-59	15	11.8	3.04	12	19.4	5.59	3	4.6	2.64
60-64	18	13.7	3.23	12	19.2	5.53	6	8.8	3.57
65-69	23	18.1	3.78	12	21.0	6.07	11	15.8	4.75
70-74	26	22.8	4.48	13	26.7	7.41	13	19.9	5.52
75-79	12	13.4	3.88	5	14.1	6.30	7	13.0	4.91
80-84	13	22.9	6.35	6	30.7	12.53	7	18.8	7.11
85+	13	27.1	7.50	3	24.4	14.09	10	28.0	8.84
Total deaths	164			94			70		
Connecticut									
. Crude rate		5.6	-		6.6	-		4.7	-
. AAMR (1940 Std. mil.)		3.8	0.33		5.2	0.56		2.6	0.36
. AAMR (1970 Std. mil.)		4.5	0.36		6.0	0.62		3.2	0.41
. AAMR (2000 Std. mil.)		5.3	0.42		7.0	0.73		3.8	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.7	-		4.2	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.54	1	1.1	1.05	0	0.0	0.00
10-14	2	1.2	0.82	1	1.1	1.14	1	1.2	1.19
15-19	1	0.6	0.60	1	1.2	1.17	0	0.0	0.00
20-24	1	0.5	0.50	1	1.0	0.99	0	0.0	0.00
25-29	2	0.9	0.62	1	0.9	0.87	1	0.9	0.88
30-34	5	1.9	0.85	3	2.3	1.32	2	1.5	1.08
35-39	6	2.4	0.98	5	4.0	1.81	1	0.8	0.80
40-44	7	3.2	1.19	6	5.5	2.24	1	0.9	0.88
45-49	12	6.1	1.75	6	6.2	2.52	6	6.0	2.44
50-54	7	4.7	1.79	2	2.8	1.96	5	6.6	2.97
55-59	14	11.0	2.93	8	12.9	4.57	6	9.1	3.74
60-64	13	9.9	2.75	8	12.8	4.52	5	7.3	3.26
65-69	24	18.9	3.86	12	21.0	6.07	12	17.2	4.96
70-74	33	29.0	5.04	18	37.0	8.72	15	23.0	5.93
75-79	39	43.6	6.99	19	53.6	12.29	20	37.1	8.30
80-84	32	56.4	9.96	15	76.7	19.81	17	45.7	11.07
85+	39	81.2	13.00	17	138.3	33.54	22	61.5	13.12
Total deaths	238			124			114		
Connecticut									
. Crude rate		8.1	-		8.7	-		7.6	-
. AAMR (1940 Std. mil.)		4.5	0.34		5.6	0.55		3.6	0.41
. AAMR (1970 Std. mil.)		5.8	0.40		7.5	0.68		4.7	0.48
. AAMR (2000 Std. mil.)		7.5	0.49		9.9	0.92		6.1	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.8	-		6.3	-		4.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.54	0	0.0	0.00	1	1.1	1.11
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
25-29	3	1.3	0.76	1	0.9	0.87	2	1.8	1.25
30-34	8	3.1	1.08	4	3.1	1.53	4	3.1	1.53
35-39	11	4.4	1.33	8	6.5	2.28	3	2.4	1.39
40-44	17	7.7	1.86	11	10.1	3.04	6	5.3	2.17
45-49	22	11.1	2.38	14	14.4	3.85	8	8.0	2.82
50-54	43	29.1	4.44	28	38.8	7.33	15	19.9	5.14
55-59	76	59.6	6.84	42	67.8	10.46	34	51.8	8.89
60-64	152	115.9	9.40	85	135.8	14.73	67	97.7	11.94
65-69	223	175.7	11.77	120	210.2	19.19	103	147.5	14.54
70-74	328	287.9	15.90	178	366.0	27.43	150	229.7	18.75
75-79	409	457.7	22.63	203	572.2	40.16	206	382.4	26.64
80-84	342	602.3	32.57	147	752.0	62.03	195	523.7	37.51
85+	488	1015.6	45.97	129	1049.3	92.39	359	1004.0	52.99
Total deaths	2,124			970			1,154		
Connecticut									
. Crude rate		72.7	-		68.4	-		76.8	-
. AAMR (1940 Std. mil.)		32.6	0.81		39.4	1.34		27.4	1.00
. AAMR (1970 Std. mil.)		47.9	1.08		57.3	1.85		41.0	1.30
. AAMR (2000 Std. mil.)		65.4	1.43		77.6	2.57		56.8	1.69
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): No comparison is available  
. THE U.S. HP 2000 target: CT rate is not significantly different



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	1	0.8	0.78	0	0.0	0.00	1	1.5	1.52
60-64	4	3.0	1.52	3	4.8	2.77	1	1.5	1.46
65-69	12	9.5	2.73	8	14.0	4.96	4	5.7	2.86
70-74	13	11.4	3.16	7	14.4	5.44	6	9.2	3.75
75-79	25	28.0	5.60	15	42.3	10.92	10	18.6	5.87
80-84	45	79.3	11.81	19	97.2	22.30	26	69.8	13.69
85+	81	168.6	18.73	22	178.9	38.15	59	165.0	21.48
Total deaths	181			74			107		
Connecticut									
. Crude rate		6.2	-		5.2	-		7.1	-
. AAMR (1940 Std. mil.)		1.9	0.16		2.4	0.29		1.5	0.18
. AAMR (1970 Std. mil.)		3.5	0.26		4.3	0.50		2.9	0.30
. AAMR (2000 Std. mil.)		5.6	0.42		6.8	0.82		4.9	0.48
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.2	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	6.5	1.79	7	6.8	2.56	6	6.1	2.50
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	0	0.0	0.00	1	1.2	1.19
15-19	1	0.6	0.60	1	1.2	1.17	0	0.0	0.00
20-24	8	4.0	1.43	2	2.0	1.40	6	6.2	2.52
25-29	15	6.6	1.70	9	7.9	2.62	6	5.3	2.16
30-34	25	9.6	1.91	19	14.5	3.33	6	4.6	1.87
35-39	48	19.3	2.78	36	29.1	4.84	12	9.6	2.77
40-44	90	40.5	4.27	70	64.1	7.66	20	17.7	3.96
45-49	157	79.5	6.35	110	113.2	10.79	47	46.9	6.84
50-54	212	143.7	9.87	150	207.9	16.97	62	82.3	10.45
55-59	326	255.6	14.16	234	377.7	24.69	92	140.3	14.63
60-64	543	414.0	17.77	362	578.3	30.39	181	264.0	19.62
65-69	913	719.5	23.81	576	1009.2	42.05	337	482.7	26.29
70-74	1,346	1181.3	32.20	820	1686.0	58.88	526	805.5	35.12
75-79	1,717	1921.6	46.37	880	2480.4	83.61	837	1553.6	53.70
80-84	1,969	3467.8	78.15	848	4338.3	148.98	1,121	3010.8	89.92
85+	4,088	8507.5	133.06	1,149	9346.0	275.72	2,939	8219.1	151.61
Total deaths	11,472			5,273			6,199		
Connecticut									
. Crude rate		392.7	-		371.9	-		412.3	-
. AAMR (1940 Std. mil.)		155.1	1.72		206.6	3.06		114.7	1.89
. AAMR (1970 Std. mil.)		243.0	2.36		313.7	4.35		191.1	2.63
. AAMR (2000 Std. mil.)		357.8	3.35		449.2	6.42		294.7	3.78
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		181.3	-		237.6	-		136.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is higher  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Diseases of the heart (390-398,402,404-429)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	12	6.0	1.72	6	5.8	2.37	6	6.1	2.50
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	0	0.0	0.00	1	1.2	1.19
15-19	1	0.6	0.60	1	1.2	1.17	0	0.0	0.00
20-24	6	3.0	1.24	1	1.0	0.99	5	5.1	2.30
25-29	12	5.3	1.52	9	7.9	2.62	3	2.6	1.53
30-34	21	8.0	1.75	18	13.8	3.24	3	2.3	1.32
35-39	39	15.7	2.51	30	24.2	4.42	9	7.2	2.40
40-44	78	35.1	3.97	61	55.9	7.16	17	15.0	3.65
45-49	126	63.8	5.69	96	98.8	10.08	30	29.9	5.47
50-54	180	122.0	9.09	133	184.3	15.98	47	62.4	9.10
55-59	284	222.7	13.21	209	337.4	23.34	75	114.4	13.21
60-64	476	362.9	16.63	326	520.8	28.84	150	218.8	17.86
65-69	788	621.0	22.12	504	883.0	39.33	284	406.8	24.14
70-74	1,098	963.7	29.08	676	1389.9	53.46	422	646.2	31.46
75-79	1,366	1528.8	41.36	704	1984.3	74.79	662	1228.8	47.76
80-84	1,501	2643.5	68.23	663	3391.8	131.73	838	2250.7	77.75
85+	3,137	6528.3	116.56	896	7288.1	243.48	2,241	6267.1	132.39
Total deaths	9,126			4,333			4,793		
Connecticut									
. Crude rate		312.4	-		305.6	-		318.8	-
. AAMR (1940 Std. mil.)		126.7	1.57		173.1	2.82		90.1	1.68
. AAMR (1970 Std. mil.)		195.7	2.13		258.6	3.95		149.1	2.34
. AAMR (2000 Std. mil.)		284.5	2.99		365.3	5.76		228.1	3.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		144.3	-		195.2	-		103.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available









## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	3	1.5	0.88	2	2.1	1.45	1	1.0	1.00
50-54	3	2.0	1.17	2	2.8	1.96	1	1.3	1.33
55-59	2	1.6	1.11	2	3.2	2.28	0	0.0	0.00
60-64	8	6.1	2.16	5	8.0	3.57	3	4.4	2.53
65-69	11	8.7	2.61	6	10.5	4.29	5	7.2	3.20
70-74	8	7.0	2.48	4	8.2	4.11	4	6.1	3.06
75-79	19	21.3	4.88	8	22.5	7.97	11	20.4	6.16
80-84	26	45.8	8.98	8	40.9	14.47	18	48.3	11.39
85+	36	74.9	12.49	13	105.7	29.33	23	64.3	13.41
Total deaths	116			50			66		
Connecticut									
. Crude rate		4.0	-		3.5	-		4.4	-
. AAMR (1940 Std. mil.)		1.6	0.18		2.0	0.30		1.3	0.20
. AAMR (1970 Std. mil.)		2.5	0.24		3.0	0.43		2.1	0.28
. AAMR (2000 Std. mil.)		3.6	0.34		4.4	0.65		3.2	0.39
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.2	-		1.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.50	1	1.0	0.99	0	0.0	0.00
25-29	1	0.4	0.44	0	0.0	0.00	1	0.9	0.88
30-34	3	1.1	0.66	1	0.8	0.76	2	1.5	1.08
35-39	6	2.4	0.98	4	3.2	1.61	2	1.6	1.13
40-44	10	4.5	1.42	8	7.3	2.59	2	1.8	1.25
45-49	21	10.6	2.32	6	6.2	2.52	15	15.0	3.86
50-54	25	16.9	3.39	11	15.2	4.60	14	18.6	4.96
55-59	26	20.4	4.00	16	25.8	6.46	10	15.2	4.82
60-64	41	31.3	4.88	20	31.9	7.14	21	30.6	6.68
65-69	75	59.1	6.83	39	68.3	10.94	36	51.6	8.59
70-74	171	150.1	11.48	95	195.3	20.04	76	116.4	13.35
75-79	249	278.7	17.66	120	338.2	30.88	129	239.4	21.08
80-84	336	591.8	32.28	129	659.9	58.11	207	556.0	38.64
85+	688	1431.8	54.59	173	1407.2	106.99	515	1440.2	63.46
Total deaths	1,653			623			1,030		
Connecticut									
. Crude rate		56.6	-		43.9	-		68.5	-
. AAMR (1940 Std. mil.)		19.5	0.57		21.8	0.93		17.9	0.73
. AAMR (1970 Std. mil.)		33.0	0.84		36.4	1.46		30.7	1.03
. AAMR (2000 Std. mil.)		51.8	1.28		56.2	2.33		48.8	1.54
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.2	-		28.6	-		24.1	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Atherosclerosis (440)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.5	0.45	1	0.9	0.92	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	2	1.6	1.11	0	0.0	0.00	2	3.0	2.16
60-64	3	2.3	1.32	2	3.2	2.26	1	1.5	1.46
65-69	7	5.5	2.09	5	8.8	3.92	2	2.9	2.03
70-74	11	9.7	2.91	5	10.3	4.60	6	9.2	3.75
75-79	21	23.5	5.13	12	33.8	9.76	9	16.7	5.57
80-84	33	58.1	10.12	13	66.5	18.45	20	53.7	12.01
85+	113	235.2	22.12	26	211.5	41.48	87	243.3	26.08
Total deaths	191			64			127		
Connecticut									
. Crude rate		6.5	-		4.5	-		8.4	-
. AAMR (1940 Std. mil.)		1.8	0.15		2.0	0.26		1.6	0.19
. AAMR (1970 Std. mil.)		3.5	0.26		3.7	0.47		3.3	0.31
. AAMR (2000 Std. mil.)		6.0	0.44		6.2	0.80		5.8	0.52
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		2.8	-		2.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.5	1.11	3	2.9	1.68	2	2.0	1.45
5-9	1	0.5	0.54	1	1.1	1.05	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	1	1.2	1.17	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	5	2.0	0.90	2	1.6	1.14	3	2.4	1.39
40-44	2	0.9	0.64	1	0.9	0.92	1	0.9	0.88
45-49	12	6.1	1.75	10	10.3	3.25	2	2.0	1.41
50-54	9	6.1	2.03	7	9.7	3.67	2	2.7	1.88
55-59	12	9.4	2.72	7	11.3	4.27	5	7.6	3.41
60-64	20	15.2	3.41	8	12.8	4.52	12	17.5	5.05
65-69	46	36.3	5.35	26	45.6	8.93	20	28.6	6.41
70-74	84	73.7	8.04	54	111.0	15.11	30	45.9	8.39
75-79	133	148.8	12.91	75	211.4	24.41	58	107.7	14.14
80-84	211	371.6	25.58	102	521.8	51.67	109	292.8	28.04
85+	553	1150.8	48.94	188	1529.2	111.53	365	1020.8	53.43
Total deaths	1,096			487			609		
Connecticut									
. Crude rate		37.5	-		34.4	-		40.5	-
. AAMR (1940 Std. mil.)		11.8	0.43		16.1	0.79		9.1	0.49
. AAMR (1970 Std. mil.)		21.1	0.66		28.7	1.31		16.9	0.74
. AAMR (2000 Std. mil.)		34.5	1.04		46.9	2.20		28.2	1.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.6	-		16.7	-		10.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.60	0	0.0	0.00	1	1.2	1.24
20-24	1	0.5	0.50	0	0.0	0.00	1	1.0	1.03
25-29	2	0.9	0.62	0	0.0	0.00	2	1.8	1.25
30-34	1	0.4	0.38	1	0.8	0.76	0	0.0	0.00
35-39	1	0.4	0.40	1	0.8	0.81	0	0.0	0.00
40-44	4	1.8	0.90	2	1.8	1.30	2	1.8	1.25
45-49	5	2.5	1.13	2	2.1	1.45	3	3.0	1.73
50-54	5	3.4	1.52	2	2.8	1.96	3	4.0	2.30
55-59	24	18.8	3.84	11	17.8	5.35	13	19.8	5.50
60-64	59	45.0	5.86	34	54.3	9.31	25	36.5	7.29
65-69	129	101.7	8.95	80	140.2	15.67	49	70.2	10.03
70-74	179	157.1	11.74	95	195.3	20.04	84	128.6	14.03
75-79	241	269.7	17.37	134	377.7	32.63	107	198.6	19.20
80-84	204	359.3	25.15	113	578.1	54.38	91	244.4	25.62
85+	267	555.6	34.01	111	902.9	85.70	156	436.3	34.93
Total deaths	1,123			586			537		
Connecticut									
. Crude rate		38.4	-		41.3	-		35.7	-
. AAMR (1940 Std. mil.)		15.9	0.54		21.3	0.92		12.5	0.66
. AAMR (1970 Std. mil.)		24.4	0.75		34.0	1.41		18.9	0.88
. AAMR (2000 Std. mil.)		34.2	1.02		49.1	2.10		26.2	1.14
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		19.9	-		26.4	-		15.4	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	5	2.2	0.98	2	1.7	1.23	3	2.6	1.53
30-34	6	2.3	0.94	5	3.8	1.71	1	0.8	0.76
35-39	20	8.0	1.80	12	9.7	2.80	8	6.4	2.26
40-44	16	7.2	1.80	12	11.0	3.17	4	3.5	1.77
45-49	21	10.6	2.32	17	17.5	4.24	4	4.0	2.00
50-54	21	14.2	3.11	18	24.9	5.88	3	4.0	2.30
55-59	29	22.7	4.22	21	33.9	7.40	8	12.2	4.31
60-64	30	22.9	4.18	17	27.2	6.59	13	19.0	5.26
65-69	37	29.2	4.79	18	31.5	7.43	19	27.2	6.24
70-74	24	21.1	4.30	18	37.0	8.72	6	9.2	3.75
75-79	28	31.3	5.92	14	39.5	10.55	14	26.0	6.95
80-84	20	35.2	7.88	10	51.2	16.18	10	26.9	8.49
85+	6	12.5	5.10	3	24.4	14.09	3	8.4	4.84
Total deaths	263			167			96		
Connecticut									
. Crude rate		9.0	-		11.8	-		6.4	-
. AAMR (1940 Std. mil.)		6.6	0.43		9.3	0.74		4.1	0.46
. AAMR (1970 Std. mil.)		7.3	0.46		10.3	0.81		4.7	0.50
. AAMR (2000 Std. mil.)		8.5	0.53		12.2	0.96		5.4	0.56
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		8.1	-		11.6	-		4.7	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different







## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	12	6.0	1.72	10	9.7	3.06	2	2.0	1.45
5-9	2	1.1	0.76	2	2.1	1.49	0	0.0	0.00
10-14	6	3.5	1.43	6	6.8	2.79	0	0.0	0.00
15-19	41	24.7	3.86	30	35.1	6.40	11	13.7	4.12
20-24	63	31.8	4.01	56	55.5	7.42	7	7.2	2.72
25-29	50	21.9	3.10	39	34.0	5.45	11	9.7	2.93
30-34	67	25.6	3.13	59	45.1	5.87	8	6.1	2.16
35-39	57	22.9	3.03	47	37.9	5.53	10	8.0	2.53
40-44	47	21.2	3.09	36	33.0	5.50	11	9.7	2.93
45-49	27	13.7	2.63	19	19.5	4.48	8	8.0	2.82
50-54	26	17.6	3.46	20	27.7	6.20	6	8.0	3.25
55-59	23	18.0	3.76	16	25.8	6.46	7	10.7	4.03
60-64	42	32.0	4.94	28	44.7	8.45	14	20.4	5.46
65-69	41	32.3	5.05	23	40.3	8.40	18	25.8	6.08
70-74	59	51.8	6.74	33	67.9	11.81	26	39.8	7.81
75-79	54	60.4	8.22	26	73.3	14.37	28	52.0	9.82
80-84	64	112.7	14.09	27	138.1	26.58	37	99.4	16.34
85+	119	247.6	22.70	34	276.6	47.43	85	237.7	25.78
Total deaths	800			511			289		
Connecticut									
. Crude rate		27.4	-		36.0	-		19.2	-
. AAMR (1940 Std. mil.)		20.7	0.83		31.5	1.48		10.1	0.77
. AAMR (1970 Std. mil.)		22.0	0.83		32.3	1.49		12.2	0.81
. AAMR (2000 Std. mil.)		25.9	0.92		37.1	1.69		15.5	0.94
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		29.5	-		43.1	-		16.4	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.0	1.22	6	5.8	2.37	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.82	2	2.3	1.61	0	0.0	0.00
15-19	30	18.1	3.30	21	24.6	5.36	9	11.2	3.73
20-24	46	23.2	3.42	41	40.7	6.35	5	5.1	2.30
25-29	29	12.7	2.36	22	19.2	4.09	7	6.2	2.33
30-34	28	10.7	2.02	24	18.3	3.74	4	3.1	1.53
35-39	19	7.6	1.75	15	12.1	3.13	4	3.2	1.60
40-44	14	6.3	1.68	9	8.2	2.75	5	4.4	1.98
45-49	11	5.6	1.68	8	8.2	2.91	3	3.0	1.73
50-54	15	10.2	2.63	12	16.6	4.80	3	4.0	2.30
55-59	9	7.1	2.35	5	8.1	3.61	4	6.1	3.05
60-64	17	13.0	3.14	12	19.2	5.53	5	7.3	3.26
65-69	11	8.7	2.61	4	7.0	3.50	7	10.0	3.79
70-74	19	16.7	3.83	7	14.4	5.44	12	18.4	5.30
75-79	14	15.7	4.19	4	11.3	5.64	10	18.6	5.87
80-84	13	22.9	6.35	5	25.6	11.44	8	21.5	7.60
85+	8	16.6	5.89	6	48.8	19.92	2	5.6	3.95
Total deaths	291			203			88		
Connecticut									
. Crude rate		10.0	-		14.3	-		5.9	-
. AAMR (1940 Std. mil.)		9.6	0.60		14.4	1.05		4.7	0.58
. AAMR (1970 Std. mil.)		9.4	0.58		13.9	1.01		5.0	0.58
. AAMR (2000 Std. mil.)		9.7	0.57		14.2	1.02		5.3	0.58
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.9	-		22.2	-		9.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is lower  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.85	2	2.3	1.65	0	0.0	0.00
20-24	2	1.0	0.71	2	2.0	1.40	0	0.0	0.00
25-29	1	0.4	0.44	1	0.9	0.87	0	0.0	0.00
30-34	2	0.8	0.54	2	1.5	1.08	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	4	1.8	0.90	3	2.7	1.59	1	0.9	0.88
45-49	2	1.0	0.72	1	1.0	1.03	1	1.0	1.00
50-54	3	2.0	1.17	1	1.4	1.39	2	2.7	1.88
55-59	6	4.7	1.92	5	8.1	3.61	1	1.5	1.52
60-64	10	7.6	2.41	6	9.6	3.91	4	5.8	2.92
65-69	5	3.9	1.76	4	7.0	3.50	1	1.4	1.43
70-74	14	12.3	3.28	9	18.5	6.17	5	7.7	3.42
75-79	18	20.1	4.75	12	33.8	9.76	6	11.1	4.55
80-84	29	51.1	9.48	11	56.3	16.97	18	48.3	11.39
85+	72	149.8	17.66	20	162.7	36.38	52	145.4	20.17
Total deaths	170			79			91		
Connecticut									
. Crude rate		5.8	-		5.6	-		6.1	-
. AAMR (1940 Std. mil.)		2.4	0.23		3.3	0.42		1.6	0.22
. AAMR (1970 Std. mil.)		3.6	0.30		4.8	0.55		2.7	0.31
. AAMR (2000 Std. mil.)		5.4	0.42		6.9	0.80		4.3	0.46
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		3.5	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	4	2.4	1.21	3	3.5	2.03	1	1.2	1.24
20-24	1	0.5	0.50	1	1.0	0.99	0	0.0	0.00
25-29	2	0.9	0.62	1	0.9	0.87	1	0.9	0.88
30-34	5	1.9	0.85	5	3.8	1.71	0	0.0	0.00
35-39	3	1.2	0.70	2	1.6	1.14	1	0.8	0.80
40-44	5	2.3	1.01	5	4.6	2.05	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	1	0.7	0.68	0	0.0	0.00	1	1.3	1.33
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.8	0.76	0	0.0	0.00	1	1.5	1.46
65-69	2	1.6	1.11	1	1.8	1.75	1	1.4	1.43
70-74	5	4.4	1.96	5	10.3	4.60	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	2	3.5	2.49	0	0.0	0.00	2	5.4	3.80
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	32			24			8		
Connecticut									
. Crude rate		1.1	-		1.7	-		0.5	-
. AAMR (1940 Std. mil.)		1.0	0.19		1.6	0.34		-	-
. AAMR (1970 Std. mil.)		1.0	0.19		1.5	0.33		-	-
. AAMR (2000 Std. mil.)		1.1	0.19		1.6	0.33		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.6	-		2.7	-		0.5	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is not significantly different

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.5	0.50	1	1.0	0.97	0	0.0	0.00
5-9	1	0.5	0.54	1	1.1	1.05	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.0	0.71	1	1.0	0.99	1	1.0	1.03
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.5	0.45	1	0.9	0.92	0	0.0	0.00
45-49	1	0.5	0.51	0	0.0	0.00	1	1.0	1.00
50-54	1	0.7	0.68	1	1.4	1.39	0	0.0	0.00
55-59	2	1.6	1.11	1	1.6	1.61	1	1.5	1.52
60-64	1	0.8	0.76	1	1.6	1.60	0	0.0	0.00
65-69	3	2.4	1.37	2	3.5	2.48	1	1.4	1.43
70-74	1	0.9	0.88	0	0.0	0.00	1	1.5	1.53
75-79	4	4.5	2.24	2	5.6	3.99	2	3.7	2.63
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	18			11			7		
Connecticut									
. Crude rate		0.6	-		0.8	-		0.5	-
. AAMR (1940 Std. mil.)		0.5	0.13		-	-		-	-
. AAMR (1970 Std. mil.)		0.6	0.14		-	-		-	-
. AAMR (2000 Std. mil.)		0.6	0.14		-	-		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.3	-		1.7	-		1.0	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: CT rate is not significantly different  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.86	1	1.0	0.97	2	2.0	1.45
5-9	1	0.5	0.54	0	0.0	0.00	1	1.1	1.11
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	13	7.8	2.17	11	12.9	3.88	2	2.5	1.76
20-24	17	8.6	2.08	14	13.9	3.71	3	3.1	1.78
25-29	19	8.3	1.91	16	14.0	3.49	3	2.6	1.53
30-34	18	6.9	1.62	16	12.2	3.06	2	1.5	1.08
35-39	11	4.4	1.33	5	4.0	1.81	6	4.8	1.96
40-44	3	1.4	0.78	1	0.9	0.92	2	1.8	1.25
45-49	5	2.5	1.13	5	5.1	2.30	0	0.0	0.00
50-54	2	1.4	0.96	2	2.8	1.96	0	0.0	0.00
55-59	2	1.6	1.11	1	1.6	1.61	1	1.5	1.52
60-64	1	0.8	0.76	1	1.6	1.60	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	2	1.8	1.24	1	2.1	2.06	1	1.5	1.53
75-79	2	2.2	1.58	1	2.8	2.82	1	1.9	1.86
80-84	1	1.8	1.76	0	0.0	0.00	1	2.7	2.69
85+	1	2.1	2.08	0	0.0	0.00	1	2.8	2.80
Total deaths	102			76			26		
Connecticut									
. Crude rate		3.5	-		5.4	-		1.7	-
. AAMR (1940 Std. mil.)		3.8	0.39		5.8	0.68		1.7	0.36
. AAMR (1970 Std. mil.)		3.4	0.36		5.1	0.62		1.6	0.34
. AAMR (2000 Std. mil.)		3.4	0.34		5.0	0.58		1.7	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.6	-		16.8	-		4.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	1.5	0.86	1	1.0	0.97	2	2.0	1.45
5-9	1	0.5	0.54	0	0.0	0.00	1	1.1	1.11
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	13	7.8	2.17	11	12.9	3.88	2	2.5	1.76
20-24	17	8.6	2.08	14	13.9	3.71	3	3.1	1.78
25-29	18	7.9	1.86	15	13.1	3.38	3	2.6	1.53
30-34	18	6.9	1.62	16	12.2	3.06	2	1.5	1.08
35-39	11	4.4	1.33	5	4.0	1.81	6	4.8	1.96
40-44	3	1.4	0.78	1	0.9	0.92	2	1.8	1.25
45-49	5	2.5	1.13	5	5.1	2.30	0	0.0	0.00
50-54	2	1.4	0.96	2	2.8	1.96	0	0.0	0.00
55-59	2	1.6	1.11	1	1.6	1.61	1	1.5	1.52
60-64	1	0.8	0.76	1	1.6	1.60	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	2	1.8	1.24	1	2.1	2.06	1	1.5	1.53
75-79	2	2.2	1.58	1	2.8	2.82	1	1.9	1.86
80-84	1	1.8	1.76	0	0.0	0.00	1	2.7	2.69
85+	1	2.1	2.08	0	0.0	0.00	1	2.8	2.80
Total deaths	101			75			26		
Connecticut									
. Crude rate		3.5	-		5.3	-		1.7	-
. AAMR (1940 Std. mil.)		3.7	0.39		5.7	0.68		1.7	0.36
. AAMR (1970 Std. mil.)		3.4	0.36		5.1	0.62		1.6	0.34
. AAMR (2000 Std. mil.)		3.3	0.34		4.9	0.58		1.7	0.34
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.5	-		16.5	-		4.4	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower



## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	3.0	1.22	6	5.8	2.37	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.2	0.82	2	2.3	1.61	0	0.0	0.00
15-19	29	17.5	3.25	20	23.4	5.23	9	11.2	3.73
20-24	44	22.2	3.35	39	38.7	6.19	5	5.1	2.30
25-29	29	12.7	2.36	22	19.2	4.09	7	6.2	2.33
30-34	28	10.7	2.02	24	18.3	3.74	4	3.1	1.53
35-39	19	7.6	1.75	15	12.1	3.13	4	3.2	1.60
40-44	14	6.3	1.68	9	8.2	2.75	5	4.4	1.98
45-49	11	5.6	1.68	8	8.2	2.91	3	3.0	1.73
50-54	15	10.2	2.63	12	16.6	4.80	3	4.0	2.30
55-59	9	7.1	2.35	5	8.1	3.61	4	6.1	3.05
60-64	16	12.2	3.05	11	17.6	5.30	5	7.3	3.26
65-69	11	8.7	2.61	4	7.0	3.50	7	10.0	3.79
70-74	19	16.7	3.83	7	14.4	5.44	12	18.4	5.30
75-79	13	14.5	4.04	4	11.3	5.64	9	16.7	5.57
80-84	13	22.9	6.35	5	25.6	11.44	8	21.5	7.60
85+	7	14.6	5.51	5	40.7	18.19	2	5.6	3.95
Total deaths	285			198			87		
Connecticut									
. Crude rate		9.8	-		14.0	-		5.8	-
. AAMR (1940 Std. mil.)		9.4	0.60		14.0	1.04		4.7	0.58
. AAMR (1970 Std. mil.)		9.2	0.58		13.5	1.00		4.9	0.58
. AAMR (2000 Std. mil.)		9.5	0.57		13.8	1.00		5.3	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.6	-		21.8	-		9.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available





## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.54	0	0.0	0.00	1	1.1	1.11
10-14	2	1.2	0.82	2	2.3	1.61	0	0.0	0.00
15-19	16	9.6	2.41	16	18.7	4.68	0	0.0	0.00
20-24	36	18.2	3.03	31	30.7	5.52	5	5.1	2.30
25-29	30	13.2	2.40	27	23.6	4.54	3	2.6	1.53
30-34	19	7.3	1.67	19	14.5	3.33	0	0.0	0.00
35-39	19	7.6	1.75	13	10.5	2.91	6	4.8	1.96
40-44	14	6.3	1.68	12	11.0	3.17	2	1.8	1.25
45-49	15	7.6	1.96	13	13.4	3.71	2	2.0	1.41
50-54	7	4.7	1.79	6	8.3	3.39	1	1.3	1.33
55-59	7	5.5	2.07	6	9.7	3.95	1	1.5	1.52
60-64	7	5.3	2.02	6	9.6	3.91	1	1.5	1.46
65-69	8	6.3	2.23	7	12.3	4.64	1	1.4	1.43
70-74	10	8.8	2.78	9	18.5	6.17	1	1.5	1.53
75-79	6	6.7	2.74	6	16.9	6.90	0	0.0	0.00
80-84	2	3.5	2.49	2	10.2	7.23	0	0.0	0.00
85+	3	6.2	3.60	3	24.4	14.09	0	0.0	0.00
Total deaths	202			178			24		
Connecticut									
. Crude rate		6.9	-		12.6	-		1.6	-
. AAMR (1940 Std. mil.)		6.9	0.51		12.4	0.97		1.6	0.34
. AAMR (1970 Std. mil.)		6.5	0.48		11.7	0.92		1.5	0.32
. AAMR (2000 Std. mil.)		6.7	0.48		12.2	0.93		1.6	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.0	-		25.8	-		4.1	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: CT rate is lower

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	6	3.6	1.48	6	7.0	2.86	0	0.0	0.00
20-24	16	8.1	2.02	15	14.9	3.84	1	1.0	1.03
25-29	18	7.9	1.86	16	14.0	3.49	2	1.8	1.25
30-34	7	2.7	1.01	7	5.4	2.02	0	0.0	0.00
35-39	11	4.4	1.33	8	6.5	2.28	3	2.4	1.39
40-44	13	5.9	1.62	11	10.1	3.04	2	1.8	1.25
45-49	10	5.1	1.60	8	8.2	2.91	2	2.0	1.41
50-54	6	4.1	1.66	5	6.9	3.10	1	1.3	1.33
55-59	6	4.7	1.92	5	8.1	3.61	1	1.5	1.52
60-64	5	3.8	1.70	5	8.0	3.57	0	0.0	0.00
65-69	8	6.3	2.23	7	12.3	4.64	1	1.4	1.43
70-74	9	7.9	2.63	9	18.5	6.17	0	0.0	0.00
75-79	6	6.7	2.74	6	16.9	6.90	0	0.0	0.00
80-84	2	3.5	2.49	2	10.2	7.23	0	0.0	0.00
85+	3	6.2	3.60	3	24.4	14.09	0	0.0	0.00
Total deaths	127			114			13		
Connecticut									
. Crude rate		4.3	-		8.0	-		0.9	-
. AAMR (1940 Std. mil.)		4.0	0.38		7.4	0.72		-	-
. AAMR (1970 Std. mil.)		3.9	0.36		7.3	0.70		-	-
. AAMR (2000 Std. mil.)		4.2	0.38		8.0	0.76		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.7	-		11.8	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.54	0	0.0	0.00	1	1.1	1.11
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	10	6.0	1.91	10	11.7	3.70	0	0.0	0.00
20-24	14	7.1	1.89	11	10.9	3.29	3	3.1	1.78
25-29	11	4.8	1.46	10	8.7	2.76	1	0.9	0.88
30-34	12	4.6	1.32	12	9.2	2.65	0	0.0	0.00
35-39	8	3.2	1.14	5	4.0	1.81	3	2.4	1.39
40-44	1	0.5	0.45	1	0.9	0.92	0	0.0	0.00
45-49	5	2.5	1.13	5	5.1	2.30	0	0.0	0.00
50-54	1	0.7	0.68	1	1.4	1.39	0	0.0	0.00
55-59	1	0.8	0.78	1	1.6	1.61	0	0.0	0.00
60-64	1	0.8	0.76	1	1.6	1.60	0	0.0	0.00
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	66			58			8		
Connecticut									
. Crude rate		2.3	-		4.1	-		0.5	-
. AAMR (1940 Std. mil.)		2.6	0.33		4.5	0.61		-	-
. AAMR (1970 Std. mil.)		2.3	0.30		4.0	0.55		-	-
. AAMR (2000 Std. mil.)		2.2	0.28		3.9	0.51		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.4	-		12.5	-		2.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

-----All poisoning (E850-E869, E950-E952, E962, E972, E980-E982)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	3	1.8	1.04	1	1.2	1.17	2	2.5	1.76
20-24	14	7.1	1.89	14	13.9	3.71	0	0.0	0.00
25-29	20	8.8	1.96	16	14.0	3.49	4	3.5	1.76
30-34	36	13.8	2.29	32	24.5	4.32	4	3.1	1.53
35-39	41	16.5	2.57	34	27.4	4.71	7	5.6	2.12
40-44	30	13.5	2.46	21	19.2	4.20	9	8.0	2.65
45-49	16	8.1	2.03	12	12.3	3.56	4	4.0	2.00
50-54	6	4.1	1.66	5	6.9	3.10	1	1.3	1.33
55-59	5	3.9	1.75	3	4.8	2.80	2	3.0	2.16
60-64	3	2.3	1.32	1	1.6	1.60	2	2.9	2.06
65-69	7	5.5	2.09	7	12.3	4.64	0	0.0	0.00
70-74	6	5.3	2.15	3	6.2	3.56	3	4.6	2.65
75-79	5	5.6	2.50	1	2.8	2.82	4	7.4	3.71
80-84	2	3.5	2.49	1	5.1	5.12	1	2.7	2.69
85+	6	12.5	5.10	5	40.7	18.19	1	2.8	2.80
Total deaths	201			157			44		
Connecticut									
. Crude rate		6.9	-		11.1	-		2.9	-
. AAMR (1940 Std. mil.)		6.1	0.45		9.9	0.81		2.5	0.40
. AAMR (1970 Std. mil.)		5.4	0.40		8.7	0.72		2.3	0.37
. AAMR (2000 Std. mil.)		6.5	0.46		10.6	0.86		2.8	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.3	-		7.1	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.5	0.50	1	1.0	0.99	0	0.0	0.00
25-29	4	1.8	0.88	2	1.7	1.23	2	1.8	1.25
30-34	12	4.6	1.32	9	6.9	2.29	3	2.3	1.32
35-39	23	9.2	1.93	16	12.9	3.23	7	5.6	2.12
40-44	17	7.7	1.86	12	11.0	3.17	5	4.4	1.98
45-49	18	9.1	2.15	15	15.4	3.98	3	3.0	1.73
50-54	16	10.8	2.71	13	18.0	5.00	3	4.0	2.30
55-59	23	18.0	3.76	18	29.1	6.85	5	7.6	3.41
60-64	13	9.9	2.75	10	16.0	5.05	3	4.4	2.53
65-69	23	18.1	3.78	14	24.5	6.56	9	12.9	4.30
70-74	11	9.7	2.91	6	12.3	5.04	5	7.7	3.42
75-79	10	11.2	3.54	8	22.5	7.97	2	3.7	2.63
80-84	6	10.6	4.31	5	25.6	11.44	1	2.7	2.69
85+	1	2.1	2.08	1	8.1	8.13	0	0.0	0.00
Total deaths	178			130			48		
Connecticut									
. Crude rate		6.1	-		9.2	-		3.2	-
. AAMR (1940 Std. mil.)		5.0	0.39		7.6	0.69		2.5	0.38
. AAMR (1970 Std. mil.)		5.0	0.39		7.9	0.71		2.5	0.37
. AAMR (2000 Std. mil.)		5.9	0.45		9.3	0.83		2.9	0.43
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.8	-		11.0	-		3.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is lower  
. THE U.S. HP 2000 target: No comparison is available

## Connecticut White Resident Age-Specific Mortality Rates by Sex, 1992

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to  
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives  
for  
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.6	0.58	1	1.1	1.14	0	0.0	0.00
15-19	2	1.2	0.85	1	1.2	1.17	1	1.2	1.24
20-24	11	5.6	1.67	11	10.9	3.29	0	0.0	0.00
25-29	16	7.0	1.76	12	10.5	3.02	4	3.5	1.76
30-34	29	11.1	2.06	27	20.6	3.97	2	1.5	1.08
35-39	32	12.9	2.27	26	21.0	4.12	6	4.8	1.96
40-44	19	8.6	1.96	12	11.0	3.17	7	6.2	2.34
45-49	14	7.1	1.90	10	10.3	3.25	4	4.0	2.00
50-54	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
55-59	3	2.4	1.36	1	1.6	1.61	2	3.0	2.16
60-64	2	1.5	1.08	1	1.6	1.60	1	1.5	1.46
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	2	1.8	1.24	1	2.1	2.06	1	1.5	1.53
75-79	2	2.2	1.58	0	0.0	0.00	2	3.7	2.63
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	2	4.2	2.94	2	16.3	11.50	0	0.0	0.00
Total deaths	135			105			30		
Connecticut									
. Crude rate		4.6	-		7.4	-		2.0	-
. AAMR (1940 Std. mil.)		4.3	0.38		6.8	0.67		1.8	0.34
. AAMR (1970 Std. mil.)		3.6	0.33		5.7	0.57		1.6	0.31
. AAMR (2000 Std. mil.)		4.3	0.38		6.8	0.68		2.0	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.3	-		5.9	-		2.7	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:  
. The CT HP 2000 target: No comparison is available  
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different  
. THE U.S. HP 2000 target: CT rate is higher