

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Septicemia (038)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	0.7	0.28	4	0.8	0.41	3	0.7	0.38
5-9	3	0.3	0.18	0	0.0	0.00	3	0.6	0.36
10-14	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
15-19	1	0.1	0.12	0	0.0	0.00	1	0.2	0.25
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	4	0.4	0.21	4	0.8	0.41	0	0.0	0.00
30-34	4	0.3	0.17	1	0.2	0.17	3	0.5	0.29
35-39	15	1.2	0.30	8	1.2	0.44	7	1.1	0.41
40-44	22	1.9	0.40	13	2.2	0.62	9	1.5	0.50
45-49	26	2.5	0.49	18	3.5	0.83	8	1.5	0.54
50-54	40	4.6	0.73	26	6.1	1.20	14	3.2	0.85
55-59	36	5.5	0.91	22	6.9	1.46	14	4.2	1.11
60-64	62	10.8	1.38	37	13.4	2.21	25	8.4	1.69
65-69	114	19.2	1.80	63	23.1	2.91	51	15.8	2.22
70-74	163	29.3	2.29	88	36.6	3.90	75	23.7	2.74
75-79	279	58.9	3.52	150	78.0	6.37	129	45.8	4.03
80-84	388	121.8	6.18	165	142.8	11.12	223	109.8	7.35
85+	611	221.8	8.97	190	255.8	18.56	421	209.3	10.20
Total deaths	1,776			789			987		
Connecticut									
. Crude rate		12.3	-		11.2	-		13.3	-
. AAMR (1940 Std. mil.)		4.5	0.13		5.6	0.22		3.8	0.16
. AAMR (1970 Std. mil.)		7.0	0.18		8.5	0.31		6.0	0.21
. AAMR (2000 Std. mil.)		10.3	0.25		12.5	0.45		8.9	0.29
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.9	-		4.4	-		3.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	18	1.9	0.45	10	2.1	0.66	8	1.8	0.62
5-9	27	2.8	0.53	18	3.6	0.84	9	1.9	0.63
10-14	22	2.4	0.51	14	3.0	0.80	8	1.8	0.63
15-19	21	2.5	0.55	12	2.8	0.80	9	2.2	0.74
20-24	52	6.4	0.89	34	8.2	1.41	18	4.6	1.08
25-29	67	7.0	0.85	41	8.5	1.32	26	5.4	1.07
30-34	157	13.1	1.04	81	13.5	1.50	76	12.7	1.46
35-39	314	24.4	1.37	137	21.3	1.82	177	27.4	2.06
40-44	540	45.8	1.97	242	41.7	2.68	298	49.8	2.88
45-49	915	88.8	2.94	409	80.5	3.98	506	96.9	4.31
50-54	1,496	173.1	4.47	679	160.3	6.15	817	185.3	6.48
55-59	1,839	279.9	6.53	973	303.6	9.73	866	257.2	8.74
60-64	2,700	472.1	9.09	1,441	523.5	13.79	1,259	424.4	11.96
65-69	4,125	694.1	10.81	2,266	831.8	17.47	1,859	577.6	13.40
70-74	5,320	954.8	13.09	2,818	1171.0	22.06	2,502	790.4	15.80
75-79	5,758	1215.0	16.01	2,944	1531.1	28.22	2,814	999.2	18.84
80-84	4,822	1513.5	21.80	2,396	2073.5	42.36	2,426	1194.9	24.26
85+	4,954	1798.6	25.55	2,034	2738.1	60.71	2,920	1451.6	26.86
Total deaths	33,147			16,549			16,598		
Connecticut									
. Crude rate		229.3	-		235.3	-		223.6	-
. AAMR (1940 Std. mil.)		118.6	0.75		137.9	1.16		105.3	0.98
. AAMR (1970 Std. mil.)		157.9	0.91		189.9	1.50		137.3	1.16
. AAMR (2000 Std. mil.)		199.2	1.10		245.4	1.93		171.6	1.36
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		131.6	-		159.6	-		111.1	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
20-24	1	0.1	0.12	1	0.2	0.24	0	0.0	0.00
25-29	3	0.3	0.18	1	0.2	0.21	2	0.4	0.30
30-34	11	0.9	0.28	6	1.0	0.41	5	0.8	0.37
35-39	15	1.2	0.30	8	1.2	0.44	7	1.1	0.41
40-44	50	4.2	0.60	28	4.8	0.91	22	3.7	0.78
45-49	57	5.5	0.73	36	7.1	1.18	21	4.0	0.88
50-54	113	13.1	1.23	73	17.2	2.02	40	9.1	1.43
55-59	154	23.4	1.89	92	28.7	2.99	62	18.4	2.34
60-64	239	41.8	2.70	138	50.1	4.27	101	34.0	3.39
65-69	423	71.2	3.46	241	88.5	5.70	182	56.5	4.19
70-74	554	99.4	4.22	307	127.6	7.28	247	78.0	4.96
75-79	624	131.7	5.27	299	155.5	8.99	325	115.4	6.40
80-84	593	186.1	7.64	263	227.6	14.03	330	162.5	8.95
85+	772	280.3	10.09	249	335.2	21.24	523	260.0	11.37
Total deaths	3,610			1,743			1,867		
Connecticut									
. Crude rate		25.0	-		24.8	-		25.2	-
. AAMR (1940 Std. mil.)		11.5	0.22		14.1	0.37		9.4	0.27
. AAMR (1970 Std. mil.)		16.1	0.28		19.8	0.48		13.4	0.34
. AAMR (2000 Std. mil.)		21.3	0.36		26.1	0.63		18.0	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.9	-		15.7	-		10.7	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Rectal cancer (154)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
30-34	1	0.1	0.08	0	0.0	0.00	1	0.2	0.17
35-39	2	0.2	0.11	2	0.3	0.22	0	0.0	0.00
40-44	12	1.0	0.29	7	1.2	0.46	5	0.8	0.37
45-49	12	1.2	0.34	11	2.2	0.65	1	0.2	0.19
50-54	30	3.5	0.63	22	5.2	1.11	8	1.8	0.64
55-59	34	5.2	0.89	22	6.9	1.46	12	3.6	1.03
60-64	44	7.7	1.16	25	9.1	1.82	19	6.4	1.47
65-69	75	12.6	1.46	41	15.1	2.35	34	10.6	1.81
70-74	116	20.8	1.93	72	29.9	3.53	44	13.9	2.10
75-79	114	24.1	2.25	53	27.6	3.79	61	21.7	2.77
80-84	84	26.4	2.88	38	32.9	5.33	46	22.7	3.34
85+	104	37.8	3.70	34	45.8	7.85	70	34.8	4.16
Total deaths	630			328			302		
Connecticut									
. Crude rate		4.4	-		4.7	-		4.1	-
. AAMR (1940 Std. mil.)		2.2	0.10		2.8	0.17		1.6	0.12
. AAMR (1970 Std. mil.)		3.0	0.12		3.8	0.21		2.3	0.14
. AAMR (2000 Std. mil.)		3.8	0.15		4.8	0.27		3.0	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		2.6	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
30-34	3	0.3	0.14	3	0.5	0.29	0	0.0	0.00
35-39	9	0.7	0.23	7	1.1	0.41	2	0.3	0.22
40-44	16	1.4	0.34	12	2.1	0.60	4	0.7	0.33
45-49	52	5.0	0.70	30	5.9	1.08	22	4.2	0.90
50-54	74	8.6	1.00	39	9.2	1.47	35	7.9	1.34
55-59	104	15.8	1.55	64	20.0	2.50	40	11.9	1.88
60-64	136	23.8	2.04	75	27.2	3.15	61	20.6	2.63
65-69	234	39.4	2.57	138	50.7	4.31	96	29.8	3.04
70-74	334	59.9	3.28	160	66.5	5.26	174	55.0	4.17
75-79	340	71.7	3.89	140	72.8	6.15	200	71.0	5.02
80-84	271	85.1	5.17	118	102.1	9.40	153	75.4	6.09
85+	271	98.4	5.98	78	105.0	11.89	193	95.9	6.91
Total deaths	1,845			864			981		
Connecticut									
. Crude rate		12.8	-		12.3	-		13.2	-
. AAMR (1940 Std. mil.)		6.4	0.17		7.5	0.27		5.5	0.21
. AAMR (1970 Std. mil.)		8.7	0.21		10.0	0.35		7.6	0.26
. AAMR (2000 Std. mil.)		11.0	0.26		12.6	0.43		9.8	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.3	-		7.4	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Lung cancer (162.2-.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.1	0.10	1	0.2	0.21	0	0.0	0.00
30-34	9	0.8	0.25	5	0.8	0.37	4	0.7	0.33
35-39	37	2.9	0.47	17	2.6	0.64	20	3.1	0.69
40-44	108	9.2	0.88	49	8.4	1.21	59	9.9	1.28
45-49	211	20.5	1.41	109	21.4	2.05	102	19.5	1.93
50-54	392	45.3	2.29	192	45.3	3.27	200	45.4	3.21
55-59	577	87.8	3.66	333	103.9	5.69	244	72.5	4.64
60-64	917	160.3	5.29	528	191.8	8.35	389	131.1	6.65
65-69	1,373	231.0	6.24	806	295.9	10.42	567	176.2	7.40
70-74	1,631	292.7	7.25	946	393.1	12.78	685	216.4	8.27
75-79	1,549	326.9	8.31	871	453.0	15.35	678	240.8	9.25
80-84	1,196	375.4	10.85	642	555.6	21.93	554	272.9	11.59
85+	746	270.8	9.92	386	519.6	26.45	360	179.0	9.43
Total deaths	8,747			4,885			3,862		
Connecticut									
. Crude rate		60.5	-		69.4	-		52.0	-
. AAMR (1940 Std. mil.)		33.4	0.40		42.2	0.65		26.7	0.49
. AAMR (1970 Std. mil.)		43.7	0.49		57.1	0.83		34.3	0.59
. AAMR (2000 Std. mil.)		52.8	0.57		70.7	1.02		41.0	0.67
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		38.6	-		54.1	-		26.6	-
. HP 2000 (1940 Std. mil.)		42.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	1	0.1	0.08	-	-	-	1	0.2	0.17
45-49	4	0.4	0.19	-	-	-	4	0.8	0.38
50-54	6	0.7	0.28	-	-	-	6	1.4	0.56
55-59	8	1.2	0.43	-	-	-	8	2.4	0.84
60-64	15	2.6	0.68	-	-	-	15	5.1	1.31
65-69	25	4.2	0.84	-	-	-	25	7.8	1.55
70-74	36	6.5	1.08	-	-	-	36	11.4	1.90
75-79	44	9.3	1.40	-	-	-	44	15.6	2.36
80-84	37	11.6	1.91	-	-	-	37	18.2	3.00
85+	31	11.3	2.02	-	-	-	31	15.4	2.77
Total deaths	207			-			207		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.10
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.12
. AAMR (2000 Std. mil.)		-	-		-	-		2.1	0.15
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	1	0.1	0.12	-	-	-	1	0.3	0.25
25-29	1	0.1	0.10	-	-	-	1	0.2	0.21
30-34	4	0.3	0.17	-	-	-	4	0.7	0.33
35-39	4	0.3	0.16	-	-	-	4	0.6	0.31
40-44	14	1.2	0.32	-	-	-	14	2.3	0.63
45-49	36	3.5	0.58	-	-	-	36	6.9	1.15
50-54	72	8.3	0.98	-	-	-	72	16.3	1.92
55-59	49	7.5	1.07	-	-	-	49	14.6	2.08
60-64	74	12.9	1.50	-	-	-	74	24.9	2.90
65-69	102	17.2	1.70	-	-	-	102	31.7	3.14
70-74	133	23.9	2.07	-	-	-	133	42.0	3.64
75-79	127	26.8	2.38	-	-	-	127	45.1	4.00
80-84	105	33.0	3.22	-	-	-	105	51.7	5.05
85+	135	49.0	4.22	-	-	-	135	67.1	5.78
Total deaths	857			-			857		
Connecticut									
. Crude rate		-	-		-	-		11.5	-
. AAMR (1940 Std. mil.)		-	-		-	-		5.9	0.24
. AAMR (1970 Std. mil.)		-	-		-	-		7.5	0.28
. AAMR (2000 Std. mil.)		-	-		-	-		9.1	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		6.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	0.5	0.24	3	0.6	0.36	2	0.4	0.31
5-9	8	0.8	0.29	6	1.2	0.49	2	0.4	0.30
10-14	5	0.5	0.24	3	0.6	0.37	2	0.4	0.32
15-19	2	0.2	0.17	0	0.0	0.00	2	0.5	0.35
20-24	5	0.6	0.28	5	1.2	0.54	0	0.0	0.00
25-29	4	0.4	0.21	4	0.8	0.41	0	0.0	0.00
30-34	12	1.0	0.29	6	1.0	0.41	6	1.0	0.41
35-39	20	1.6	0.35	15	2.3	0.60	5	0.8	0.35
40-44	24	2.0	0.42	15	2.6	0.67	9	1.5	0.50
45-49	53	5.1	0.71	32	6.3	1.11	21	4.0	0.88
50-54	66	7.6	0.94	35	8.3	1.40	31	7.0	1.26
55-59	62	9.4	1.20	32	10.0	1.77	30	8.9	1.63
60-64	54	9.4	1.28	36	13.1	2.18	18	6.1	1.43
65-69	94	15.8	1.63	51	18.7	2.62	43	13.4	2.04
70-74	102	18.3	1.81	43	17.9	2.72	59	18.6	2.43
75-79	80	16.9	1.89	42	21.8	3.37	38	13.5	2.19
80-84	70	22.0	2.63	34	29.4	5.05	36	17.7	2.96
85+	48	17.4	2.52	21	28.3	6.17	27	13.4	2.58
Total deaths	714			383			331		
Connecticut									
. Crude rate		4.9	-		5.4	-		4.5	-
. AAMR (1940 Std. mil.)		3.3	0.14		3.9	0.21		2.7	0.17
. AAMR (1970 Std. mil.)		3.9	0.15		4.7	0.24		3.2	0.19
. AAMR (2000 Std. mil.)		4.5	0.17		5.5	0.28		3.7	0.21
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.5	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	8	0.9	0.30	3	0.6	0.36	5	1.1	0.49
5-9	11	1.1	0.34	5	1.0	0.44	6	1.3	0.51
10-14	7	0.8	0.29	6	1.3	0.52	1	0.2	0.22
15-19	6	0.7	0.29	3	0.7	0.40	3	0.7	0.43
20-24	18	2.2	0.53	12	2.9	0.84	6	1.5	0.62
25-29	13	1.4	0.37	10	2.1	0.65	3	0.6	0.36
30-34	16	1.3	0.33	9	1.5	0.50	7	1.2	0.44
35-39	23	1.8	0.37	12	1.9	0.54	11	1.7	0.51
40-44	36	3.1	0.51	20	3.4	0.77	16	2.7	0.67
45-49	20	1.9	0.43	8	1.6	0.56	12	2.3	0.66
50-54	44	5.1	0.77	24	5.7	1.16	20	4.5	1.01
55-59	47	7.2	1.04	26	8.1	1.59	21	6.2	1.36
60-64	95	16.6	1.70	51	18.5	2.59	44	14.8	2.24
65-69	133	22.4	1.94	96	35.2	3.60	37	11.5	1.89
70-74	197	35.4	2.52	113	47.0	4.42	84	26.5	2.90
75-79	224	47.3	3.16	125	65.0	5.81	99	35.2	3.53
80-84	182	57.1	4.23	83	71.8	7.88	99	48.8	4.90
85+	196	71.2	5.08	83	111.7	12.26	113	56.2	5.28
Total deaths	1,276			689			587		
Connecticut									
. Crude rate		8.8	-		9.8	-		7.9	-
. AAMR (1940 Std. mil.)		4.8	0.16		6.1	0.25		3.8	0.19
. AAMR (1970 Std. mil.)		6.2	0.18		8.0	0.31		4.8	0.22
. AAMR (2000 Std. mil.)		7.7	0.22		10.2	0.39		6.1	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.9	-		6.2	-		3.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	0.4	0.21	2	0.5	0.34	1	0.3	0.25
25-29	6	0.6	0.25	3	0.6	0.36	3	0.6	0.36
30-34	29	2.4	0.45	8	1.3	0.47	21	3.5	0.77
35-39	42	3.3	0.50	29	4.5	0.84	13	2.0	0.56
40-44	98	8.3	0.84	65	11.2	1.39	33	5.5	0.96
45-49	166	16.1	1.25	98	19.3	1.95	68	13.0	1.58
50-54	273	31.6	1.91	165	39.0	3.03	108	24.5	2.36
55-59	448	68.2	3.22	273	85.2	5.16	175	52.0	3.93
60-64	680	118.9	4.56	387	140.6	7.15	293	98.8	5.77
65-69	1,169	196.7	5.75	641	235.3	9.29	528	164.0	7.14
70-74	1,631	292.7	7.25	861	357.8	12.19	770	243.2	8.77
75-79	2,105	444.2	9.68	1,072	557.5	17.03	1,033	366.8	11.41
80-84	2,111	662.6	14.42	980	848.1	27.09	1,131	557.0	16.56
85+	2,568	932.3	18.40	822	1106.5	38.60	1,746	868.0	20.77
Total deaths	11,331			5,408			5,923		
Connecticut									
. Crude rate		78.4	-		76.9	-		79.8	-
. AAMR (1940 Std. mil.)		34.0	0.37		41.6	0.62		28.2	0.46
. AAMR (1970 Std. mil.)		49.3	0.49		60.5	0.83		41.3	0.59
. AAMR (2000 Std. mil.)		66.6	0.63		81.7	1.13		56.4	0.75
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetic ketoacidosis (DKA) (250.1)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	2	0.2	0.15	1	0.2	0.21	1	0.2	0.21
30-34	3	0.3	0.14	2	0.3	0.24	1	0.2	0.17
35-39	4	0.3	0.16	3	0.5	0.27	1	0.2	0.15
40-44	4	0.3	0.17	2	0.3	0.24	2	0.3	0.24
45-49	2	0.2	0.14	2	0.4	0.28	0	0.0	0.00
50-54	3	0.3	0.20	3	0.7	0.41	0	0.0	0.00
55-59	3	0.5	0.26	2	0.6	0.44	1	0.3	0.30
60-64	1	0.2	0.17	1	0.4	0.36	0	0.0	0.00
65-69	3	0.5	0.29	2	0.7	0.52	1	0.3	0.31
70-74	7	1.3	0.47	3	1.2	0.72	4	1.3	0.63
75-79	5	1.1	0.47	2	1.0	0.74	3	1.1	0.62
80-84	6	1.9	0.77	3	2.6	1.50	3	1.5	0.85
85+	6	2.2	0.89	0	0.0	0.00	6	3.0	1.22
Total deaths	50			27			23		
Connecticut									
. Crude rate		0.3	-		0.4	-		0.3	-
. AAMR (1940 Std. mil.)		0.2	0.03		0.3	0.06		0.1	0.04
. AAMR (1970 Std. mil.)		0.2	0.04		0.3	0.06		0.2	0.04
. AAMR (2000 Std. mil.)		0.3	0.04		0.4	0.07		0.2	0.05
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		0.6	-		0.5	-		0.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Alzheimer's diseases (331.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.1	0.08	1	0.2	0.17	0	0.0	0.00
45-49	2	0.2	0.14	0	0.0	0.00	2	0.4	0.27
50-54	2	0.2	0.16	1	0.2	0.24	1	0.2	0.23
55-59	11	1.7	0.50	3	0.9	0.54	8	2.4	0.84
60-64	15	2.6	0.68	9	3.3	1.09	6	2.0	0.83
65-69	37	6.2	1.02	22	8.1	1.72	15	4.7	1.20
70-74	70	12.6	1.50	26	10.8	2.12	44	13.9	2.10
75-79	171	36.1	2.76	63	32.8	4.13	108	38.4	3.69
80-84	278	87.3	5.23	100	86.5	8.65	178	87.7	6.57
85+	642	233.1	9.20	149	200.6	16.43	493	245.1	11.04
Total deaths	1,229			374			855		
Connecticut									
. Crude rate		8.5	-		5.3	-		11.5	-
. AAMR (1940 Std. mil.)		2.2	0.07		2.1	0.12		2.3	0.10
. AAMR (1970 Std. mil.)		4.2	0.12		3.9	0.20		4.3	0.16
. AAMR (2000 Std. mil.)		7.0	0.20		6.4	0.33		7.2	0.25
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		2.6	-		2.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cardiovascular disease (390-459)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	54	5.8	0.78	31	6.4	1.16	23	5.0	1.05
5-9	6	0.6	0.25	3	0.6	0.34	3	0.6	0.36
10-14	11	1.2	0.36	5	1.1	0.48	6	1.3	0.55
15-19	16	1.9	0.48	13	3.0	0.83	3	0.7	0.43
20-24	28	3.5	0.66	14	3.4	0.90	14	3.6	0.95
25-29	69	7.2	0.86	40	8.2	1.30	29	6.1	1.13
30-34	125	10.4	0.93	81	13.5	1.50	44	7.4	1.11
35-39	261	20.2	1.25	173	26.9	2.04	88	13.6	1.45
40-44	462	39.2	1.82	333	57.4	3.14	129	21.6	1.90
45-49	779	75.6	2.71	570	112.1	4.70	209	40.0	2.77
50-54	1,104	127.7	3.84	796	187.9	6.66	308	69.9	3.98
55-59	1,538	234.1	5.97	1,109	346.1	10.39	429	127.4	6.15
60-64	2,241	391.8	8.28	1,510	548.5	14.12	731	246.4	9.11
65-69	3,938	662.6	10.56	2,477	909.2	18.27	1,461	453.9	11.88
70-74	5,941	1066.2	13.83	3,483	1447.3	24.52	2,458	776.5	15.66
75-79	8,611	1817.1	19.58	4,543	2362.7	35.05	4,068	1444.5	22.65
80-84	10,674	3350.4	32.43	4,764	4122.7	59.73	5,910	2910.8	37.86
85+	22,456	8152.9	54.41	6,328	8518.5	107.09	16,128	8017.9	63.13
Total deaths	58,314			26,273			32,041		
Connecticut									
. Crude rate		403.4	-		373.5	-		431.7	-
. AAMR (1940 Std. mil.)		145.7	0.74		190.3	1.30		109.3	0.82
. AAMR (1970 Std. mil.)		229.1	1.01		288.9	1.81		182.7	1.14
. AAMR (2000 Std. mil.)		338.8	1.41		413.9	2.60		283.1	1.61
. HP 2000 (1940 Std. mil.)		84.4	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		177.7	-		231.3	-		134.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.2	0.15	2	0.4	0.29	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
15-19	4	0.5	0.24	4	0.9	0.46	0	0.0	0.00
20-24	2	0.2	0.18	1	0.2	0.24	1	0.3	0.25
25-29	16	1.7	0.42	13	2.7	0.74	3	0.6	0.36
30-34	41	3.4	0.53	33	5.5	0.96	8	1.3	0.47
35-39	112	8.7	0.82	85	13.2	1.43	27	4.2	0.80
40-44	261	22.1	1.37	209	36.0	2.49	52	8.7	1.20
45-49	476	46.2	2.12	388	76.3	3.88	88	16.9	1.80
50-54	732	84.7	3.13	582	137.4	5.70	150	34.0	2.78
55-59	996	151.6	4.80	754	235.3	8.57	242	71.9	4.62
60-64	1,429	249.9	6.61	1,017	369.4	11.58	412	138.9	6.84
65-69	2,440	410.6	8.31	1,637	600.9	14.85	803	249.5	8.80
70-74	3,481	624.7	10.59	2,173	903.0	19.37	1,308	413.2	11.42
75-79	4,964	1047.5	14.87	2,745	1427.6	27.25	2,219	788.0	16.73
80-84	5,864	1840.6	24.04	2,771	2398.0	45.55	3,093	1523.4	27.39
85+	12,208	4432.2	40.11	3,603	4850.2	80.80	8,605	4277.9	46.12
Total deaths	33,029			16,018			17,011		
Connecticut									
. Crude rate		228.5	-		227.7	-		229.2	-
. AAMR (1940 Std. mil.)		84.9	0.57		118.9	1.03		57.2	0.59
. AAMR (1970 Std. mil.)		131.9	0.77		177.6	1.42		96.6	0.82
. AAMR (2000 Std. mil.)		192.4	1.06		250.4	2.01		149.9	1.17
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		110.2	-		151.7	-		76.6	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.1	0.08	1	0.2	0.17	0	0.0	0.00
35-39	1	0.1	0.08	1	0.2	0.16	0	0.0	0.00
40-44	6	0.5	0.21	5	0.9	0.39	1	0.2	0.17
45-49	6	0.6	0.24	4	0.8	0.39	2	0.4	0.27
50-54	15	1.7	0.45	9	2.1	0.71	6	1.4	0.56
55-59	12	1.8	0.53	7	2.2	0.83	5	1.5	0.66
60-64	25	4.4	0.87	14	5.1	1.36	11	3.7	1.12
65-69	41	6.9	1.08	24	8.8	1.80	17	5.3	1.28
70-74	84	15.1	1.64	38	15.8	2.56	46	14.5	2.14
75-79	92	19.4	2.02	38	19.8	3.21	54	19.2	2.61
80-84	113	35.5	3.34	42	36.3	5.61	71	35.0	4.15
85+	287	104.2	6.15	72	96.9	11.42	215	106.9	7.29
Total deaths	684			255			429		
Connecticut									
. Crude rate		4.7	-		3.6	-		5.8	-
. AAMR (1940 Std. mil.)		1.6	0.08		1.8	0.12		1.5	0.10
. AAMR (1970 Std. mil.)		2.6	0.11		2.8	0.18		2.5	0.13
. AAMR (2000 Std. mil.)		4.0	0.15		4.1	0.26		3.8	0.19
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		2.5	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	11	1.2	0.35	8	1.7	0.59	3	0.7	0.38
5-9	1	0.1	0.10	1	0.2	0.20	0	0.0	0.00
10-14	2	0.2	0.15	1	0.2	0.21	1	0.2	0.22
15-19	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
20-24	2	0.2	0.18	1	0.2	0.24	1	0.3	0.25
25-29	5	0.5	0.23	2	0.4	0.29	3	0.6	0.36
30-34	23	1.9	0.40	13	2.2	0.60	10	1.7	0.53
35-39	34	2.6	0.45	21	3.3	0.71	13	2.0	0.56
40-44	61	5.2	0.66	29	5.0	0.93	32	5.3	0.95
45-49	95	9.2	0.95	49	9.6	1.38	46	8.8	1.30
50-54	100	11.6	1.16	48	11.3	1.64	52	11.8	1.64
55-59	163	24.8	1.94	107	33.4	3.23	56	16.6	2.22
60-64	208	36.4	2.52	125	45.4	4.06	83	28.0	3.07
65-69	454	76.4	3.59	234	85.9	5.62	220	68.4	4.61
70-74	775	139.1	5.00	395	164.1	8.26	380	120.0	6.16
75-79	1,344	283.6	7.74	620	322.5	12.95	724	257.1	9.55
80-84	1,844	578.8	13.48	728	630.0	23.35	1,116	549.7	16.45
85+	3,800	1379.6	22.38	938	1262.7	41.23	2,862	1422.8	26.60
Total deaths	8,923			3,321			5,602		
Connecticut									
. Crude rate		61.7	-		47.2	-		75.5	-
. AAMR (1940 Std. mil.)		19.9	0.26		21.9	0.42		18.1	0.32
. AAMR (1970 Std. mil.)		33.2	0.37		35.5	0.62		31.0	0.46
. AAMR (2000 Std. mil.)		51.4	0.54		53.5	0.95		49.1	0.67
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		26.5	-		29.0	-		24.5	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Aortic aneurysm (441)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.1	0.12	1	0.2	0.24	0	0.0	0.00
25-29	1	0.1	0.10	1	0.2	0.21	0	0.0	0.00
30-34	2	0.2	0.12	2	0.3	0.24	0	0.0	0.00
35-39	8	0.6	0.22	6	0.9	0.38	2	0.3	0.22
40-44	6	0.5	0.21	5	0.9	0.39	1	0.2	0.17
45-49	5	0.5	0.22	2	0.4	0.28	3	0.6	0.33
50-54	13	1.5	0.42	10	2.4	0.75	3	0.7	0.39
55-59	23	3.5	0.73	21	6.6	1.43	2	0.6	0.42
60-64	35	6.1	1.03	28	10.2	1.92	7	2.4	0.89
65-69	125	21.0	1.88	94	34.5	3.56	31	9.6	1.73
70-74	176	31.6	2.38	116	48.2	4.48	60	19.0	2.45
75-79	203	42.8	3.01	113	58.8	5.53	90	32.0	3.37
80-84	201	63.1	4.45	101	87.4	8.70	100	49.3	4.93
85+	244	88.6	5.67	113	152.1	14.31	131	65.1	5.69
Total deaths	1,043			613			430		
Connecticut									
. Crude rate		7.2	-		8.7	-		5.8	-
. AAMR (1940 Std. mil.)		2.9	0.11		4.5	0.20		1.7	0.10
. AAMR (1970 Std. mil.)		4.4	0.14		6.8	0.28		2.7	0.14
. AAMR (2000 Std. mil.)		6.0	0.19		9.3	0.38		3.9	0.19
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.3	-		5.3	-		2.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic obstructive pulmonary disease (COPD) (490-496)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	0.2	0.15	1	0.2	0.21	1	0.2	0.22
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
15-19	6	0.7	0.29	2	0.5	0.33	4	1.0	0.49
20-24	6	0.7	0.30	4	1.0	0.48	2	0.5	0.36
25-29	3	0.3	0.18	2	0.4	0.29	1	0.2	0.21
30-34	7	0.6	0.22	3	0.5	0.29	4	0.7	0.33
35-39	17	1.3	0.32	6	0.9	0.38	11	1.7	0.51
40-44	25	2.1	0.42	11	1.9	0.57	14	2.3	0.63
45-49	34	3.3	0.57	14	2.8	0.74	20	3.8	0.86
50-54	65	7.5	0.93	29	6.8	1.27	36	8.2	1.36
55-59	125	19.0	1.70	47	14.7	2.14	78	23.2	2.62
60-64	267	46.7	2.86	126	45.8	4.08	141	47.5	4.00
65-69	565	95.1	4.00	290	106.5	6.25	275	85.4	5.15
70-74	890	159.7	5.35	430	178.7	8.62	460	145.3	6.78
75-79	1,256	265.0	7.48	601	312.6	12.75	655	232.6	9.09
80-84	1,217	382.0	10.95	565	488.9	20.57	652	321.1	12.58
85+	1,386	503.2	13.52	587	790.2	32.62	799	397.2	14.05
Total deaths	5,872			2,719			3,153		
Connecticut									
. Crude rate		40.6	-		38.7	-		42.5	-
. AAMR (1940 Std. mil.)		16.1	0.25		18.4	0.39		14.9	0.33
. AAMR (1970 Std. mil.)		24.5	0.34		29.3	0.57		22.0	0.43
. AAMR (2000 Std. mil.)		34.0	0.45		42.3	0.83		30.0	0.54
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		21.1	-		26.9	-		17.0	-
. HP 2000 (1940 Std. mil.)		25.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.1	0.12	1	0.2	0.23	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	6	0.6	0.25	3	0.6	0.36	3	0.6	0.36
30-34	20	1.7	0.37	13	2.2	0.60	7	1.2	0.44
35-39	54	4.2	0.57	35	5.4	0.92	19	2.9	0.67
40-44	105	8.9	0.87	79	13.6	1.53	26	4.3	0.85
45-49	147	14.3	1.18	109	21.4	2.05	38	7.3	1.18
50-54	142	16.4	1.38	112	26.4	2.50	30	6.8	1.24
55-59	120	18.3	1.67	85	26.5	2.88	35	10.4	1.76
60-64	121	21.2	1.92	82	29.8	3.29	39	13.1	2.11
65-69	176	29.6	2.23	109	40.0	3.83	67	20.8	2.54
70-74	193	34.6	2.49	106	44.0	4.28	87	27.5	2.95
75-79	162	34.2	2.69	69	35.9	4.32	93	33.0	3.42
80-84	104	32.6	3.20	46	39.8	5.87	58	28.6	3.75
85+	55	20.0	2.69	19	25.6	5.87	36	17.9	2.98
Total deaths	1,407			869			538		
Connecticut									
. Crude rate		9.7	-		12.4	-		7.2	-
. AAMR (1940 Std. mil.)		6.6	0.19		9.3	0.33		4.1	0.20
. AAMR (1970 Std. mil.)		7.6	0.21		10.5	0.36		4.9	0.22
. AAMR (2000 Std. mil.)		8.9	0.24		12.3	0.42		5.9	0.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.8	-		11.3	-		4.8	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									CT rate is higher

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----End stage renal disease (ESRD) (581-588,591,593.8,593.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	6	0.6	0.26	3	0.6	0.36	3	0.7	0.38
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	0.2	0.17	2	0.5	0.33	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	7	0.7	0.27	5	1.0	0.46	2	0.4	0.30
30-34	4	0.3	0.17	3	0.5	0.29	1	0.2	0.17
35-39	16	1.2	0.31	4	0.6	0.31	12	1.9	0.54
40-44	15	1.3	0.33	9	1.6	0.52	6	1.0	0.41
45-49	24	2.3	0.48	15	3.0	0.76	9	1.7	0.57
50-54	29	3.4	0.62	19	4.5	1.03	10	2.3	0.72
55-59	39	5.9	0.95	19	5.9	1.36	20	5.9	1.33
60-64	63	11.0	1.39	37	13.4	2.21	26	8.8	1.72
65-69	119	20.0	1.84	62	22.8	2.89	57	17.7	2.35
70-74	210	37.7	2.60	114	47.4	4.44	96	30.3	3.10
75-79	301	63.5	3.66	162	84.3	6.62	139	49.4	4.19
80-84	366	114.9	6.00	191	165.3	11.96	175	86.2	6.52
85+	629	228.4	9.11	275	370.2	22.32	354	176.0	9.35
Total deaths	1,830			920			910		
Connecticut									
. Crude rate		12.7	-		13.1	-		12.3	-
. AAMR (1940 Std. mil.)		4.7	0.13		6.1	0.22		3.8	0.16
. AAMR (1970 Std. mil.)		7.3	0.18		9.9	0.33		5.8	0.21
. AAMR (2000 Std. mil.)		10.6	0.25		14.9	0.50		8.4	0.28
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.2	-		6.4	-		4.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	67	7.1	0.87	42	8.7	1.34	25	5.5	1.10
5-9	35	3.6	0.60	28	5.6	1.05	7	1.5	0.55
10-14	56	6.1	0.82	32	6.8	1.21	24	5.3	1.09
15-19	227	27.0	1.79	155	35.7	2.87	72	17.7	2.09
20-24	268	33.2	2.03	199	48.1	3.41	69	17.6	2.12
25-29	288	29.9	1.76	232	47.8	3.14	56	11.7	1.57
30-34	355	29.6	1.57	279	46.4	2.78	76	12.7	1.46
35-39	369	28.6	1.49	288	44.8	2.64	81	12.5	1.39
40-44	327	27.7	1.53	265	45.6	2.80	62	10.4	1.32
45-49	268	26.0	1.59	197	38.8	2.76	71	13.6	1.61
50-54	164	19.0	1.48	120	28.3	2.59	44	10.0	1.50
55-59	131	19.9	1.74	81	25.3	2.81	50	14.9	2.10
60-64	151	26.4	2.15	93	33.8	3.50	58	19.6	2.57
65-69	176	29.6	2.23	111	40.7	3.87	65	20.2	2.50
70-74	285	51.1	3.03	153	63.6	5.14	132	41.7	3.63
75-79	357	75.3	3.99	194	100.9	7.24	163	57.9	4.53
80-84	437	137.2	6.56	205	177.4	12.39	232	114.3	7.50
85+	733	266.1	9.83	291	391.7	22.96	442	219.7	10.45
Total deaths	4,694			2,965			1,729		
Connecticut									
. Crude rate		32.5	-		42.2	-		23.3	-
. AAMR (1940 Std. mil.)		24.4	0.42		35.3	0.71		13.9	0.44
. AAMR (1970 Std. mil.)		25.7	0.41		36.6	0.70		15.8	0.44
. AAMR (2000 Std. mil.)		30.3	0.45		43.5	0.81		19.0	0.47
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.2	-		43.9	-		17.1	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	1.5	0.40	6	1.2	0.51	8	1.8	0.62
5-9	18	1.8	0.43	16	3.2	0.79	2	0.4	0.30
10-14	36	3.9	0.65	21	4.5	0.98	15	3.3	0.86
15-19	168	20.0	1.54	106	24.4	2.37	62	15.3	1.94
20-24	181	22.4	1.67	131	31.7	2.77	50	12.7	1.80
25-29	148	15.4	1.26	117	24.1	2.23	31	6.5	1.17
30-34	122	10.2	0.92	88	14.6	1.56	34	5.7	0.97
35-39	107	8.3	0.80	72	11.2	1.32	35	5.4	0.92
40-44	84	7.1	0.78	59	10.2	1.32	25	4.2	0.84
45-49	96	9.3	0.95	68	13.4	1.62	28	5.4	1.01
50-54	74	8.6	1.00	54	12.7	1.73	20	4.5	1.01
55-59	46	7.0	1.03	26	8.1	1.59	20	5.9	1.33
60-64	65	11.4	1.41	35	12.7	2.15	30	10.1	1.85
65-69	52	8.7	1.21	32	11.7	2.08	20	6.2	1.39
70-74	79	14.2	1.60	41	17.0	2.66	38	12.0	1.95
75-79	74	15.6	1.82	36	18.7	3.12	38	13.5	2.19
80-84	73	22.9	2.68	39	33.8	5.40	34	16.7	2.87
85+	58	21.1	2.76	26	35.0	6.86	32	15.9	2.81
Total deaths	1,495			973			522		
Connecticut									
. Crude rate		10.3	-		13.8	-		7.0	-
. AAMR (1940 Std. mil.)		10.3	0.29		14.0	0.48		6.6	0.33
. AAMR (1970 Std. mil.)		10.1	0.28		13.6	0.46		6.7	0.32
. AAMR (2000 Std. mil.)		10.3	0.27		14.1	0.46		6.8	0.31
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		16.1	-		22.5	-		9.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
5-9	1	0.1	0.10	1	0.2	0.20	0	0.0	0.00
10-14	2	0.2	0.15	1	0.2	0.21	1	0.2	0.22
15-19	3	0.4	0.21	3	0.7	0.40	0	0.0	0.00
20-24	12	1.5	0.43	11	2.7	0.80	1	0.3	0.25
25-29	6	0.6	0.25	5	1.0	0.46	1	0.2	0.21
30-34	13	1.1	0.30	11	1.8	0.55	2	0.3	0.24
35-39	12	0.9	0.27	10	1.6	0.49	2	0.3	0.22
40-44	16	1.4	0.34	13	2.2	0.62	3	0.5	0.29
45-49	21	2.0	0.44	15	3.0	0.76	6	1.1	0.47
50-54	18	2.1	0.49	12	2.8	0.82	6	1.4	0.56
55-59	17	2.6	0.63	14	4.4	1.17	3	0.9	0.51
60-64	21	3.7	0.80	16	5.8	1.45	5	1.7	0.75
65-69	45	7.6	1.13	31	11.4	2.04	14	4.3	1.16
70-74	78	14.0	1.58	42	17.5	2.69	36	11.4	1.90
75-79	116	24.5	2.27	69	35.9	4.32	47	16.7	2.43
80-84	183	57.4	4.25	82	71.0	7.84	101	49.7	4.95
85+	405	147.0	7.31	143	192.5	16.10	262	130.3	8.05
Total deaths	970			479			491		
Connecticut									
. Crude rate		6.7	-		6.8	-		6.6	-
. AAMR (1940 Std. mil.)		2.5	0.10		3.6	0.19		1.6	0.10
. AAMR (1970 Std. mil.)		3.8	0.13		5.3	0.25		2.7	0.14
. AAMR (2000 Std. mil.)		5.7	0.18		7.8	0.36		4.3	0.20
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.3	-		3.7	-		1.6	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	14	1.5	0.40	11	2.3	0.69	3	0.7	0.38
5-9	3	0.3	0.18	3	0.6	0.34	0	0.0	0.00
10-14	2	0.2	0.15	0	0.0	0.00	2	0.4	0.32
15-19	15	1.8	0.46	13	3.0	0.83	2	0.5	0.35
20-24	7	0.9	0.33	7	1.7	0.64	0	0.0	0.00
25-29	13	1.4	0.37	12	2.5	0.71	1	0.2	0.21
30-34	15	1.3	0.32	15	2.5	0.64	0	0.0	0.00
35-39	15	1.2	0.30	12	1.9	0.54	3	0.5	0.27
40-44	7	0.6	0.22	5	0.9	0.39	2	0.3	0.24
45-49	5	0.5	0.22	4	0.8	0.39	1	0.2	0.19
50-54	5	0.6	0.26	5	1.2	0.53	0	0.0	0.00
55-59	9	1.4	0.46	8	2.5	0.88	1	0.3	0.30
60-64	4	0.7	0.35	2	0.7	0.51	2	0.7	0.48
65-69	7	1.2	0.45	5	1.8	0.82	2	0.6	0.44
70-74	3	0.5	0.31	1	0.4	0.42	2	0.6	0.45
75-79	7	1.5	0.56	5	2.6	1.16	2	0.7	0.50
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	4	1.5	0.73	3	4.0	2.33	1	0.5	0.50
Total deaths	135			111			24		
Connecticut									
. Crude rate		0.9	-		1.6	-		0.3	-
. AAMR (1940 Std. mil.)		1.0	0.09		1.6	0.16		0.3	0.07
. AAMR (1970 Std. mil.)		0.9	0.08		1.5	0.15		0.3	0.07
. AAMR (2000 Std. mil.)		0.9	0.08		1.6	0.15		0.3	0.07
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		2.5	-		0.4	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	1.4	0.38	7	1.5	0.55	6	1.3	0.54
5-9	6	0.6	0.25	3	0.6	0.34	3	0.6	0.36
10-14	5	0.5	0.24	5	1.1	0.48	0	0.0	0.00
15-19	6	0.7	0.29	1	0.2	0.23	5	1.2	0.55
20-24	6	0.7	0.30	5	1.2	0.54	1	0.3	0.25
25-29	5	0.5	0.23	3	0.6	0.36	2	0.4	0.30
30-34	4	0.3	0.17	3	0.5	0.29	1	0.2	0.17
35-39	8	0.6	0.22	6	0.9	0.38	2	0.3	0.22
40-44	8	0.7	0.24	6	1.0	0.42	2	0.3	0.24
45-49	11	1.1	0.32	7	1.4	0.52	4	0.8	0.38
50-54	7	0.8	0.31	4	0.9	0.47	3	0.7	0.39
55-59	10	1.5	0.48	5	1.6	0.70	5	1.5	0.66
60-64	7	1.2	0.46	4	1.5	0.73	3	1.0	0.58
65-69	6	1.0	0.41	4	1.5	0.73	2	0.6	0.44
70-74	12	2.2	0.62	6	2.5	1.02	6	1.9	0.77
75-79	8	1.7	0.60	3	1.6	0.90	5	1.8	0.79
80-84	11	3.5	1.04	6	5.2	2.12	5	2.5	1.10
85+	9	3.3	1.09	3	4.0	2.33	6	3.0	1.22
Total deaths	142			81			61		
Connecticut									
. Crude rate		1.0	-		1.2	-		0.8	-
. AAMR (1940 Std. mil.)		0.8	0.08		1.0	0.12		0.7	0.10
. AAMR (1970 Std. mil.)		0.9	0.08		1.1	0.13		0.7	0.10
. AAMR (2000 Std. mil.)		1.0	0.08		1.2	0.13		0.8	0.10
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.2	-		1.7	-		1.0	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	19	2.1	0.48	15	3.2	0.83	4	0.9	0.45
15-19	51	6.1	0.85	41	9.5	1.48	10	2.5	0.78
20-24	94	11.7	1.20	79	19.1	2.15	15	3.8	0.99
25-29	107	11.1	1.08	90	18.6	1.96	17	3.6	0.86
30-34	159	13.3	1.05	123	20.5	1.84	36	6.0	1.00
35-39	158	12.3	0.97	121	18.8	1.71	37	5.7	0.94
40-44	154	13.1	1.05	120	20.7	1.89	34	5.7	0.97
45-49	113	11.0	1.03	82	16.1	1.78	31	5.9	1.07
50-54	97	11.2	1.14	68	16.1	1.95	29	6.6	1.22
55-59	75	11.4	1.32	58	18.1	2.38	17	5.0	1.22
60-64	71	12.4	1.47	59	21.4	2.79	12	4.0	1.17
65-69	69	11.6	1.40	50	18.4	2.60	19	5.9	1.35
70-74	57	10.2	1.35	45	18.7	2.79	12	3.8	1.09
75-79	48	10.1	1.46	39	20.3	3.25	9	3.2	1.07
80-84	53	16.6	2.29	40	34.6	5.47	13	6.4	1.78
85+	38	13.8	2.24	32	43.1	7.62	6	3.0	1.22
Total deaths	1,363			1,062			301		
Connecticut									
. Crude rate		9.4	-		15.1	-		4.1	-
. AAMR (1940 Std. mil.)		8.6	0.25		13.7	0.44		3.6	0.23
. AAMR (1970 Std. mil.)		8.2	0.23		13.2	0.42		3.4	0.21
. AAMR (2000 Std. mil.)		9.1	0.25		15.0	0.46		3.9	0.23
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.3	-		18.7	-		4.1	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is higher
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide (E960-E969)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	18	1.9	0.45	7	1.5	0.55	11	2.4	0.73
5-9	4	0.4	0.20	2	0.4	0.28	2	0.4	0.30
10-14	7	0.8	0.29	4	0.9	0.43	3	0.7	0.39
15-19	57	6.8	0.90	46	10.6	1.56	11	2.7	0.82
20-24	68	8.4	1.02	61	14.7	1.89	7	1.8	0.67
25-29	57	5.9	0.78	41	8.5	1.32	16	3.4	0.84
30-34	37	3.1	0.51	26	4.3	0.85	11	1.8	0.55
35-39	31	2.4	0.43	24	3.7	0.76	7	1.1	0.41
40-44	36	3.1	0.51	27	4.7	0.89	9	1.5	0.50
45-49	15	1.5	0.38	9	1.8	0.59	6	1.1	0.47
50-54	27	3.1	0.60	18	4.2	1.00	9	2.0	0.68
55-59	11	1.7	0.50	9	2.8	0.94	2	0.6	0.42
60-64	8	1.4	0.49	5	1.8	0.81	3	1.0	0.58
65-69	11	1.9	0.56	8	2.9	1.04	3	0.9	0.54
70-74	8	1.4	0.51	4	1.7	0.83	4	1.3	0.63
75-79	4	0.8	0.42	1	0.5	0.52	3	1.1	0.62
80-84	5	1.6	0.70	2	1.7	1.22	3	1.5	0.85
85+	2	0.7	0.51	1	1.3	1.35	1	0.5	0.50
Total deaths	406			295			111		
Connecticut									
. Crude rate		2.8	-		4.2	-		1.5	-
. AAMR (1940 Std. mil.)		3.2	0.17		4.8	0.29		1.6	0.16
. AAMR (1970 Std. mil.)		3.0	0.16		4.5	0.27		1.5	0.16
. AAMR (2000 Std. mil.)		2.9	0.15		4.3	0.25		1.5	0.15
. HP 2000 (1940 Std. mil.)		5.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.1	-		15.9	-		4.0	-
. HP 2000 (1940 Std. mil.)		7.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All injury (E800-E999)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	87	9.3	0.99	50	10.4	1.47	37	8.1	1.33
5-9	39	4.0	0.64	30	6.0	1.09	9	1.9	0.63
10-14	82	9.0	0.99	51	10.9	1.53	31	6.9	1.24
15-19	336	40.0	2.18	243	56.0	3.59	93	22.9	2.38
20-24	434	53.8	2.58	341	82.4	4.46	93	23.7	2.46
25-29	467	48.5	2.25	374	77.1	3.99	93	19.5	2.02
30-34	570	47.5	1.99	439	73.0	3.48	131	21.9	1.91
35-39	587	45.5	1.88	452	70.3	3.30	135	20.9	1.80
40-44	543	46.1	1.98	433	74.6	3.58	110	18.4	1.75
45-49	416	40.4	1.98	299	58.8	3.40	117	22.4	2.07
50-54	294	34.0	1.98	212	50.0	3.44	82	18.6	2.05
55-59	220	33.5	2.26	148	46.2	3.80	72	21.4	2.52
60-64	231	40.4	2.66	157	57.0	4.55	74	24.9	2.90
65-69	260	43.7	2.71	171	62.8	4.80	89	27.7	2.93
70-74	351	63.0	3.36	203	84.4	5.92	148	46.8	3.84
75-79	410	86.5	4.27	235	122.2	7.97	175	62.1	4.70
80-84	497	156.0	7.00	249	215.5	13.66	248	122.1	7.76
85+	773	280.6	10.09	324	436.2	24.23	449	223.2	10.53
Total deaths	6,597			4,411			2,186		
Connecticut									
. Crude rate		45.6	-		62.7	-		29.5	-
. AAMR (1940 Std. mil.)		37.0	0.52		54.9	0.89		19.8	0.53
. AAMR (1970 Std. mil.)		37.6	0.50		55.2	0.87		21.3	0.52
. AAMR (2000 Std. mil.)		43.2	0.54		64.0	0.97		25.0	0.55
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		53.0	-		80.9	-		25.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	13	1.4	0.38	6	1.2	0.51	7	1.5	0.58
5-9	16	1.6	0.41	14	2.8	0.74	2	0.4	0.30
10-14	36	3.9	0.65	21	4.5	0.98	15	3.3	0.86
15-19	165	19.7	1.53	103	23.7	2.34	62	15.3	1.94
20-24	177	21.9	1.65	127	30.7	2.72	50	12.7	1.80
25-29	143	14.9	1.24	113	23.3	2.19	30	6.3	1.15
30-34	117	9.8	0.90	82	13.6	1.51	35	5.8	0.99
35-39	103	8.0	0.79	69	10.7	1.29	34	5.3	0.90
40-44	82	7.0	0.77	57	9.8	1.30	25	4.2	0.84
45-49	96	9.3	0.95	68	13.4	1.62	28	5.4	1.01
50-54	73	8.4	0.99	53	12.5	1.72	20	4.5	1.01
55-59	46	7.0	1.03	26	8.1	1.59	20	5.9	1.33
60-64	62	10.8	1.38	34	12.4	2.12	28	9.4	1.78
65-69	53	8.9	1.23	33	12.1	2.11	20	6.2	1.39
70-74	76	13.6	1.56	40	16.6	2.63	36	11.4	1.90
75-79	73	15.4	1.80	36	18.7	3.12	37	13.1	2.16
80-84	71	22.3	2.64	37	32.0	5.26	34	16.7	2.87
85+	56	20.3	2.72	25	33.7	6.73	31	15.4	2.77
Total deaths	1,458			944			514		
Connecticut									
. Crude rate		10.1	-		13.4	-		6.9	-
. AAMR (1940 Std. mil.)		10.1	0.29		13.6	0.47		6.5	0.33
. AAMR (1970 Std. mil.)		9.9	0.28		13.2	0.45		6.6	0.32
. AAMR (2000 Std. mil.)		10.1	0.27		13.7	0.45		6.7	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.8	-		22.0	-		9.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
5-9	1	0.1	0.10	1	0.2	0.20	0	0.0	0.00
10-14	2	0.2	0.15	1	0.2	0.21	1	0.2	0.22
15-19	3	0.4	0.21	3	0.7	0.40	0	0.0	0.00
20-24	12	1.5	0.43	11	2.7	0.80	1	0.3	0.25
25-29	9	0.9	0.31	7	1.4	0.55	2	0.4	0.30
30-34	15	1.3	0.32	13	2.2	0.60	2	0.3	0.24
35-39	16	1.2	0.31	13	2.0	0.56	3	0.5	0.27
40-44	17	1.4	0.35	14	2.4	0.64	3	0.5	0.29
45-49	25	2.4	0.49	17	3.3	0.81	8	1.5	0.54
50-54	20	2.3	0.52	13	3.1	0.85	7	1.6	0.60
55-59	16	2.4	0.61	13	4.1	1.13	3	0.9	0.51
60-64	24	4.2	0.86	19	6.9	1.58	5	1.7	0.75
65-69	42	7.1	1.09	31	11.4	2.04	11	3.4	1.03
70-74	64	11.5	1.44	39	16.2	2.60	25	7.9	1.58
75-79	96	20.3	2.07	59	30.7	3.99	37	13.1	2.16
80-84	122	38.3	3.47	61	52.8	6.76	61	30.0	3.85
85+	239	86.8	5.61	87	117.1	12.56	152	75.6	6.13
Total deaths	724			402			322		
Connecticut									
. Crude rate		5.0	-		5.7	-		4.3	-
. AAMR (1940 Std. mil.)		2.2	0.10		3.4	0.19		1.3	0.10
. AAMR (1970 Std. mil.)		3.1	0.12		4.5	0.23		1.9	0.12
. AAMR (2000 Std. mil.)		4.3	0.16		6.3	0.32		2.9	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.5	-		3.4	-		1.4	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3, E922.8-.9, E955.0-.4, E965.0-.4, E970, E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
5-9	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
10-14	8	0.9	0.31	7	1.5	0.57	1	0.2	0.22
15-19	60	7.1	0.92	53	12.2	1.68	7	1.7	0.65
20-24	105	13.0	1.27	98	23.7	2.39	7	1.8	0.67
25-29	89	9.2	0.98	75	15.5	1.79	14	2.9	0.78
30-34	95	7.9	0.81	85	14.1	1.53	10	1.7	0.53
35-39	75	5.8	0.67	64	9.9	1.24	11	1.7	0.51
40-44	79	6.7	0.75	69	11.9	1.43	10	1.7	0.53
45-49	53	5.1	0.71	41	8.1	1.26	12	2.3	0.66
50-54	52	6.0	0.83	42	9.9	1.53	10	2.3	0.72
55-59	39	5.9	0.95	35	10.9	1.85	4	1.2	0.59
60-64	30	5.2	0.96	29	10.5	1.96	1	0.3	0.34
65-69	36	6.1	1.01	31	11.4	2.04	5	1.6	0.69
70-74	30	5.4	0.98	26	10.8	2.12	4	1.3	0.63
75-79	22	4.6	0.99	22	11.4	2.44	0	0.0	0.00
80-84	23	7.2	1.51	22	19.0	4.06	1	0.5	0.49
85+	20	7.3	1.62	19	25.6	5.87	1	0.5	0.50
Total deaths	818			719			99		
Connecticut									
. Crude rate		5.7	-		10.2	-		1.3	-
. AAMR (1940 Std. mil.)		5.7	0.21		10.1	0.40		1.4	0.15
. AAMR (1970 Std. mil.)		5.3	0.20		9.6	0.38		1.3	0.13
. AAMR (2000 Std. mil.)		5.6	0.20		10.3	0.39		1.3	0.14
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		14.9	-		26.1	-		4.2	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.1	0.11	1	0.2	0.21	0	0.0	0.00
5-9	1	0.1	0.10	0	0.0	0.00	1	0.2	0.21
10-14	3	0.3	0.19	2	0.4	0.30	1	0.2	0.22
15-19	39	4.6	0.74	33	7.6	1.32	6	1.5	0.60
20-24	58	7.2	0.94	55	13.3	1.79	3	0.8	0.44
25-29	41	4.3	0.67	32	6.6	1.17	9	1.9	0.63
30-34	22	1.8	0.39	19	3.2	0.72	3	0.5	0.29
35-39	20	1.6	0.35	17	2.6	0.64	3	0.5	0.27
40-44	16	1.4	0.34	15	2.6	0.67	1	0.2	0.17
45-49	10	1.0	0.31	6	1.2	0.48	4	0.8	0.38
50-54	17	2.0	0.48	12	2.8	0.82	5	1.1	0.51
55-59	6	0.9	0.37	5	1.6	0.70	1	0.3	0.30
60-64	2	0.3	0.25	1	0.4	0.36	1	0.3	0.34
65-69	3	0.5	0.29	0	0.0	0.00	3	0.9	0.54
70-74	2	0.4	0.25	0	0.0	0.00	2	0.6	0.45
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	241			198			43		
Connecticut									
. Crude rate		1.7	-		2.8	-		0.6	-
. AAMR (1940 Std. mil.)		2.1	0.14		3.4	0.25		0.7	0.11
. AAMR (1970 Std. mil.)		1.9	0.13		3.1	0.23		0.6	0.10
. AAMR (2000 Std. mil.)		1.8	0.12		2.9	0.21		0.6	0.09
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.5	-		12.4	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	0.2	0.18	2	0.5	0.34	0	0.0	0.00
25-29	7	0.7	0.27	7	1.4	0.55	0	0.0	0.00
30-34	27	2.3	0.43	22	3.7	0.78	5	0.8	0.37
35-39	54	4.2	0.57	41	6.4	1.00	13	2.0	0.56
40-44	89	7.5	0.80	67	11.5	1.41	22	3.7	0.78
45-49	118	11.5	1.05	95	18.7	1.92	23	4.4	0.92
50-54	99	11.5	1.15	78	18.4	2.08	21	4.8	1.04
55-59	64	9.7	1.22	47	14.7	2.14	17	5.0	1.22
60-64	86	15.0	1.62	64	23.2	2.91	22	7.4	1.58
65-69	86	14.5	1.56	52	19.1	2.65	34	10.6	1.81
70-74	58	10.4	1.37	39	16.2	2.60	19	6.0	1.38
75-79	44	9.3	1.40	28	14.6	2.75	16	5.7	1.42
80-84	19	6.0	1.37	16	13.8	3.46	3	1.5	0.85
85+	10	3.6	1.15	8	10.8	3.81	2	1.0	0.70
Total deaths	763			566			197		
Connecticut									
. Crude rate		5.3	-		8.0	-		2.7	-
. AAMR (1940 Std. mil.)		4.2	0.16		6.5	0.28		2.0	0.15
. AAMR (1970 Std. mil.)		4.3	0.16		6.8	0.29		2.1	0.15
. AAMR (2000 Std. mil.)		5.0	0.18		7.9	0.33		2.4	0.17
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.9	-		10.9	-		3.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1994-1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	3	0.3	0.18	1	0.2	0.21	2	0.4	0.31
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.1	0.11	0	0.0	0.00	1	0.2	0.22
15-19	16	1.9	0.48	12	2.8	0.80	4	1.0	0.49
20-24	57	7.1	0.94	41	9.9	1.55	16	4.1	1.02
25-29	106	11.0	1.07	86	17.7	1.91	20	4.2	0.94
30-34	197	16.4	1.17	145	24.1	2.00	52	8.7	1.20
35-39	247	19.2	1.22	188	29.2	2.13	59	9.1	1.19
40-44	213	18.1	1.24	169	29.1	2.24	44	7.4	1.11
45-49	121	11.7	1.07	83	16.3	1.79	38	7.3	1.18
50-54	36	4.2	0.69	24	5.7	1.16	12	2.7	0.79
55-59	27	4.1	0.79	12	3.7	1.08	15	4.5	1.15
60-64	11	1.9	0.58	5	1.8	0.81	6	2.0	0.83
65-69	8	1.3	0.48	2	0.7	0.52	6	1.9	0.76
70-74	14	2.5	0.67	8	3.3	1.18	6	1.9	0.77
75-79	11	2.3	0.70	6	3.1	1.27	5	1.8	0.79
80-84	12	3.8	1.09	4	3.5	1.73	8	3.9	1.39
85+	9	3.3	1.09	2	2.7	1.90	7	3.5	1.32
Total deaths	1,089			788			301		
Connecticut									
. Crude rate		7.5	-		11.2	-		4.1	-
. AAMR (1940 Std. mil.)		7.0	0.22		10.3	0.38		3.7	0.23
. AAMR (1970 Std. mil.)		6.0	0.19		8.7	0.32		3.3	0.20
. AAMR (2000 Std. mil.)		7.2	0.22		10.6	0.38		3.9	0.23
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.0	-		7.0	-		3.0	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: CT rate is higher