

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Causes of Death for the Age-Specific Mortality Tables

	Page		Page
All causes	2	Hypertension with or without renal disease	33
Infectious diseases and parasitic disease	3	Hypertension-related	34
Septicemia	4	Cerebrovascular disease	35
Septicemia-related	5	Atherosclerosis	36
HIV infection	6	Atherosclerosis-related	37
All cancer	7	Aortic aneurysm	38
Oral cavity or pharynx	8	Pneumonia & influenza	39
Colorectal cancer	9	Chronic obstructive pulmonary disease	40
Colon cancer	10	COPD-related	41
Rectal cancer	11	Asthma	42
Pancreatic cancer	12	Chronic liver disease and cirrhosis	43
Lung and other respiratory cancer	13	Nephritis, nephrotic syndrome/nephrosis	44
Lung cancer	14	Nephritis-related	45
Malignant melanoma of skin	15	End stage renal disease	46
Female breast cancer	16	Unintentional injuries	47
Cervical cancer	17	Motor vehicle accidents	48
Endometrial cancer	18	Falls and fall related injuries	49
Ovarian cancer	19	Drowning	50
Prostate cancer	20	Residential fires	51
Bladder cancer	21	Suicide	52
Brain & other CNS cancer	22	Homicide and legal intervention	53
Leukemia	23	Homicide	54
Diabetes mellitus	24	All injury	55
Diabetes-related	25	All motor vehicle accidents	56
Diabetic ketoacidosis	26	All falls	57
Alzheimer's diseases	27	All firearms	58
Cardiovascular disease	28	Suicide	59
Diseases of the heart	29	Homicide	60
Heart disease-related	30	All poisoning	61
Coronary heart disease	31	Alcohol-induced	62
Congestive heart failure	32	Drug-induced	63

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Infectious diseases and parasitic disease (1-139)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	7	3.9	1.47	4	4.3	2.17	3	3.4	1.98
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	2	1.1	0.75	0	0.0	0.00	2	2.2	1.53
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	3	2.0	1.14	1	1.3	1.29	2	2.7	1.91
25-29	6	3.4	1.38	6	6.7	2.74	0	0.0	0.00
30-34	19	8.4	1.94	12	10.7	3.09	7	6.2	2.35
35-39	24	9.3	1.91	14	10.9	2.92	10	7.8	2.46
40-44	34	13.8	2.36	26	21.4	4.20	8	6.4	2.25
45-49	35	16.9	2.86	25	24.5	4.90	10	9.5	3.01
50-54	23	12.5	2.60	18	19.9	4.70	5	5.3	2.38
55-59	24	16.9	3.46	17	24.6	5.96	7	9.6	3.65
60-64	25	22.5	4.51	14	26.1	6.98	11	19.2	5.79
65-69	31	27.6	4.97	21	40.5	8.84	10	16.6	5.25
70-74	59	54.5	7.09	28	59.5	11.24	31	50.6	9.09
75-79	74	77.0	8.95	40	101.2	16.00	34	60.1	10.31
80-84	95	140.7	14.43	38	151.4	24.56	57	134.3	17.79
85+	143	242.4	20.27	49	295.0	42.14	94	221.8	22.87
Total deaths	604			313			291		
Connecticut									
. Crude rate		21.0	-		22.3	-		19.7	-
. AAMR (1940 Std. mil.)		10.7	0.52		13.9	0.87		7.7	0.60
. AAMR (1970 Std. mil.)		13.3	0.58		17.0	0.98		10.2	0.69
. AAMR (2000 Std. mil.)		17.9	0.73		22.7	1.29		14.0	0.85
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.8	-		16.7	-		9.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----HIV infection (042-044)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.53	0	0.0	0.00	1	1.1	1.08
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.3	0.93	1	1.3	1.29	1	1.4	1.35
25-29	5	2.8	1.26	5	5.6	2.50	0	0.0	0.00
30-34	14	6.2	1.66	10	8.9	2.82	4	3.5	1.77
35-39	18	7.0	1.65	12	9.4	2.71	6	4.7	1.90
40-44	22	8.9	1.90	18	14.8	3.49	4	3.2	1.59
45-49	16	7.7	1.93	13	12.7	3.53	3	2.9	1.65
50-54	7	3.8	1.44	7	7.8	2.93	0	0.0	0.00
55-59	10	7.1	2.23	8	11.6	4.09	2	2.8	1.95
60-64	1	0.9	0.90	1	1.9	1.87	0	0.0	0.00
65-69	1	0.9	0.89	1	1.9	1.93	0	0.0	0.00
70-74	1	0.9	0.92	1	2.1	2.12	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	99			77			22		
Connecticut									
. Crude rate		3.4	-		5.5	-		1.5	-
. AAMR (1940 Std. mil.)		3.1	0.32		4.9	0.56		1.4	0.32
. AAMR (1970 Std. mil.)		2.8	0.29		4.3	0.50		1.3	0.29
. AAMR (2000 Std. mil.)		3.3	0.33		5.2	0.60		1.5	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.6	-		7.2	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All cancer (140-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	5	2.8	1.25	3	3.3	1.88	2	2.3	1.62
5-9	7	3.6	1.36	2	2.0	1.41	5	5.3	2.35
10-14	3	1.6	0.92	2	2.1	1.47	1	1.1	1.08
15-19	2	1.2	0.82	1	1.1	1.12	1	1.2	1.19
20-24	9	5.9	1.98	6	7.7	3.16	3	4.1	2.34
25-29	8	4.5	1.59	4	4.5	2.24	4	4.5	2.26
30-34	30	13.3	2.43	15	13.4	3.45	15	13.3	3.43
35-39	56	21.8	2.91	28	21.9	4.13	28	21.7	4.11
40-44	106	42.9	4.17	52	42.8	5.94	54	43.0	5.86
45-49	180	86.9	6.48	79	77.4	8.70	101	96.2	9.57
50-54	311	168.8	9.57	141	156.2	13.16	170	180.9	13.87
55-59	381	268.8	13.77	201	290.6	20.50	180	248.0	18.49
60-64	496	447.4	20.09	262	488.8	30.20	234	408.6	26.71
65-69	760	677.8	24.59	432	833.4	40.10	328	544.0	30.04
70-74	1,019	940.5	29.46	521	1106.6	48.48	498	812.9	36.43
75-79	1,175	1222.6	35.67	600	1518.0	61.97	575	1016.2	42.38
80-84	1,019	1508.7	47.26	524	2087.6	91.20	495	1166.4	52.42
85+	1,013	1716.9	53.95	403	2426.1	120.85	610	1439.1	58.27
Total deaths	6,580			3,276			3,304		
Connecticut									
. Crude rate		228.3	-		233.7	-		223.2	-
. AAMR (1940 Std. mil.)		115.3	1.64		133.0	2.54		102.7	2.15
. AAMR (1970 Std. mil.)		154.0	2.01		182.8	3.26		134.6	2.56
. AAMR (2000 Std. mil.)		194.5	2.41		235.6	4.14		168.7	3.00
. HP 2000 (1940 Std. mil.)		120.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		123.5	-		147.7	-		105.5	-
. HP 2000 (1940 Std. mil.)		130.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Oral cavity or pharynx (140.0-149.9)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.44	1	0.9	0.89	0	0.0	0.00
35-39	4	1.6	0.78	4	3.1	1.56	0	0.0	0.00
40-44	1	0.4	0.40	0	0.0	0.00	1	0.8	0.80
45-49	1	0.5	0.48	1	1.0	0.98	0	0.0	0.00
50-54	4	2.2	1.09	3	3.3	1.92	1	1.1	1.06
55-59	10	7.1	2.23	6	8.7	3.54	4	5.5	2.76
60-64	4	3.6	1.80	2	3.7	2.64	2	3.5	2.47
65-69	13	11.6	3.22	9	17.4	5.79	4	6.6	3.32
70-74	5	4.6	2.06	2	4.2	3.00	3	4.9	2.83
75-79	18	18.7	4.41	10	25.3	8.00	8	14.1	5.00
80-84	12	17.8	5.13	7	27.9	10.54	5	11.8	5.27
85+	8	13.6	4.79	2	12.0	8.51	6	14.2	5.78
Total deaths	81			47			34		
Connecticut									
. Crude rate		2.8	-		3.4	-		2.3	-
. AAMR (1940 Std. mil.)		1.5	0.19		2.1	0.33		1.0	0.21
. AAMR (1970 Std. mil.)		2.0	0.23		2.6	0.39		1.4	0.26
. AAMR (2000 Std. mil.)		2.4	0.27		3.3	0.48		1.7	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.0	-		3.1	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Colorectal cancer (153.0-154.3,154.8,159.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	3	1.3	0.77	1	0.9	0.89	2	1.8	1.25
35-39	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
40-44	11	4.5	1.34	5	4.1	1.84	6	4.8	1.95
45-49	9	4.3	1.45	3	2.9	1.70	6	5.7	2.33
50-54	19	10.3	2.37	13	14.4	3.99	6	6.4	2.61
55-59	30	21.2	3.86	17	24.6	5.96	13	17.9	4.97
60-64	42	37.9	5.85	19	35.4	8.13	23	40.2	8.37
65-69	81	72.2	8.03	44	84.9	12.80	37	61.4	10.09
70-74	92	84.9	8.85	51	108.3	15.17	41	66.9	10.45
75-79	133	138.4	12.00	60	151.8	19.60	73	129.0	15.10
80-84	136	201.4	17.27	62	247.0	31.37	74	174.4	20.27
85+	150	254.2	20.76	45	270.9	40.38	105	247.7	24.17
Total deaths	707			321			386		
Connecticut									
. Crude rate		24.5	-		22.9	-		26.1	-
. AAMR (1940 Std. mil.)		10.8	0.48		12.2	0.75		9.7	0.62
. AAMR (1970 Std. mil.)		15.4	0.61		17.5	0.99		13.8	0.78
. AAMR (2000 Std. mil.)		20.4	0.77		23.2	1.30		18.4	0.96
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		12.0	-		14.4	-		10.0	-
. HP 2000 (1940 Std. mil.)		13.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pancreatic cancer (157)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.6	0.56	0	0.0	0.00	1	1.1	1.13
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
40-44	4	1.6	0.81	3	2.5	1.43	1	0.8	0.80
45-49	12	5.8	1.67	6	5.9	2.40	6	5.7	2.33
50-54	17	9.2	2.24	7	7.8	2.93	10	10.6	3.36
55-59	20	14.1	3.16	12	17.3	5.01	8	11.0	3.90
60-64	23	20.7	4.33	17	31.7	7.69	6	10.5	4.28
65-69	49	43.7	6.24	31	59.8	10.74	18	29.9	7.04
70-74	63	58.1	7.33	29	61.6	11.44	34	55.5	9.52
75-79	66	68.7	8.45	28	70.8	13.39	38	67.2	10.89
80-84	68	100.7	12.21	36	143.4	23.90	32	75.4	13.33
85+	65	110.2	13.66	19	114.4	26.24	46	108.5	16.00
Total deaths	389			189			200		
Connecticut									
. Crude rate		13.5	-		13.5	-		13.5	-
. AAMR (1940 Std. mil.)		6.5	0.38		7.8	0.62		5.4	0.47
. AAMR (1970 Std. mil.)		8.9	0.48		10.7	0.79		7.5	0.58
. AAMR (2000 Std. mil.)		11.4	0.58		13.5	0.99		9.8	0.71
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.2	-		7.3	-		5.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Malignant melanoma of skin (172)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	2	0.9	0.63	1	0.9	0.89	1	0.9	0.89
35-39	2	0.8	0.55	0	0.0	0.00	2	1.6	1.10
40-44	3	1.2	0.70	3	2.5	1.43	0	0.0	0.00
45-49	3	1.4	0.84	3	2.9	1.70	0	0.0	0.00
50-54	7	3.8	1.44	5	5.5	2.48	2	2.1	1.50
55-59	7	4.9	1.87	3	4.3	2.50	4	5.5	2.76
60-64	5	4.5	2.02	3	5.6	3.23	2	3.5	2.47
65-69	10	8.9	2.82	8	15.4	5.46	2	3.3	2.35
70-74	13	12.0	3.33	7	14.9	5.62	6	9.8	4.00
75-79	11	11.4	3.45	6	15.2	6.20	5	8.8	3.95
80-84	11	16.3	4.91	3	12.0	6.90	8	18.9	6.66
85+	12	20.3	5.87	5	30.1	13.46	7	16.5	6.24
Total deaths	86			47			39		
Connecticut									
. Crude rate		3.0	-		3.4	-		2.6	-
. AAMR (1940 Std. mil.)		1.7	0.20		2.2	0.34		1.2	0.24
. AAMR (1970 Std. mil.)		2.1	0.23		2.7	0.40		1.5	0.27
. AAMR (2000 Std. mil.)		2.6	0.28		3.3	0.49		2.0	0.32
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.9	-		2.7	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Female breast cancer (174)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	2	0.9	0.63	-	-	-	2	1.8	1.25
35-39	10	3.9	1.23	-	-	-	10	7.8	2.46
40-44	15	6.1	1.57	-	-	-	15	12.0	3.09
45-49	31	15.0	2.69	-	-	-	31	29.5	5.30
50-54	47	25.5	3.72	-	-	-	47	50.0	7.29
55-59	31	21.9	3.93	-	-	-	31	42.7	7.67
60-64	43	38.8	5.91	-	-	-	43	75.1	11.45
65-69	55	49.1	6.61	-	-	-	55	91.2	12.30
70-74	89	82.1	8.71	-	-	-	89	145.3	15.40
75-79	57	59.3	7.86	-	-	-	57	100.7	13.34
80-84	65	96.2	11.94	-	-	-	65	153.2	19.00
85+	97	164.4	16.69	-	-	-	97	228.8	23.23
Total deaths	542			-			542		
Connecticut									
. Crude rate		-	-		-	-		36.6	-
. AAMR (1940 Std. mil.)		-	-		-	-		18.8	0.95
. AAMR (1970 Std. mil.)		-	-		-	-		23.4	1.09
. AAMR (2000 Std. mil.)		-	-		-	-		28.7	1.26
. HP 2000 (1940 Std. mil.)		-	-		-	-		23.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		18.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		20.6	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									CT rate is lower
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cervical cancer (180)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	1	0.4	0.44	-	-	-	1	0.9	0.89
35-39	3	1.2	0.67	-	-	-	3	2.3	1.35
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	2	1.0	0.68	-	-	-	2	1.9	1.35
50-54	5	2.7	1.21	-	-	-	5	5.3	2.38
55-59	1	0.7	0.71	-	-	-	1	1.4	1.38
60-64	2	1.8	1.28	-	-	-	2	3.5	2.47
65-69	4	3.6	1.78	-	-	-	4	6.6	3.32
70-74	1	0.9	0.92	-	-	-	1	1.6	1.63
75-79	2	2.1	1.47	-	-	-	2	3.5	2.50
80-84	4	5.9	2.96	-	-	-	4	9.4	4.71
85+	5	8.5	3.79	-	-	-	5	11.8	5.28
Total deaths	30			-			30		
Connecticut									
. Crude rate		-	-		-	-		2.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.25
. AAMR (1970 Std. mil.)		-	-		-	-		1.3	0.26
. AAMR (2000 Std. mil.)		-	-		-	-		1.7	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.1	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		2.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		1.3	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Endometrial cancer (182)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	0	0.0	0.00	-	-	-	0	0.0	0.00
40-44	0	0.0	0.00	-	-	-	0	0.0	0.00
45-49	2	1.0	0.68	-	-	-	2	1.9	1.35
50-54	1	0.5	0.54	-	-	-	1	1.1	1.06
55-59	4	2.8	1.41	-	-	-	4	5.5	2.76
60-64	4	3.6	1.80	-	-	-	4	7.0	3.49
65-69	2	1.8	1.26	-	-	-	2	3.3	2.35
70-74	3	2.8	1.60	-	-	-	3	4.9	2.83
75-79	7	7.3	2.75	-	-	-	7	12.4	4.68
80-84	10	14.8	4.68	-	-	-	10	23.6	7.45
85+	8	13.6	4.79	-	-	-	8	18.9	6.67
Total deaths	41			-			41		
Connecticut									
. Crude rate		-	-		-	-		2.8	-
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	0.23
. AAMR (1970 Std. mil.)		-	-		-	-		1.6	0.28
. AAMR (2000 Std. mil.)		-	-		-	-		2.1	0.33
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		1.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is not significantly different
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Ovarian cancer (183)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	-	-	-	0	0.0	0.00
5-9	0	0.0	0.00	-	-	-	0	0.0	0.00
10-14	0	0.0	0.00	-	-	-	0	0.0	0.00
15-19	0	0.0	0.00	-	-	-	0	0.0	0.00
20-24	0	0.0	0.00	-	-	-	0	0.0	0.00
25-29	0	0.0	0.00	-	-	-	0	0.0	0.00
30-34	0	0.0	0.00	-	-	-	0	0.0	0.00
35-39	2	0.8	0.55	-	-	-	2	1.6	1.10
40-44	4	1.6	0.81	-	-	-	4	3.2	1.59
45-49	6	2.9	1.18	-	-	-	6	5.7	2.33
50-54	17	9.2	2.24	-	-	-	17	18.1	4.39
55-59	13	9.2	2.54	-	-	-	13	17.9	4.97
60-64	15	13.5	3.49	-	-	-	15	26.2	6.76
65-69	28	25.0	4.72	-	-	-	28	46.4	8.78
70-74	29	26.8	4.97	-	-	-	29	47.3	8.79
75-79	26	27.1	5.31	-	-	-	26	46.0	9.01
80-84	21	31.1	6.78	-	-	-	21	49.5	10.80
85+	31	52.5	9.44	-	-	-	31	73.1	13.13
Total deaths	192			-			192		
Connecticut									
. Crude rate		-	-		-	-		13.0	-
. AAMR (1940 Std. mil.)		-	-		-	-		6.7	0.56
. AAMR (1970 Std. mil.)		-	-		-	-		8.4	0.66
. AAMR (2000 Std. mil.)		-	-		-	-		10.2	0.75
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		5.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Bladder cancer (188)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	1	0.4	0.40	0	0.0	0.00	1	0.8	0.80
45-49	1	0.5	0.48	1	1.0	0.98	0	0.0	0.00
50-54	4	2.2	1.09	4	4.4	2.22	0	0.0	0.00
55-59	3	2.1	1.22	3	4.3	2.50	0	0.0	0.00
60-64	8	7.2	2.55	8	14.9	5.28	0	0.0	0.00
65-69	16	14.3	3.57	12	23.1	6.68	4	6.6	3.32
70-74	24	22.2	4.52	12	25.5	7.36	12	19.6	5.65
75-79	44	45.8	6.90	33	83.5	14.53	11	19.4	5.86
80-84	38	56.3	9.13	29	115.5	21.45	9	21.2	7.07
85+	41	69.5	10.85	23	138.5	28.87	18	42.5	10.01
Total deaths	180			125			55		
Connecticut									
. Crude rate		6.2	-		8.9	-		3.7	-
. AAMR (1940 Std. mil.)		2.4	0.21		4.2	0.42		1.1	0.18
. AAMR (1970 Std. mil.)		3.7	0.29		6.5	0.60		1.7	0.25
. AAMR (2000 Std. mil.)		5.1	0.38		9.2	0.83		2.5	0.34
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.2	-		3.6	-		1.1	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Brain & other CNS cancer (191-192)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	1	1.1	1.09	0	0.0	0.00
5-9	1	0.5	0.51	1	1.0	1.00	0	0.0	0.00
10-14	2	1.1	0.75	1	1.0	1.04	1	1.1	1.08
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	1	0.7	0.66	1	1.3	1.29	0	0.0	0.00
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	2	0.9	0.63	1	0.9	0.89	1	0.9	0.89
35-39	6	2.3	0.95	4	3.1	1.56	2	1.6	1.10
40-44	5	2.0	0.91	5	4.1	1.84	0	0.0	0.00
45-49	10	4.8	1.53	6	5.9	2.40	4	3.8	1.90
50-54	13	7.1	1.96	7	7.8	2.93	6	6.4	2.61
55-59	14	9.9	2.64	8	11.6	4.09	6	8.3	3.38
60-64	11	9.9	2.99	8	14.9	5.28	3	5.2	3.02
65-69	18	16.1	3.78	13	25.1	6.96	5	8.3	3.71
70-74	18	16.6	3.92	2	4.2	3.00	16	26.1	6.53
75-79	16	16.6	4.16	11	27.8	8.39	5	8.8	3.95
80-84	12	17.8	5.13	6	23.9	9.76	6	14.1	5.77
85+	12	20.3	5.87	6	36.1	14.75	6	14.2	5.78
Total deaths	143			82			61		
Connecticut									
. Crude rate		5.0	-		5.9	-		4.1	-
. AAMR (1940 Std. mil.)		3.3	0.30		4.2	0.50		2.4	0.35
. AAMR (1970 Std. mil.)		3.8	0.34		5.0	0.56		2.9	0.39
. AAMR (2000 Std. mil.)		4.5	0.38		5.8	0.65		3.4	0.44
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		3.5	-		4.3	-		2.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Leukemia (204-208)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.79	0	0.0	0.00	2	2.3	1.62
5-9	4	2.0	1.02	0	0.0	0.00	4	4.2	2.10
10-14	1	0.5	0.53	1	1.0	1.04	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.3	0.93	2	2.6	1.82	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	6	2.7	1.09	3	2.7	1.54	3	2.7	1.54
35-39	6	2.3	0.95	4	3.1	1.56	2	1.6	1.10
40-44	9	3.6	1.21	3	2.5	1.43	6	4.8	1.95
45-49	3	1.4	0.84	0	0.0	0.00	3	2.9	1.65
50-54	12	6.5	1.88	6	6.6	2.71	6	6.4	2.61
55-59	11	7.8	2.34	6	8.7	3.54	5	6.9	3.08
60-64	21	18.9	4.13	11	20.5	6.19	10	17.5	5.52
65-69	15	13.4	3.45	11	21.2	6.40	4	6.6	3.32
70-74	40	36.9	5.84	27	57.3	11.04	13	21.2	5.89
75-79	55	57.2	7.72	38	96.1	15.60	17	30.0	7.29
80-84	34	50.3	8.63	15	59.8	15.43	19	44.8	10.27
85+	31	52.5	9.44	12	72.2	20.85	19	44.8	10.28
Total deaths	252			139			113		
Connecticut									
. Crude rate		8.7	-		9.9	-		7.6	-
. AAMR (1940 Std. mil.)		4.7	0.34		5.8	0.54		4.0	0.46
. AAMR (1970 Std. mil.)		6.1	0.41		7.7	0.67		4.9	0.52
. AAMR (2000 Std. mil.)		7.5	0.48		9.9	0.84		6.0	0.58
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.6	-		5.9	-		3.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes mellitus (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	1	0.4	0.44	0	0.0	0.00	1	0.9	0.89
35-39	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
40-44	7	2.8	1.07	3	2.5	1.43	4	3.2	1.59
45-49	9	4.3	1.45	4	3.9	1.96	5	4.8	2.13
50-54	19	10.3	2.37	12	13.3	3.84	7	7.4	2.81
55-59	23	16.2	3.38	15	21.7	5.60	8	11.0	3.90
60-64	42	37.9	5.85	16	29.8	7.46	26	45.4	8.90
65-69	59	52.6	6.85	32	61.7	10.91	27	44.8	8.62
70-74	76	70.1	8.05	41	87.1	13.60	35	57.1	9.66
75-79	96	99.9	10.19	49	124.0	17.71	47	83.1	12.12
80-84	99	146.6	14.73	44	175.3	26.43	55	129.6	17.47
85+	148	250.8	20.62	46	276.9	40.83	102	240.6	23.83
Total deaths	581			264			317		
Connecticut									
. Crude rate		20.2	-		18.8	-		21.4	-
. AAMR (1940 Std. mil.)		8.8	0.43		10.0	0.67		7.9	0.57
. AAMR (1970 Std. mil.)		12.5	0.56		14.3	0.90		11.3	0.71
. AAMR (2000 Std. mil.)		16.8	0.70		19.3	1.20		15.1	0.87
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.6	-		15.2	-		12.3	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Diabetes-related (250)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	1	1.1	1.09	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.3	0.93	1	1.3	1.29	1	1.4	1.35
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	1	0.4	0.44	0	0.0	0.00	1	0.9	0.89
35-39	6	2.3	0.95	4	3.1	1.56	2	1.6	1.10
40-44	19	7.7	1.77	13	10.7	2.97	6	4.8	1.95
45-49	28	13.5	2.56	18	17.6	4.15	10	9.5	3.01
50-54	51	27.7	3.88	33	36.6	6.36	18	19.2	4.51
55-59	97	68.4	6.95	65	94.0	11.66	32	44.1	7.79
60-64	143	129.0	10.79	84	156.7	17.10	59	103.0	13.41
65-69	208	185.5	12.86	113	218.0	20.51	95	157.6	16.17
70-74	323	298.1	16.59	183	388.7	28.73	140	228.5	19.31
75-79	437	454.7	21.75	221	559.1	37.61	216	381.7	25.97
80-84	443	655.9	31.16	216	860.5	58.55	227	534.9	35.50
85+	566	959.3	40.32	178	1071.6	80.32	388	915.3	46.47
Total deaths	2,326			1,131			1,195		
Connecticut									
. Crude rate		80.7	-		80.7	-		80.7	-
. AAMR (1940 Std. mil.)		33.8	0.83		42.4	1.39		26.9	1.00
. AAMR (1970 Std. mil.)		49.4	1.09		61.5	1.87		40.2	1.30
. AAMR (2000 Std. mil.)		66.7	1.39		82.4	2.47		55.4	1.64
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		-	-		-	-		-	-
. HP 2000 (1940 Std. mil.)		34.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): No comparison is available
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Coronary heart disease (CHD) (402,410-414,429.2)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.12	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	3	1.7	0.97	3	3.4	1.94	0	0.0	0.00
30-34	8	3.6	1.26	6	5.3	2.18	2	1.8	1.25
35-39	20	7.8	1.74	14	10.9	2.92	6	4.7	1.90
40-44	46	18.6	2.75	38	31.3	5.08	8	6.4	2.25
45-49	82	39.6	4.37	67	65.6	8.02	15	14.3	3.69
50-54	131	71.1	6.21	104	115.2	11.30	27	28.7	5.53
55-59	211	148.9	10.25	161	232.8	18.34	50	68.9	9.74
60-64	262	236.3	14.60	193	360.0	25.92	69	120.5	14.50
65-69	431	384.4	18.51	283	545.9	32.45	148	245.5	20.18
70-74	619	571.3	22.96	391	830.4	42.00	228	372.2	24.65
75-79	905	941.6	31.30	489	1237.2	55.95	416	735.2	36.05
80-84	1,177	1742.6	50.79	548	2183.2	93.26	629	1482.1	59.09
85+	2,499	4235.6	84.73	740	4454.9	163.76	1,759	4149.7	98.94
Total deaths	6,395			3,038			3,357		
Connecticut									
. Crude rate		221.9	-		216.7	-		226.8	-
. AAMR (1940 Std. mil.)		78.7	1.22		108.6	2.20		53.7	1.26
. AAMR (1970 Std. mil.)		122.8	1.65		162.0	3.00		91.4	1.77
. AAMR (2000 Std. mil.)		179.8	2.26		227.9	4.17		142.8	2.51
. HP 2000 (1940 Std. mil.)		84.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		96.8	-		131.8	-		68.0	-
. HP 2000 (1940 Std. mil.)		100.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:

- . The CT HP 2000 target: CT rate is lower
- . The U.S. AAMR (1940 Std. mil.): CT rate is lower
- . THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Congestive heart failure (CHF) (428.0)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
40-44	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
45-49	1	0.5	0.48	0	0.0	0.00	1	1.0	0.95
50-54	4	2.2	1.09	2	2.2	1.57	2	2.1	1.50
55-59	7	4.9	1.87	3	4.3	2.50	4	5.5	2.76
60-64	11	9.9	2.99	5	9.3	4.17	6	10.5	4.28
65-69	14	12.5	3.34	10	19.3	6.10	4	6.6	3.32
70-74	38	35.1	5.69	18	38.2	9.01	20	32.6	7.30
75-79	61	63.5	8.13	36	91.1	15.18	25	44.2	8.84
80-84	115	170.3	15.88	47	187.2	27.31	68	160.2	19.43
85+	318	539.0	30.22	72	433.4	51.08	246	580.3	37.00
Total deaths	570			194			376		
Connecticut									
. Crude rate		19.8	-		13.8	-		25.4	-
. AAMR (1940 Std. mil.)		5.0	0.26		5.3	0.43		4.7	0.34
. AAMR (1970 Std. mil.)		9.3	0.41		9.5	0.69		8.9	0.51
. AAMR (2000 Std. mil.)		15.5	0.65		15.2	1.10		15.2	0.80
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		6.8	-		5.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Hypertension with or without renal disease (401,403)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
40-44	3	1.2	0.70	3	2.5	1.43	0	0.0	0.00
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	6	3.3	1.33	3	3.3	1.92	3	3.2	1.84
55-59	4	2.8	1.41	3	4.3	2.50	1	1.4	1.38
60-64	6	5.4	2.21	2	3.7	2.64	4	7.0	3.49
65-69	6	5.4	2.18	4	7.7	3.86	2	3.3	2.35
70-74	12	11.1	3.20	6	12.7	5.20	6	9.8	4.00
75-79	22	22.9	4.88	10	25.3	8.00	12	21.2	6.12
80-84	24	35.5	7.25	11	43.8	13.21	13	30.6	8.50
85+	63	106.8	13.45	15	90.3	23.32	48	113.2	16.34
Total deaths	146			57			89		
Connecticut									
. Crude rate		5.1	-		4.1	-		6.0	-
. AAMR (1940 Std. mil.)		1.7	0.18		1.9	0.29		1.5	0.22
. AAMR (1970 Std. mil.)		2.7	0.24		2.9	0.40		2.5	0.30
. AAMR (2000 Std. mil.)		4.1	0.34		4.3	0.58		3.8	0.42
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.4	-		2.5	-		2.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Cerebrovascular disease (430-438)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.12	0	0.0	0.00
20-24	1	0.7	0.66	0	0.0	0.00	1	1.4	1.35
25-29	2	1.1	0.79	0	0.0	0.00	2	2.3	1.60
30-34	4	1.8	0.89	1	0.9	0.89	3	2.7	1.54
35-39	11	4.3	1.29	7	5.5	2.07	4	3.1	1.55
40-44	10	4.0	1.28	4	3.3	1.65	6	4.8	1.95
45-49	17	8.2	1.99	8	7.8	2.77	9	8.6	2.86
50-54	19	10.3	2.37	7	7.8	2.93	12	12.8	3.69
55-59	44	31.0	4.68	29	41.9	7.79	15	20.7	5.34
60-64	37	33.4	5.49	22	41.0	8.75	15	26.2	6.76
65-69	72	64.2	7.57	47	90.7	13.23	25	41.5	8.29
70-74	151	139.4	11.34	70	148.7	17.77	81	132.2	14.69
75-79	270	280.9	17.10	125	316.2	28.29	145	256.3	21.28
80-84	403	596.7	29.72	161	641.4	50.55	242	570.2	36.65
85+	796	1349.2	47.82	195	1173.9	84.07	601	1417.8	57.83
Total deaths	1,839			677			1,162		
Connecticut									
. Crude rate		63.8	-		48.3	-		78.5	-
. AAMR (1940 Std. mil.)		19.6	0.58		21.2	0.91		18.2	0.74
. AAMR (1970 Std. mil.)		32.8	0.81		34.3	1.34		31.0	1.02
. AAMR (2000 Std. mil.)		50.8	1.19		51.5	2.00		49.3	1.47
. HP 2000 (1940 Std. mil.)		16.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		25.1	-		26.6	-		23.6	-
. HP 2000 (1940 Std. mil.)		20.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is higher
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Pneumonia & influenza (480-487)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.79	1	1.1	1.09	1	1.1	1.14
5-9	1	0.5	0.51	0	0.0	0.00	1	1.1	1.05
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	1	0.4	0.44	1	0.9	0.89	0	0.0	0.00
35-39	3	1.2	0.67	2	1.6	1.10	1	0.8	0.78
40-44	5	2.0	0.91	3	2.5	1.43	2	1.6	1.13
45-49	9	4.3	1.45	3	2.9	1.70	6	5.7	2.33
50-54	8	4.3	1.54	7	7.8	2.93	1	1.1	1.06
55-59	12	8.5	2.44	8	11.6	4.09	4	5.5	2.76
60-64	17	15.3	3.72	10	18.7	5.90	7	12.2	4.62
65-69	37	33.0	5.42	22	42.4	9.05	15	24.9	6.42
70-74	69	63.7	7.67	39	82.8	13.26	30	49.0	8.94
75-79	157	163.4	13.04	77	194.8	22.20	80	141.4	15.81
80-84	235	347.9	22.70	118	470.1	43.28	117	275.7	25.49
85+	611	1035.6	41.90	219	1318.4	89.09	392	924.8	46.71
Total deaths	1,168			511			657		
Connecticut									
. Crude rate		40.5	-		36.5	-		44.4	-
. AAMR (1940 Std. mil.)		10.8	0.40		13.8	0.69		8.9	0.48
. AAMR (1970 Std. mil.)		19.5	0.61		24.8	1.11		16.3	0.71
. AAMR (2000 Std. mil.)		31.9	0.93		40.7	1.82		27.1	1.07
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		13.2	-		16.3	-		11.0	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Asthma (493)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.53	1	1.0	1.04	0	0.0	0.00
15-19	1	0.6	0.58	0	0.0	0.00	1	1.2	1.19
20-24	2	1.3	0.93	2	2.6	1.82	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.55	1	0.8	0.78	1	0.8	0.78
40-44	5	2.0	0.91	1	0.8	0.82	4	3.2	1.59
45-49	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
50-54	5	2.7	1.21	3	3.3	1.92	2	2.1	1.50
55-59	2	1.4	1.00	1	1.4	1.45	1	1.4	1.38
60-64	4	3.6	1.80	1	1.9	1.87	3	5.2	3.02
65-69	1	0.9	0.89	0	0.0	0.00	1	1.7	1.66
70-74	3	2.8	1.60	0	0.0	0.00	3	4.9	2.83
75-79	7	7.3	2.75	2	5.1	3.58	5	8.8	3.95
80-84	6	8.9	3.63	1	4.0	3.98	5	11.8	5.27
85+	6	10.2	4.15	1	6.0	6.02	5	11.8	5.28
Total deaths	45			14			31		
Connecticut									
. Crude rate		1.6	-		1.0	-		2.1	-
. AAMR (1940 Std. mil.)		1.0	0.18		-	-		1.1	0.24
. AAMR (1970 Std. mil.)		1.2	0.19		-	-		1.3	0.27
. AAMR (2000 Std. mil.)		1.4	0.21		-	-		1.7	0.31
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.4	-		1.2	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Chronic liver disease and cirrhosis (571)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	1	1.1	1.09	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.6	0.56	0	0.0	0.00	1	1.1	1.13
30-34	6	2.7	1.09	5	4.5	1.99	1	0.9	0.89
35-39	9	3.5	1.17	4	3.1	1.56	5	3.9	1.74
40-44	20	8.1	1.81	17	14.0	3.40	3	2.4	1.38
45-49	26	12.6	2.46	16	15.7	3.92	10	9.5	3.01
50-54	22	11.9	2.55	14	15.5	4.15	8	8.5	3.01
55-59	25	17.6	3.53	18	26.0	6.13	7	9.6	3.65
60-64	25	22.5	4.51	19	35.4	8.13	6	10.5	4.28
65-69	28	25.0	4.72	14	27.0	7.22	14	23.2	6.21
70-74	37	34.1	5.61	24	51.0	10.40	13	21.2	5.89
75-79	31	32.3	5.79	8	20.2	7.16	23	40.6	8.48
80-84	27	40.0	7.69	13	51.8	14.36	14	33.0	8.82
85+	9	15.3	5.08	4	24.1	12.04	5	11.8	5.28
Total deaths	267			157			110		
Connecticut									
. Crude rate		9.3	-		11.2	-		7.4	-
. AAMR (1940 Std. mil.)		6.2	0.41		8.2	0.68		4.2	0.46
. AAMR (1970 Std. mil.)		7.1	0.45		9.3	0.76		5.0	0.51
. AAMR (2000 Std. mil.)		8.4	0.51		10.9	0.87		6.1	0.59
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.2	-		10.3	-		4.4	-
. HP 2000 (1940 Std. mil.)		6.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Nephritis, nephrotic syndrome/nephrosis (580-589)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.6	0.56	0	0.0	0.00	1	1.1	1.13
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	1	0.4	0.39	0	0.0	0.00	1	0.8	0.78
40-44	2	0.8	0.57	1	0.8	0.82	1	0.8	0.80
45-49	5	2.4	1.08	3	2.9	1.70	2	1.9	1.35
50-54	8	4.3	1.54	5	5.5	2.48	3	3.2	1.84
55-59	8	5.6	2.00	4	5.8	2.89	4	5.5	2.76
60-64	13	11.7	3.25	7	13.1	4.94	6	10.5	4.28
65-69	17	15.2	3.68	6	11.6	4.73	11	18.2	5.50
70-74	37	34.1	5.61	23	48.8	10.19	14	22.9	6.11
75-79	54	56.2	7.65	24	60.7	12.39	30	53.0	9.68
80-84	69	102.2	12.30	34	135.5	23.23	35	82.5	13.94
85+	115	194.9	18.18	44	264.9	39.93	71	167.5	19.88
Total deaths	331			151			180		
Connecticut									
. Crude rate		11.5	-		10.8	-		12.2	-
. AAMR (1940 Std. mil.)		4.1	0.28		4.8	0.44		3.7	0.37
. AAMR (1970 Std. mil.)		6.4	0.38		7.7	0.64		5.7	0.48
. AAMR (2000 Std. mil.)		9.3	0.51		11.5	0.95		8.1	0.62
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.4	-		5.4	-		3.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Unintentional injuries (E800-E949)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	9	5.0	1.67	7	7.6	2.88	2	2.3	1.62
5-9	6	3.1	1.26	5	5.0	2.23	1	1.1	1.05
10-14	7	3.7	1.40	4	4.1	2.07	3	3.2	1.87
15-19	40	23.1	3.66	33	37.0	6.45	7	8.4	3.16
20-24	49	32.4	4.62	38	49.0	7.95	11	14.9	4.49
25-29	60	33.7	4.35	50	55.9	7.91	10	11.3	3.57
30-34	89	39.5	4.19	64	57.0	7.13	25	22.2	4.43
35-39	84	32.7	3.57	59	46.1	6.00	25	19.4	3.88
40-44	65	26.3	3.26	49	40.4	5.76	16	12.7	3.19
45-49	62	29.9	3.80	43	42.1	6.42	19	18.1	4.15
50-54	30	16.3	2.97	24	26.6	5.43	6	6.4	2.61
55-59	31	21.9	3.93	19	27.5	6.30	12	16.5	4.77
60-64	28	25.3	4.77	17	31.7	7.69	11	19.2	5.79
65-69	40	35.7	5.64	26	50.2	9.84	14	23.2	6.21
70-74	56	51.7	6.91	29	61.6	11.44	27	44.1	8.48
75-79	75	78.0	9.01	40	101.2	16.00	35	61.9	10.46
80-84	97	143.6	14.58	44	175.3	26.43	53	124.9	17.15
85+	172	291.5	22.23	72	433.4	51.08	100	235.9	23.59
Total deaths	1,000			623			377		
Connecticut									
. Crude rate		34.7	-		44.4	-		25.5	-
. AAMR (1940 Std. mil.)		25.3	0.94		36.8	1.64		14.1	0.95
. AAMR (1970 Std. mil.)		26.4	0.91		37.9	1.60		15.8	0.94
. AAMR (2000 Std. mil.)		31.7	1.01		45.3	1.83		20.0	1.07
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		30.1	-		43.0	-		17.8	-
. HP 2000 (1940 Std. mil.)		29.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Motor vehicle accidents (E810-E825)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	4	2.0	1.02	4	4.0	2.00	0	0.0	0.00
10-14	2	1.1	0.75	1	1.0	1.04	1	1.1	1.08
15-19	29	16.8	3.11	23	25.8	5.38	6	7.2	2.92
20-24	31	20.5	3.68	23	29.7	6.19	8	10.8	3.83
25-29	28	15.7	2.97	25	28.0	5.59	3	3.4	1.95
30-34	34	15.1	2.59	23	20.5	4.27	11	9.8	2.94
35-39	27	10.5	2.02	17	13.3	3.22	10	7.8	2.46
40-44	14	5.7	1.52	8	6.6	2.33	6	4.8	1.95
45-49	20	9.7	2.16	14	13.7	3.66	6	5.7	2.33
50-54	10	5.4	1.72	8	8.9	3.13	2	2.1	1.50
55-59	9	6.3	2.12	3	4.3	2.50	6	8.3	3.38
60-64	9	8.1	2.71	5	9.3	4.17	4	7.0	3.49
65-69	13	11.6	3.22	6	11.6	4.73	7	11.6	4.39
70-74	16	14.8	3.69	6	12.7	5.20	10	16.3	5.16
75-79	18	18.7	4.41	9	22.8	7.59	9	15.9	5.30
80-84	20	29.6	6.62	9	35.9	11.95	11	25.9	7.81
85+	10	16.9	5.36	4	24.1	12.04	6	14.2	5.78
Total deaths	295			188			107		
Connecticut									
. Crude rate		10.2	-		13.4	-		7.2	-
. AAMR (1940 Std. mil.)		9.9	0.63		13.8	1.08		5.9	0.66
. AAMR (1970 Std. mil.)		9.5	0.60		13.1	1.01		5.9	0.64
. AAMR (2000 Std. mil.)		10.1	0.59		13.6	1.00		6.6	0.65
. HP 2000 (1940 Std. mil.)		10.8	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.6	-		21.6	-		9.9	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Falls and fall related injuries (E880-E888)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	5	2.2	0.99	4	3.6	1.78	1	0.9	0.89
35-39	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
40-44	3	1.2	0.70	3	2.5	1.43	0	0.0	0.00
45-49	7	3.4	1.28	4	3.9	1.96	3	2.9	1.65
50-54	5	2.7	1.21	4	4.4	2.22	1	1.1	1.06
55-59	3	2.1	1.22	2	2.9	2.04	1	1.4	1.38
60-64	4	3.6	1.80	2	3.7	2.64	2	3.5	2.47
65-69	12	10.7	3.09	9	17.4	5.79	3	5.0	2.87
70-74	8	7.4	2.61	4	8.5	4.25	4	6.5	3.26
75-79	22	22.9	4.88	12	30.4	8.76	10	17.7	5.59
80-84	41	60.7	9.48	17	67.7	16.43	24	56.6	11.54
85+	93	157.6	16.35	33	198.7	34.58	60	141.5	18.27
Total deaths	206			97			109		
Connecticut									
. Crude rate		7.1	-		6.9	-		7.4	-
. AAMR (1940 Std. mil.)		2.4	0.22		3.4	0.39		1.7	0.23
. AAMR (1970 Std. mil.)		3.7	0.28		4.9	0.51		2.9	0.31
. AAMR (2000 Std. mil.)		5.8	0.40		7.5	0.77		4.6	0.45
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.8	-		3.8	-		1.9	-
. HP 2000 (1940 Std. mil.)		2.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is not significantly different

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Drowning (E830,E832,E910)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.79	2	2.2	1.54	0	0.0	0.00
5-9	1	0.5	0.51	1	1.0	1.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.12	0	0.0	0.00
20-24	2	1.3	0.93	2	2.6	1.82	0	0.0	0.00
25-29	4	2.2	1.12	3	3.4	1.94	1	1.1	1.13
30-34	1	0.4	0.44	1	0.9	0.89	0	0.0	0.00
35-39	1	0.4	0.39	1	0.8	0.78	0	0.0	0.00
40-44	1	0.4	0.40	1	0.8	0.82	0	0.0	0.00
45-49	1	0.5	0.48	1	1.0	0.98	0	0.0	0.00
50-54	2	1.1	0.77	2	2.2	1.57	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.9	0.90	0	0.0	0.00	1	1.7	1.75
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	3	3.1	1.80	1	2.5	2.53	2	3.5	2.50
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	1	1.7	1.69	0	0.0	0.00	1	2.4	2.36
Total deaths	21			16			5		
Connecticut									
. Crude rate		0.7	-		1.1	-		0.3	-
. AAMR (1940 Std. mil.)		0.7	0.18		1.3	0.33		-	-
. AAMR (1970 Std. mil.)		0.7	0.17		1.2	0.31		-	-
. AAMR (2000 Std. mil.)		0.7	0.16		1.2	0.29		-	-
. HP 2000 (1940 Std. mil.)		1.0	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.7	-		2.7	-		0.6	-
. HP 2000 (1940 Std. mil.)		1.3	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Residential fires (E890-E899)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	1	0.5	0.51	0	0.0	0.00	1	1.1	1.05
10-14	2	1.1	0.75	2	2.1	1.47	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	2	1.3	0.93	2	2.6	1.82	0	0.0	0.00
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
35-39	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
40-44	2	0.8	0.57	1	0.8	0.82	1	0.8	0.80
45-49	2	1.0	0.68	0	0.0	0.00	2	1.9	1.35
50-54	1	0.5	0.54	0	0.0	0.00	1	1.1	1.06
55-59	2	1.4	1.00	0	0.0	0.00	2	2.8	1.95
60-64	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
65-69	4	3.6	1.78	3	5.8	3.34	1	1.7	1.66
70-74	2	1.8	1.31	0	0.0	0.00	2	3.3	2.31
75-79	1	1.0	1.04	0	0.0	0.00	1	1.8	1.77
80-84	3	4.4	2.56	2	8.0	5.63	1	2.4	2.36
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	25			13			12		
Connecticut									
. Crude rate		0.9	-		0.9	-		0.8	-
. AAMR (1940 Std. mil.)		0.7	0.16		-	-		-	-
. AAMR (1970 Std. mil.)		0.8	0.17		-	-		-	-
. AAMR (2000 Std. mil.)		0.8	0.17		-	-		-	-
. HP 2000 (1940 Std. mil.)		0.5	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		1.0	-		1.3	-		0.7	-
. HP 2000 (1940 Std. mil.)		1.2	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Suicide (E950-E959)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	1	0.5	0.53	0	0.0	0.00	1	1.1	1.08
15-19	10	5.8	1.83	8	9.0	3.17	2	2.4	1.69
20-24	14	9.2	2.47	11	14.2	4.28	3	4.1	2.34
25-29	16	9.0	2.25	14	15.7	4.19	2	2.3	1.60
30-34	30	13.3	2.43	20	17.8	3.98	10	8.9	2.80
35-39	34	13.2	2.27	26	20.3	3.98	8	6.2	2.20
40-44	34	13.8	2.36	26	21.4	4.20	8	6.4	2.25
45-49	19	9.2	2.10	14	13.7	3.66	5	4.8	2.13
50-54	21	11.4	2.49	15	16.6	4.29	6	6.4	2.61
55-59	12	8.5	2.44	10	14.5	4.57	2	2.8	1.95
60-64	6	5.4	2.21	5	9.3	4.17	1	1.7	1.75
65-69	10	8.9	2.82	7	13.5	5.10	3	5.0	2.87
70-74	4	3.7	1.85	3	6.4	3.68	1	1.6	1.63
75-79	12	12.5	3.60	10	25.3	8.00	2	3.5	2.50
80-84	11	16.3	4.91	8	31.9	11.27	3	7.1	4.08
85+	9	15.3	5.08	5	30.1	13.46	4	9.4	4.72
Total deaths	243			182			61		
Connecticut									
. Crude rate		8.4	-		13.0	-		4.1	-
. AAMR (1940 Std. mil.)		7.5	0.52		11.5	0.91		3.6	0.50
. AAMR (1970 Std. mil.)		7.0	0.48		10.9	0.85		3.3	0.46
. AAMR (2000 Std. mil.)		8.1	0.52		12.7	0.95		3.9	0.50
. HP 2000 (1940 Std. mil.)		6.7	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		10.4	-		17.2	-		4.0	-
. HP 2000 (1940 Std. mil.)		10.5	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: CT rate is not significantly different
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Homicide and legal intervention (E960-E978)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	2	1.1	0.79	0	0.0	0.00	2	2.3	1.62
5-9	1	0.5	0.51	1	1.0	1.00	0	0.0	0.00
10-14	2	1.1	0.75	0	0.0	0.00	2	2.2	1.53
15-19	10	5.8	1.83	9	10.1	3.37	1	1.2	1.19
20-24	12	7.9	2.29	11	14.2	4.28	1	1.4	1.35
25-29	10	5.6	1.78	7	7.8	2.96	3	3.4	1.95
30-34	5	2.2	0.99	4	3.6	1.78	1	0.9	0.89
35-39	6	2.3	0.95	4	3.1	1.56	2	1.6	1.10
40-44	10	4.0	1.28	8	6.6	2.33	2	1.6	1.13
45-49	3	1.4	0.84	3	2.9	1.70	0	0.0	0.00
50-54	5	2.7	1.21	4	4.4	2.22	1	1.1	1.06
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	5	4.5	2.02	3	5.6	3.23	2	3.5	2.47
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	1	0.9	0.92	1	2.1	2.12	0	0.0	0.00
75-79	1	1.0	1.04	0	0.0	0.00	1	1.8	1.77
80-84	1	1.5	1.48	1	4.0	3.98	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	74			56			18		
Connecticut									
. Crude rate		2.6	-		4.0	-		1.2	-
. AAMR (1940 Std. mil.)		3.0	0.37		4.6	0.64		1.4	0.34
. AAMR (1970 Std. mil.)		2.8	0.34		4.2	0.59		1.3	0.33
. AAMR (2000 Std. mil.)		2.7	0.32		4.1	0.56		1.3	0.30
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		7.3	-		11.3	-		3.2	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All motor vehicle accidents (E810-E819,E958.5,E988.5)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	4	2.0	1.02	4	4.0	2.00	0	0.0	0.00
10-14	2	1.1	0.75	1	1.0	1.04	1	1.1	1.08
15-19	29	16.8	3.11	23	25.8	5.38	6	7.2	2.92
20-24	30	19.8	3.62	22	28.4	6.05	8	10.8	3.83
25-29	26	14.6	2.86	23	25.7	5.37	3	3.4	1.95
30-34	34	15.1	2.59	23	20.5	4.27	11	9.8	2.94
35-39	26	10.1	1.99	16	12.5	3.12	10	7.8	2.46
40-44	14	5.7	1.52	8	6.6	2.33	6	4.8	1.95
45-49	20	9.7	2.16	14	13.7	3.66	6	5.7	2.33
50-54	10	5.4	1.72	8	8.9	3.13	2	2.1	1.50
55-59	9	6.3	2.12	3	4.3	2.50	6	8.3	3.38
60-64	8	7.2	2.55	5	9.3	4.17	3	5.2	3.02
65-69	14	12.5	3.34	7	13.5	5.10	7	11.6	4.39
70-74	14	12.9	3.45	6	12.7	5.20	8	13.1	4.62
75-79	18	18.7	4.41	9	22.8	7.59	9	15.9	5.30
80-84	19	28.1	6.45	8	31.9	11.27	11	25.9	7.81
85+	10	16.9	5.36	4	24.1	12.04	6	14.2	5.78
Total deaths	288			184			104		
Connecticut									
. Crude rate		10.0	-		13.1	-		7.0	-
. AAMR (1940 Std. mil.)		9.6	0.63		13.5	1.06		5.7	0.65
. AAMR (1970 Std. mil.)		9.3	0.59		12.8	1.00		5.8	0.63
. AAMR (2000 Std. mil.)		9.8	0.59		13.3	0.99		6.4	0.64
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		15.2	-		21.0	-		9.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----All falls (E880-E886,E888,E957,E968.1,E987)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	1	0.6	0.58	1	1.1	1.12	0	0.0	0.00
20-24	1	0.7	0.66	1	1.3	1.29	0	0.0	0.00
25-29	1	0.6	0.56	1	1.1	1.12	0	0.0	0.00
30-34	5	2.2	0.99	4	3.6	1.78	1	0.9	0.89
35-39	2	0.8	0.55	2	1.6	1.10	0	0.0	0.00
40-44	4	1.6	0.81	4	3.3	1.65	0	0.0	0.00
45-49	8	3.9	1.37	5	4.9	2.19	3	2.9	1.65
50-54	5	2.7	1.21	4	4.4	2.22	1	1.1	1.06
55-59	2	1.4	1.00	1	1.4	1.45	1	1.4	1.38
60-64	4	3.6	1.80	2	3.7	2.64	2	3.5	2.47
65-69	11	9.8	2.96	9	17.4	5.79	2	3.3	2.35
70-74	6	5.5	2.26	4	8.5	4.25	2	3.3	2.31
75-79	19	19.8	4.54	12	30.4	8.76	7	12.4	4.68
80-84	28	41.5	7.83	13	51.8	14.36	15	35.3	9.13
85+	50	84.7	11.98	19	114.4	26.24	31	73.1	13.13
Total deaths	147			82			65		
Connecticut									
. Crude rate		5.1	-		5.9	-		4.4	-
. AAMR (1940 Std. mil.)		2.2	0.23		3.3	0.42		1.2	0.21
. AAMR (1970 Std. mil.)		3.0	0.26		4.4	0.49		1.9	0.26
. AAMR (2000 Std. mil.)		4.2	0.35		6.1	0.68		2.9	0.36
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		2.6	-		3.7	-		1.6	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is not significantly different
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

All firearms (E922.0-.3,E922.8-.9,E955.0-.4,E965.0-.4,E970,E985.0-.4)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	8	4.6	1.64	7	7.9	2.97	1	1.2	1.19
20-24	17	11.2	2.72	15	19.3	5.00	2	2.7	1.91
25-29	13	7.3	2.03	12	13.4	3.88	1	1.1	1.13
30-34	12	5.3	1.54	11	9.8	2.95	1	0.9	0.89
35-39	14	5.5	1.46	11	8.6	2.59	3	2.3	1.35
40-44	14	5.7	1.52	12	9.9	2.85	2	1.6	1.13
45-49	4	1.9	0.97	3	2.9	1.70	1	1.0	0.95
50-54	10	5.4	1.72	9	10.0	3.32	1	1.1	1.06
55-59	9	6.3	2.12	8	11.6	4.09	1	1.4	1.38
60-64	6	5.4	2.21	5	9.3	4.17	1	1.7	1.75
65-69	4	3.6	1.78	4	7.7	3.86	0	0.0	0.00
70-74	1	0.9	0.92	1	2.1	2.12	0	0.0	0.00
75-79	4	4.2	2.08	4	10.1	5.06	0	0.0	0.00
80-84	4	5.9	2.96	4	15.9	7.97	0	0.0	0.00
85+	3	5.1	2.94	2	12.0	8.51	1	2.4	2.36
Total deaths	123			108			15		
Connecticut									
. Crude rate		4.3	-		7.7	-		1.0	-
. AAMR (1940 Std. mil.)		4.3	0.42		7.7	0.79		1.0	0.29
. AAMR (1970 Std. mil.)		4.0	0.39		7.3	0.73		1.0	0.27
. AAMR (2000 Std. mil.)		4.3	0.39		7.8	0.75		1.0	0.27
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		11.4	-		19.8	-		3.3	-
. HP 2000 (1940 Std. mil.)		11.6	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: CT rate is lower

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm suicide (E955.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	2	1.2	0.82	2	2.2	1.59	0	0.0	0.00
20-24	6	4.0	1.62	5	6.4	2.88	1	1.4	1.35
25-29	6	3.4	1.38	6	6.7	2.74	0	0.0	0.00
30-34	10	4.4	1.40	9	8.0	2.67	1	0.9	0.89
35-39	10	3.9	1.23	9	7.0	2.34	1	0.8	0.78
40-44	10	4.0	1.28	8	6.6	2.33	2	1.6	1.13
45-49	2	1.0	0.68	1	1.0	0.98	1	1.0	0.95
50-54	7	3.8	1.44	6	6.6	2.71	1	1.1	1.06
55-59	7	4.9	1.87	6	8.7	3.54	1	1.4	1.38
60-64	4	3.6	1.80	4	7.5	3.73	0	0.0	0.00
65-69	4	3.6	1.78	4	7.7	3.86	0	0.0	0.00
70-74	1	0.9	0.92	1	2.1	2.12	0	0.0	0.00
75-79	4	4.2	2.08	4	10.1	5.06	0	0.0	0.00
80-84	4	5.9	2.96	4	15.9	7.97	0	0.0	0.00
85+	3	5.1	2.94	2	12.0	8.51	1	2.4	2.36
Total deaths	80			71			9		
Connecticut									
. Crude rate		2.8	-		5.1	-		0.6	-
. AAMR (1940 Std. mil.)		2.5	0.30		4.5	0.57		-	-
. AAMR (1970 Std. mil.)		2.3	0.28		4.3	0.54		-	-
. AAMR (2000 Std. mil.)		2.7	0.30		5.0	0.59		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.4	-		11.4	-		1.7	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

-----Firearm homicide (E965.0-.4)-----

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	5	2.9	1.29	4	4.5	2.24	1	1.2	1.19
20-24	11	7.3	2.19	10	12.9	4.08	1	1.4	1.35
25-29	7	3.9	1.49	6	6.7	2.74	1	1.1	1.13
30-34	1	0.4	0.44	1	0.9	0.89	0	0.0	0.00
35-39	3	1.2	0.67	2	1.6	1.10	1	0.8	0.78
40-44	3	1.2	0.70	3	2.5	1.43	0	0.0	0.00
45-49	2	1.0	0.68	2	2.0	1.38	0	0.0	0.00
50-54	3	1.6	0.94	3	3.3	1.92	0	0.0	0.00
55-59	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
60-64	1	0.9	0.90	0	0.0	0.00	1	1.7	1.75
65-69	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
70-74	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
75-79	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
80-84	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	36			31			5		
Connecticut									
. Crude rate		1.2	-		2.2	-		0.3	-
. AAMR (1940 Std. mil.)		1.6	0.28		2.8	0.52		-	-
. AAMR (1970 Std. mil.)		1.5	0.26		2.5	0.47		-	-
. AAMR (2000 Std. mil.)		1.4	0.23		2.4	0.43		-	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		4.5	-		7.6	-		1.5	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:									
. The CT HP 2000 target:									No comparison is available
. The U.S. AAMR (1940 Std. mil.):									CT rate is lower
. THE U.S. HP 2000 target:									No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for

Alcohol-induced (291,303,305.0,357.5,425.5,535.3,571.0-.3,790.3,E860)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
20-24	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
25-29	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
30-34	8	3.6	1.26	7	6.2	2.36	1	0.9	0.89
35-39	9	3.5	1.17	8	6.2	2.21	1	0.8	0.78
40-44	20	8.1	1.81	14	11.5	3.08	6	4.8	1.95
45-49	22	10.6	2.26	16	15.7	3.92	6	5.7	2.33
50-54	21	11.4	2.49	13	14.4	3.99	8	8.5	3.01
55-59	17	12.0	2.91	11	15.9	4.80	6	8.3	3.38
60-64	13	11.7	3.25	11	20.5	6.19	2	3.5	2.47
65-69	15	13.4	3.45	7	13.5	5.10	8	13.3	4.69
70-74	11	10.2	3.06	8	17.0	6.01	3	4.9	2.83
75-79	3	3.1	1.80	3	7.6	4.38	0	0.0	0.00
80-84	6	8.9	3.63	5	19.9	8.91	1	2.4	2.36
85+	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
Total deaths	145			103			42		
Connecticut									
. Crude rate		5.0	-		7.3	-		2.8	-
. AAMR (1940 Std. mil.)		4.0	0.34		5.9	0.59		2.3	0.35
. AAMR (1970 Std. mil.)		4.1	0.35		6.0	0.60		2.3	0.37
. AAMR (2000 Std. mil.)		4.7	0.39		7.1	0.70		2.7	0.41
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		6.1	-		9.7	-		2.8	-
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is lower
. THE U.S. HP 2000 target: No comparison is available

Connecticut White Resident Age-Specific Mortality Rates by Sex, 1998

Comparison of Connecticut age-adjusted mortality rates (AAMRS) to
U.S. AAMRs and U.S. Healthy People (HP) 2000 target objectives
for
Drug-induced (292,304,305.2-.9,E850-E858,E950.0-.5,E962.0,E980.0-.5)

Age	Both Sexes			Males			Females		
	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error	No. Of Deaths	Rate per 100,000	Std. Error
0-4	1	0.6	0.56	0	0.0	0.00	1	1.1	1.14
5-9	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
10-14	0	0.0	0.00	0	0.0	0.00	0	0.0	0.00
15-19	7	4.0	1.53	5	5.6	2.51	2	2.4	1.69
20-24	16	10.6	2.64	11	14.2	4.28	5	6.8	3.03
25-29	22	12.4	2.63	18	20.1	4.75	4	4.5	2.26
30-34	46	20.4	3.01	33	29.4	5.12	13	11.5	3.20
35-39	53	20.6	2.84	35	27.3	4.62	18	14.0	3.30
40-44	51	20.7	2.89	39	32.1	5.14	12	9.6	2.76
45-49	32	15.5	2.73	19	18.6	4.27	13	12.4	3.43
50-54	10	5.4	1.72	5	5.5	2.48	5	5.3	2.38
55-59	4	2.8	1.41	3	4.3	2.50	1	1.4	1.38
60-64	3	2.7	1.56	0	0.0	0.00	3	5.2	3.02
65-69	1	0.9	0.89	0	0.0	0.00	1	1.7	1.66
70-74	3	2.8	1.60	2	4.2	3.00	1	1.6	1.63
75-79	3	3.1	1.80	2	5.1	3.58	1	1.8	1.77
80-84	4	5.9	2.96	2	8.0	5.63	2	4.7	3.33
85+	2	3.4	2.40	0	0.0	0.00	2	4.7	3.34
Total deaths	258			174			84		
Connecticut									
. Crude rate		9.0	-		12.4	-		5.7	-
. AAMR (1940 Std. mil.)		8.5	0.56		11.7	0.92		5.3	0.61
. AAMR (1970 Std. mil.)		7.3	0.48		10.0	0.79		4.7	0.54
. AAMR (2000 Std. mil.)		8.7	0.54		11.9	0.90		5.5	0.61
. HP 2000 (1940 Std. mil.)		-	-		-	-		-	-
United States									
. AAMR (1940 Std. mil.)		5.9	-		8.2	-		3.6	-
. HP 2000 (1940 Std. mil.)		3.0	-		-	-		-	-

Comparison of the CT White Resident AAMR (1940 Std. mil.) vs:
. The CT HP 2000 target: No comparison is available
. The U.S. AAMR (1940 Std. mil.): CT rate is higher
. THE U.S. HP 2000 target: CT rate is higher