

Cancer Incidence in Connecticut 1980-1996

**State of Connecticut
Department of Public Health
Connecticut Tumor Registry
410 Capitol Avenue, MS# 13TMR
P.O. Box 340308
Hartford, CT 06134-0308**

October, 1999

CANCER INCIDENCE IN CONNECTICUT, 1980-1996

Connecticut Department of Public Health, 410 Capitol Avenue, Hartford CT 06134-0308.

Supported by Contract N01-CN-67005 between the National Cancer Institute and the Connecticut Department of Public Health.

Anthony P. Polednak, Ph.D.

Epidemiologist

Connecticut Tumor Registry

Connecticut Department of Public Health, 1999

(860) 509-7163

TABLE OF CONTENTS

Introduction	1		
References	3		
Age-standardized rates	5		
Age-specific rates			
All sites combined	7	Uterine Corpus	22
Oral cavity, pharynx	8	Ovary	23
Esophagus	9	Prostate	24
Stomach	10	Testis	25
Colon	11	Bladder	26
Rectum, anus	12	Kidney	27
Liver, intrahepatic ducts	13	Brain	28
Gallbladder, other biliary	14	Thyroid	29
Pancreas	15	Hodgkin's disease	30
Larynx	16	Non-Hodgkin's lymphoma	31
Trachea, bronchus, lung	17	Multiple myeloma	32
Connective tissue	18	Leukemia	33
Melanoma of skin	19	Kaposi's sarcoma	34
Female breast	20	Unknown, ill-defined sites	35
Uterine cervix	21		
Appendix			
ICD-O-2 codes	36		

INTRODUCTION

1. Background and Methods

The Connecticut Tumor Registry was established in 1941 in the Division of Cancer Research of the State Health Department, which had started tumor clinics and registries in local hospitals in 1935-40 (Heston et al., 1986). Reporting to the registry improved over time, as shown by the decreasing proportion of cases ascertained by death certificate only, which fell to 2% by 1965-69, and the increasing proportion of cases with histological confirmation, which reached at least 90% by 1965-69. Reporting of cancer cases from hospitals and pathology laboratories in Connecticut is legally mandated. Since 1973, the registry has been part of the National Cancer Institute's Surveillance, Epidemiology and End Results (SEER) Program, and is part of the State of Connecticut Department of Public Health.

Average annual age-specific and age-standardized incidence rates for cancers diagnosed among Connecticut residents during the time period 1935-39 to 1975-79 were presented in the monograph Forty-Five Years of Cancer Incidence in Connecticut, 1935-79 (Heston et al., 1986). Five-year intervals of time are useful because age-specific incidence rates fluctuate from year to year, based on small numbers of cancers. Roush et al. (1987) provided incidence data from the Connecticut Tumor Registry for 1935-79, in 10-year intervals of time period and age. Flannery et al. (1992) reported Connecticut incidence data for 1935-86, ending with the years 1980-84 and 1985-86. For previous reports, incidence rates for 1980-84 were recalculated on the basis of updated estimates of the population of Connecticut, and rates for 1985-89 and 1990-94 were included. At the time that data were analyzed for the present report, incidence data were considered complete for cases diagnosed as late as 1996. This report includes average annual incidence age-specific cancer rates (per 100,000 population) for 1980-84, 1985-89 and 1990-94, and annual rates for 1995 and 1996.

This report includes all invasive cancers. In situ bladder tumors are also included, as in SEER reports (Miller et al., 1992, 1993), because of problems encountered in distinguishing in situ from invasive bladder cancers. A previous report (Heston et al., 1986) did not include in situ bladder tumors in the rates for all cancers combined.

Tumors in the Registry are now coded according to the second version of the International Classification of Diseases for Oncology (abbreviated as "ICD-O-2"). Groupings of ICD-O-2 codes used in this report (see Appendix) are consistent with those in previous publications from the Connecticut Tumor Registry. Lymphomas are not included with the anatomic sites but are combined into separate categories (i.e., Hodgkin's disease and non-Hodgkin's lymphoma), on the basis of histology ("Morphology") codes in ICD-O-2. Rates are also provided for Kaposi's sarcoma, because of dramatic increases in males associated with the AIDS epidemic. Kaposi's sarcoma was coded to connective tissue prior to 1980, but mainly to non-melanoma skin cancer thereafter; for this report, it has been defined by the appropriate ICD-O morphology code.

In addition to average annual incidence rates by specific age group, an overall cancer incidence rate is useful for comparisons of cancer rates by time period. The age structure of the population changes over time and cancer rates also change with age. Therefore, valid comparison of total cancer rates from different time periods requires age adjustment or "standardization." In this report, as in SEER reports, age-standardization was done by the "direct" method. That is, the age-specific cancer incidence rates for Connecticut residents were multiplied by the corresponding proportions of the total U.S. population in 1970 in 5-year age groups. Average annual age-standardized rates were analyzed for 1980-84, 1985-89, and 1990-94 (Tables 1 and 2). Cancer sites with small numbers of cancers (<25 per year) are not shown separately, but are included under "all sites combined" in Tables 1 and 2.

2. Trends

a. Recent increases in age-standardized incidence rates for all cancer sites combined have been strongly influenced by increases in smoking-related cancers in women, and in cancers (mainly, non-Hodgkin's lymphoma and Kaposi's sarcoma) related to exposure to the human immunodeficiency virus (HIV). However, rates for Kaposi's sarcoma have not been increasing since 1985-89.

b. Rates for skin melanomas, for which intense exposure to ultraviolet radiation is a risk factor, have increased steadily after 1980-84.

- c. For lung cancer, rates have declined after 1985-89 in men but not in women.
- d. Rates for cancers of the colon and rectum have been declining in both sexes after 1980-84, although rates fluctuate somewhat from year to year.
- e. Trends in incidence rates for breast cancer have been attributed largely to changes in screening. For invasive breast cancer in women, substantial increases in incidence rates occurred during the late 1980's, followed by little change since 1990.
- f. For prostate cancer, striking increases in incidence rates occurred during the late 1980's and early 1990's. These increases occurred along with rising use of prostate specific antigen (PSA) as a screening test. In Connecticut, as in certain other cancer registries, the incidence rates for prostate cancer reached a peak in 1992 and then declined for most age groups. This decline in prostate cancer incidence appears to be continuing, but rates remain higher than they were in the 1980's.

REFERENCES AND RELEVANT REPORTS

Flannery JT, Polednak AP, Fine J et al. Cancer Trends and Cancer Prevention Research in Connecticut: A Statistical and Epidemiological Report, 1935-1986. Printed in 1992.

Flannery JT, Polednak AP. Cancer Incidence in Connecticut 1980-91. November, 1994.

Heston JF, Kelly JB, Meigs JW et al. Forty-Five Years of Cancer Incidence in Connecticut: 1935-79. National Cancer Institute (NCI) Monograph No. 70. NIH Publication No. 86-2652. NCI, Bethesda MD, 1986.

Miller BA, Ries LAG, Hankey BF et al. (eds.) Cancer Statistics Review 1973-1990. NIH Publication No. 93-2789. Bethesda MD: National Cancer Institute, 1993.

Miller BA, Ries LAG, Hankey BF et al. (eds.) Cancer Statistics Review 1973-1989. NIH Publication No. 92-2789. Bethesda MD: National Cancer Institute, 1992.

Polednak AP. Trends in cancer incidence in Connecticut, 1935-1991. Cancer, 1994;74:2863-2872.

Polednak AP. Trends in prostate cancer incidence in Connecticut (1988-94), by age and race. Cancer 1997; 79:99-103.

Polednak AP. Trends in cancer incidence in Connecticut. Connecticut Medicine, 1997; 61:211-218.

Table 1. Average Annual Age-Standardized Incidence Rates per 100,000 per Year in Connecticut Residents in 1980-84, 1985-89, 1990-94, 1995 and 1996: Males

Site	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1995 Rate/yr.	1996 Rate/yr
Oral, pharynx	18.1	16.7	15.5	13.8	13.7
Esophagus	7.2	7.8	7.4	7.0	8.8
Stomach	14.2	12.9	12.0	11.8	9.5
Colon	48.7	47.8	42.8	37.6	37.9
Rectum	22.7	22.6	19.1	16.9	18.6
Liver, ducts ^a	3.5	4.3	5.0	5.5	5.3
Gallbladder ^b	2.4	2.4	1.9	2.7	2.0
Pancreas	11.2	11.0	10.4	10.5	10.9
Larynx	10.1	9.0	8.5	8.3	7.4
Lung ^c	85.4	85.2	79.1	77.2	75.2
Connective tissue ^d	2.8	2.6	2.5	2.5	2.5
Melanoma of skin	11.3	14.1	18.2	19.7	20.3
Prostate	71.3	77.3	138.7	142.5	136.4
Testis	4.4	4.2	4.8	4.5	5.3
Bladder	34.7	36.1	35.8	33.5	34.1
Kidney ^e	12.4	13.3	13.8	16.6	15.5
Brain, CNS ^f	7.3	7.7	7.7	7.1	6.8
Thyroid	2.0	2.1	2.7	2.4	3.3
Hodgkin's disease	4.1	4.1	3.9	3.6	3.8
Non-Hodgkin's lymphoma	13.9	16.3	19.1	20.8	18.9
Multiple myeloma	4.6	4.3	5.0	5.5	5.0
Leukemia	14.1	13.3	12.3	13.9	12.1
Kaposi's sarcoma	0.7	2.2	2.5	2.3	1.9
Unknown ^g	14.0	14.3	12.3	12.1	12.8
All	431.9	441.8	493.4	489.3	479.3
No. of cancers	33,074	36,678	43,153	8,690	8,562

^a Liver and intrahepatic bile ducts.

^b Gallbladder and other biliary tract.

^c Trachea, bronchus and lung.

^d ICD-O-2 490-499.

^e Kidney, pelvis and ureter.

^f Brain, meninges and cranial nerves.

^g Other and ill-defined sites (Heston et al., 1986).

Notes: Rates are for invasive cancers only, except for bladder (which includes both in situ and invasive tumors). The total rate and total number of cancers includes rarer sites not shown separately in the table. See Appendix for ICD-O-2 codes.

Age standardization involved using the 1970 standard million population.

Table 2. Average Annual Age-Standardized Incidence Rates per 100,000 per Year in Connecticut Residents in 1980-84, 1985-89, 1990-94, 1995 and 1996: Females

Site	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1995 Rate/yr.	1996 Rate/yr
Oral, pharynx	7.0	6.3	6.2	5.7	6.1
Esophagus	2.1	2.1	2.0	2.1	1.9
Stomach	5.8	5.4	5.5	4.1	4.7
Colon	36.2	33.9	30.3	31.7	26.8
Rectum	13.3	12.4	11.4	11.2	11.2
Liver, ducts	1.5	1.3	1.5	1.6	2.2
Gallbladder	2.3	2.4	2.3	2.1	1.4
Pancreas	8.3	8.4	7.9	8.7	8.2
Larynx	1.8	2.0	1.9	2.1	1.6
Lung	33.1	40.6	45.3	47.8	48.0
Connective tissue	2.1	1.7	1.8	1.6	1.3
Melanoma of skin	8.9	10.7	12.5	13.0	13.9
Breast	97.5	112.4	115.9	117.5	121.7
Cervix	8.1	7.9	7.4	7.1	8.0
Corpus ^a	23.1	20.6	23.5	26.0	23.8
Ovary	14.2	14.4	15.3	14.9	14.4
Bladder	9.1	9.2	9.8	9.6	10.4
Kidney	5.0	5.9	7.0	6.9	7.4
Brain, CNS	5.4	5.3	5.3	5.3	4.9
Thyroid	4.7	4.5	6.0	6.2	6.9
Hodgkin's disease	2.8	3.0	3.3	2.9	3.0
Non-Hodgkin's lymphoma	10.6	12.6	12.9	12.9	14.1
Multiple myeloma	3.3	3.4	3.2	3.4	3.2
Leukemia	8.3	7.8	7.5	8.2	6.9
Unknown	11.3	10.1	9.9	8.6	10.0
All	335.1	353.4	363.2	370.0	372.9
No. of cancers	34,764	38,812	41,300	8,459	8,574

See footnotes to Table 1.

^aCorpus, plus "uterus not otherwise specified."

Includes only invasive cancers, except for in situ bladder tumors.

See Appendix for ICD-O-2 codes.

Age-standardization involved using the 1970 U.S. standard million population.

**AVERAGE ANNUAL AGE-SPECIFIC RATES (PER 100,000 PER YEAR)
BY CANCER SITE**

1. ALL SITES COMBINED (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	18.2	23.1	20.3	122	17.1 20	31.5 36
5-9	11.2	13.5	12.2	68	8.3 10	13.3 16
10-14	12.6	10.6	12.6	65	10.0 11	9.9 11
15-19	22.3	21.8	21.3	108	22.0 22	11.7 12
20-24	37.0	32.7	36.4	212	41.5 41	36.8 35
25-29	38.0	47.4	55.5	366	62.7 73	60.9 69
30-34	57.1	67.0	74.8	548	62.9 89	63.6 88
35-39	82.0	94.0	94.8	651	84.8 122	89.2 130
40-44	137.1	145.5	149.0	905	169.6 212	164.6 212
45-49	259.0	262.6	265.3	1375	274.2 304	265.1 303
50-54	463.1	474.5	482.0	1941	555.6 495	553.7 501
55-59	824.4	827.5	907.6	3092	986.6 668	985.3 677
60-64	1303.0	1348.8	1533.3	5102	1612.0 976	1705.1 1004
65-69	1901.0	1979.1	2431.2	7405	2500.1 1478	2373.9 1399
70-74	2579.0	2693.0	3115.8	7899	3067.8 1583	2917.3 1471
75-79	3346.7	3246.2	3660.5	6710	3315.9 1302	3058.5 1253
80-84	3671.1	3795.0	4020.0	4035	3317.6 766	3611.3 861
85+	3907.8	3863.0	3973.1	2549	3585.0 518	3121.0 484
FEMALES						
0-4	19.3	21.5	16.3	93	20.8 23	24.8 27
5-9	10.5	11.6	10.2	54	6.1 7	7.9 9
10-14	11.7	9.4	13.5	67	16.1 17	11.3 12
15-19	19.1	20.7	20.8	100	28.5 27	19.8 19
20-24	28.6	30.5	35.6	202	31.2 30	32.0 29
25-29	58.7	59.0	57.5	379	78.1 91	45.1 51
30-34	98.4	86.6	94.5	700	96.7 138	84.0 117
35-39	157.8	164.4	165.7	1163	166.1 243	173.6 256
40-44	244.8	262.3	278.4	1777	274.8 360	289.0 388
45-49	415.9	417.9	443.9	2406	449.5 521	483.0 575
50-54	571.3	597.8	624.1	2657	660.9 623	670.8 641
55-59	789.3	825.1	841.1	3072	845.7 612	916.5 672
60-64	1016.4	1110.3	1134.4	4174	1176.7 781	1142.0 735
65-69	1303.4	1404.1	1411.7	5276	1423.7 1012	1400.7 975
70-74	1554.7	1671.1	1703.7	5816	1800.0 1230	1740.7 1170
75-79	1749.1	1843.9	1943.4	5450	1885.0 1103	2004.6 1197
80-84	1996.4	2087.4	2133.3	4126	2082.8 869	2198.9 919
85+	2009.8	2045.4	2055.2	3788	1928.5 772	1871.4 782

2. **ORAL CAVITY AND PHARYNX** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.2	0.2	1	0.0 0	0.0 0
10-14	0.8	0.2	0.4	2	0.9 1	0.0 0
15-19	0.1	0.3	1.2	6	0.0 0	0.0 0
20-24	0.0	0.6	0.4	2	3.0 3	1.1 1
25-29	1.2	0.7	0.3	2	1.7 2	0.0 0
30-34	2.5	2.4	0.7	5	0.0 0	2.2 3
35-39	3.9	3.1	4.8	33	2.8 4	4.8 7
40-44	8.7	8.6	7.4	45	8.0 10	10.1 13
45-49	19.7	21.1	20.5	106	18.9 21	13.1 15
50-54	36.8	34.1	31.0	125	25.8 23	26.5 24
55-59	58.4	48.0	47.0	160	42.8 29	45.1 31
60-64	71.4	68.6	57.4	191	38.0 23	56.0 33
65-69	77.7	73.3	76.8	234	64.3 38	50.9 30
70-74	85.1	78.6	70.6	179	83.3 43	55.5 28
75-79	80.2	82.8	63.3	116	61.1 24	61.0 25
80-84	64.5	56.1	58.8	59	56.3 13	88.1 21
85+	100.3	72.2	71.7	46	34.6 5	58.0 9
FEMALES						
0-4	0.0	0.0	0.4	2	0.9 1	0.0 0
5-9	0.4	0.2	0.2	1	0.0 0	0.0 0
10-14	0.5	0.0	0.0	0	0.0 0	0.0 0
15-19	0.2	0.5	0.2	1	2.1 2	0.0 0
20-24	0.6	0.4	0.5	3	1.0 1	1.1 1
25-29	1.0	0.7	0.9	6	2.6 3	1.8 2
30-34	1.3	1.3	1.6	12	0.0 0	0.0 0
35-39	0.9	2.8	1.7	12	2.1 3	1.4 2
40-44	5.1	2.8	4.9	31	3.1 4	3.0 4
45-49	11.5	7.1	6.1	33	10.4 12	10.1 12
50-54	11.0	11.4	11.5	49	4.2 4	10.5 10
55-59	19.7	15.1	17.5	64	19.4 14	25.9 19
60-64	24.5	28.1	19.6	72	16.6 11	18.6 12
65-69	31.4	27.2	32.4	121	16.9 12	17.2 12
70-74	31.2	29.1	28.4	97	32.2 22	29.8 20
75-79	29.4	28.2	23.5	66	22.2 13	28.5 17
80-84	30.6	25.4	30.0	58	21.6 9	23.9 10
85+	27.9	29.7	31.5	58	35.0 14	33.5 14

3. **ESOPHAGUS** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.2	1	0.0 0	0.0 0
20-24	0.0	0.1	0.0	0	0.0 0	0.0 0
25-29	0.1	0.0	0.0	0	0.0 0	0.0 0
30-34	0.5	0.1	0.4	3	0.0 0	0.7 1
35-39	0.4	0.8	0.7	5	0.7 1	2.1 3
40-44	2.2	2.9	2.6	16	0.8 1	7.0 9
45-49	6.2	7.4	6.0	31	6.3 7	6.1 7
50-54	11.8	13.1	10.4	42	10.1 9	11.1 10
55-59	19.5	19.8	17.9	61	20.7 14	27.7 19
60-64	26.7	29.1	27.7	92	23.1 14	32.3 19
65-69	33.0	44.9	37.1	113	35.5 21	37.3 22
70-74	35.8	46.0	42.2	107	38.8 20	31.7 16
75-79	53.0	35.6	51.3	94	53.5 21	48.8 20
80-84	39.7	47.0	45.8	46	30.3 7	88.1 21
85+	47.3	30.7	34.3	22	48.5 7	64.5 10
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.0	0.0	0.0	0	0.0 0	0.0 0
25-29	0.0	0.0	0.2	1	0.0 0	0.0 0
30-34	0.0	0.0	0.1	1	0.0 0	0.0 0
35-39	0.0	0.2	0.0	0	0.7 1	0.7 1
40-44	0.2	0.5	0.6	4	0.0 0	0.7 1
45-49	2.4	0.9	1.3	7	0.9 1	0.0 0
50-54	3.4	2.6	2.1	9	3.2 3	2.1 2
55-59	7.0	4.8	5.8	21	5.5 4	8.2 6
60-64	6.4	8.2	7.9	29	9.0 6	9.3 6
65-69	8.7	10.1	9.4	35	7.0 5	2.9 2
70-74	9.7	14.7	13.2	45	13.2 9	10.4 7
75-79	16.1	10.0	11.4	32	13.7 8	11.7 7
80-84	11.3	17.1	11.4	22	16.8 7	12.0 5
85+	20.4	17.0	15.2	28	22.5 9	23.9 10

4. **STOMACH** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.1	0.0	0.2	1	0.0 0	0.0 0
25-29	0.3	0.1	0.6	4	0.0 0	0.0 0
30-34	0.3	0.7	0.8	6	0.0 0	0.0 0
35-39	2.1	1.8	0.4	3	0.0 0	0.0 0
40-44	2.5	3.4	4.0	24	4.8 6	3.1 4
45-49	9.7	10.1	7.5	39	7.2 8	7.9 9
50-54	12.5	13.4	13.7	55	16.8 15	8.8 8
55-59	26.5	22.4	22.0	75	16.3 11	17.5 12
60-64	36.1	38.1	34.3	114	31.4 19	30.6 18
65-69	67.7	57.7	53.8	164	50.8 30	32.2 19
70-74	95.0	75.6	82.4	209	83.3 43	61.5 31
75-79	112.0	96.7	76.4	140	101.9 40	90.3 37
80-84	140.3	129.9	111.6	112	95.3 22	100.7 24
85+	160.9	164.0	157.4	101	145.3 21	77.4 12
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.1	0.0	0.0	0	0.0 0	0.0 0
20-24	0.0	0.0	0.0	0	0.0 0	0.0 0
25-29	0.0	0.1	0.5	3	0.0 0	0.9 1
30-34	0.8	0.6	0.7	5	1.4 2	1.4 2
35-39	0.7	0.9	2.0	14	0.0 0	0.7 1
40-44	1.4	2.0	0.8	5	0.8 1	0.7 1
45-49	2.9	2.4	2.2	12	2.6 3	2.5 3
50-54	7.2	6.3	7.3	31	5.3 5	3.1 3
55-59	7.5	7.9	9.6	35	5.5 4	4.1 3
60-64	11.2	13.0	12.8	47	6.0 4	12.4 8
65-69	20.1	17.1	22.2	83	15.5 11	21.6 15
70-74	40.1	33.6	32.5	111	30.7 21	34.2 23
75-79	51.9	50.1	44.9	126	25.6 15	40.2 24
80-84	66.6	70.9	64.1	124	55.1 23	59.8 25
85+	85.1	66.5	71.1	131	67.5 27	50.3 21

5. COLON (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.2	0.0	0.0	0	0.0 0	0.0 0
15-19	0.1	0.3	0.0	0	0.0 0	0.0 0
20-24	0.6	0.4	0.5	3	0.0 0	1.1 1
25-29	0.8	1.0	1.2	8	0.9 1	0.0 0
30-34	1.8	2.0	1.4	10	4.2 6	2.9 4
35-39	5.3	4.0	3.2	22	4.9 7	6.2 9
40-44	8.3	8.7	6.3	38	9.6 12	11.7 15
45-49	18.4	18.5	18.0	93	9.9 11	16.6 19
50-54	34.9	32.9	35.0	141	39.3 35	36.5 33
55-59	76.3	77.0	73.7	251	66.5 45	43.7 30
60-64	139.2	144.0	123.5	411	99.1 60	115.5 68
65-69	218.7	206.9	201.3	613	165.8 98	199.0 117
70-74	313.2	319.6	268.6	681	242.3 125	226.1 114
75-79	467.7	413.4	365.5	670	333.6 131	346.6 142
80-84	518.3	560.4	470.2	472	372.5 86	423.6 101
85+	504.4	573.4	528.4	339	484.5 70	322.4 50
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.2	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.2	1	1.1 1	0.0 0
20-24	0.1	0.3	0.5	3	0.0 0	0.0 0
25-29	0.3	0.6	0.9	6	1.7 2	0.0 0
30-34	2.5	2.9	1.5	11	0.7 1	0.7 1
35-39	5.4	5.0	4.9	34	5.5 8	2.7 4
40-44	7.1	8.0	8.2	52	9.2 12	8.2 11
45-49	19.6	15.3	15.9	86	17.3 20	16.0 19
50-54	37.0	32.9	26.3	191	29.7 28	27.2 26
55-59	57.6	55.9	52.3	191	51.1 37	55.9 41
60-64	95.5	90.6	88.1	324	76.8 51	68.4 44
65-69	146.6	143.7	127.1	475	144.9 103	84.8 59
70-74	233.5	202.4	183.7	627	226.8 155	162.2 109
75-79	325.7	282.0	251.4	705	258.1 151	237.8 142
80-84	414.8	399.2	349.0	675	330.8 138	327.8 137
85+	429.3	444.8	370.6	683	344.7 138	351.8 147

6. **RECTUM** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.2	0.1	1	0.0 0	0.0 0
20-24	0.1	0.0	0.0	0	0.0 0	0.0 0
25-29	1.1	0.3	0.8	5	0.0 0	0.9 1
30-34	0.0	0.9	1.0	7	0.7 1	2.9 4
35-39	1.5	1.9	2.3	16	2.1 3	1.4 2
40-44	6.3	4.4	6.1	37	2.4 3	7.8 10
45-49	7.8	11.2	10.2	53	9.0 10	12.3 14
50-54	28.4	30.0	19.4	78	19.1 17	29.8 27
55-59	43.7	46.0	40.2	137	34.0 23	36.4 25
60-64	80.7	71.7	67.0	223	57.8 35	57.7 34
65-69	115.5	107.7	102.4	312	91.3 54	79.8 47
70-74	142.9	154.4	121.5	308	108.5 56	128.9 65
75-79	168.2	165.6	135.3	248	96.8 38	122.1 50
80-84	177.3	200.4	147.5	148	168.9 39	142.6 34
85+	219.9	178.7	138.7	89	152.3 22	148.3 23
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.1	0.0	0.2	1	0.0 0	0.0 0
20-24	0.3	0.0	0.2	1	0.0 0	0.0 0
25-29	0.0	0.1	0.6	4	1.7 2	0.0 0
30-34	0.6	0.6	1.6	12	0.7 1	2.9 4
35-39	1.8	2.3	1.7	12	2.1 3	0.7 1
40-44	3.1	4.8	3.9	25	6.9 9	2.2 3
45-49	10.1	6.8	7.4	40	7.8 9	13.4 16
50-54	18.6	13.7	18.1	77	15.9 15	12.6 12
55-59	26.8	22.6	24.1	88	20.7 15	27.3 20
60-64	44.6	41.5	42.7	157	28.6 19	46.6 30
65-69	61.4	58.0	47.4	177	60.5 43	51.7 36
70-74	84.2	76.9	65.0	222	67.3 46	58.0 39
75-79	87.3	94.5	76.7	215	61.5 36	63.6 38
80-84	115.0	113.6	86.9	168	105.5 44	81.4 34
85+	117.6	107.4	94.4	174	92.4 37	76.6 32

7. LIVER AND INTRAHEPATIC BILE DUCTS (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	1.0	0.6	0.2	1	0.0	0	0.9	1
5-9	0.0	0.0	0.2	1	0.0	0	0.9	1
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.2	1	0.0	0	0.0	0
20-24	0.1	0.0	0.2	1	0.0	0	0.0	0
25-29	0.0	0.3	0.5	3	0.0	0	0.0	0
30-34	0.2	0.1	1.0	7	0.0	0	0.0	0
35-39	0.3	0.6	0.7	5	1.4	2	0.0	0
40-44	0.4	1.5	2.1	13	2.4	3	1.6	2
45-49	1.3	2.1	4.4	23	6.3	7	3.5	4
50-54	4.2	4.0	6.7	27	6.7	6	8.8	8
55-59	9.9	9.2	11.2	38	10.3	7	10.2	7
60-64	9.3	17.6	10.5	35	19.8	12	27.2	16
65-69	18.0	21.4	26.3	80	27.1	16	18.7	11
70-74	20.4	26.3	30.4	77	46.5	24	21.8	11
75-79	22.8	31.4	32.2	59	25.5	10	29.3	12
80-84	35.4	33.5	38.9	39	8.7	2	29.4	7
85+	21.0	21.3	34.3	22	55.4	8	45.1	7
FEMALES								
0-4	0.6	0.4	0.5	3	0.9	1	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.2	0.2	0.0	0	0.0	0	0.0	0
15-19	0.3	0.3	0.2	1	0.0	0	0.0	0
20-24	0.1	0.0	0.2	1	0.0	0	0.0	0
25-29	0.5	0.0	0.5	3	0.0	0	0.0	0
30-34	0.2	0.0	0.0	0	0.0	0	0.0	0
35-39	0.7	0.2	0.1	1	1.4	2	0.7	1
40-44	0.8	0.9	0.3	2	0.0	0	0.0	0
45-49	0.2	1.1	1.1	6	0.0	0	0.8	1
50-54	1.8	1.0	1.4	6	4.2	4	4.2	4
55-59	2.6	2.5	1.9	7	1.4	1	4.1	3
60-64	4.9	2.4	2.2	8	3.0	2	9.3	6
65-69	3.8	6.0	5.9	22	5.6	4	8.6	6
70-74	8.2	9.0	10.8	37	11.7	8	13.4	9
75-79	12.1	9.5	10.3	29	12.0	7	15.1	9
80-84	9.3	11.1	18.1	35	14.4	6	21.5	9
85+	12.1	11.9	14.7	27	20.0	8	14.4	6

8. GALLBLADDER AND OTHER BILIARY TRACT (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.0	0	0.0	0	0.0	0
20-24	0.1	0.0	0.2	1	0.0	0	0.0	0
25-29	0.0	0.0	0.0	0	0.0	0	0.0	0
30-34	0.0	0.1	0.0	0	0.0	0	0.0	0
35-39	0.2	0.2	0.3	2	0.7	1	1.4	2
40-44	0.2	0.6	0.7	4	0.8	1	0.0	0
45-49	0.5	1.2	0.2	1	0.0	0	1.8	2
50-54	1.0	1.6	2.2	9	1.1	1	0.0	0
55-59	4.8	3.9	1.2	4	4.4	3	4.4	3
60-64	8.0	8.1	4.2	14	11.6	7	5.1	3
65-69	8.1	9.3	8.9	27	8.5	5	11.9	7
70-74	18.0	16.2	10.7	27	11.6	6	7.9	4
75-79	20.5	18.9	18.0	33	28.0	11	22.0	9
80-84	20.5	25.5	23.9	24	26.0	6	8.4	2
85+	45.5	34.0	40.5	26	62.3	9	32.2	5
FEMALES								
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.0	0	0.0	0	0.0	0
20-24	0.0	0.0	0.0	0	0.0	0	0.0	0
25-29	0.0	0.1	0.2	1	0.0	0	0.0	0
30-34	0.0	0.0	0.0	0	0.0	0	0.0	0
35-39	0.0	0.0	0.6	4	0.0	0	0.7	1
40-44	0.4	0.9	0.3	2	0.8	1	0.0	0
45-49	1.7	0.9	0.4	2	0.9	1	0.0	0
50-54	1.6	2.0	2.4	10	0.0	0	3.1	3
55-59	2.2	3.2	4.1	15	1.4	1	1.4	1
60-64	5.8	7.7	6.0	22	6.0	4	7.8	5
65-69	13.2	7.8	7.8	29	12.7	9	2.9	2
70-74	19.7	13.6	16.7	57	16.1	11	13.4	9
75-79	27.0	26.3	17.8	50	22.2	13	6.7	4
80-84	36.0	29.1	29.0	56	16.8	7	4.8	2
85+	48.4	32.7	27.7	51	27.5	11	26.3	11

9. **PANCREAS** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.4	0.0	0.2	1	0.0 0	0.0 0
25-29	0.0	0.1	0.0	0	0.0 0	0.0 0
30-34	0.5	0.3	0.3	2	2.1 3	0.0 0
35-39	1.5	1.6	1.0	7	1.4 2	1.4 2
40-44	3.6	4.7	1.7	10	4.8 6	1.6 2
45-49	6.8	5.3	6.0	31	6.3 7	11.4 13
50-54	11.3	13.1	12.4	50	14.6 13	11.1 10
55-59	22.9	21.4	22.9	78	23.6 16	26.2 18
60-64	33.4	35.3	30.4	101	26.4 16	40.8 24
65-69	46.6	59.9	49.6	151	47.4 287	50.9 30
70-74	70.7	70.4	73.0	185	65.9 34	75.4 38
75-79	96.4	74.2	76.4	140	68.8 27	51.3 21
80-84	107.1	93.3	91.7	92	91.0 21	96.5 23
85+	109.8	90.2	85.7	55	110.7 16	77.4 12
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.0	0.0	0.2	1	0.0 0	0.0 0
25-29	0.1	0.0	0.3	2	0.0 0	0.0 0
30-34	0.5	0.1	0.1	1	0.0 0	0.0 0
35-39	1.0	0.6	1.0	7	0.0 0	0.7 1
40-44	1.2	1.5	1.6	10	1.5 2	1.5 2
45-49	3.7	3.5	3.1	17	6.0 7	7.6 9
50-54	6.7	9.1	6.6	28	8.5 8	6.3 6
55-59	15.4	16.2	14.5	53	20.7 15	19.1 14
60-64	28.2	25.3	24.5	90	37.7 25	17.1 11
65-69	34.1	41.7	39.0	142	23.9 17	28.7 20
70-74	52.9	54.8	51.0	174	52.7 36	59.5 40
75-79	59.9	68.2	65.3	183	64.9 38	97.1 58
80-84	102.1	65.2	70.8	137	83.9 35	62.2 26
85+	95.7	79.0	95.6	178	104.9 42	69.4 29

10. **LARYNX** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.0	0	0.0	0	0.0	0
20-24	0.0	0.0	0.0	0	0.0	0	0.0	0
25-29	0.1	0.0	0.0	0	0.0	0	0.0	0
30-34	0.5	0.3	0.4	3	0.7	1	0.0	0
35-39	0.7	1.4	1.3	9	0.0	0	0.7	1
40-44	4.7	4.5	2.1	13	3.2	4	3.1	4
45-49	10.4	5.3	7.3	38	4.5	5	7.0	8
50-54	22.1	14.2	14.9	60	12.4	11	13.3	12
55-59	27.0	32.0	22.6	77	23.6	16	14.6	10
60-64	41.7	35.6	36.4	121	33.0	20	22.1	13
65-69	56.2	43.3	41.4	126	54.1	32	52.6	31
70-74	46.7	39.1	46.6	118	40.7	21	49.6	25
75-79	43.8	61.3	50.7	93	45.8	18	31.7	13
80-84	49.4	48.2	43.8	44	47.6	11	33.6	8
85+	30.3	23.8	35.9	23	34.6	5	38.7	6
FEMALES								
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.0	0	0.0	0	0.0	0
20-24	0.0	0.0	0.0	0	0.0	0	0.0	0
25-29	0.0	0.0	0.0	0	0.0	0	0.0	0
30-34	0.2	0.0	0.3	2	0.0	0	0.7	1
35-39	0.2	0.0	0.0	0	0.0	0	1.4	2
40-44	1.6	0.7	0.6	4	0.8	1	0.7	1
45-49	2.2	1.8	2.2	12	1.7	2	0.8	1
50-54	3.9	5.0	4.7	20	3.2	3	3.1	3
55-59	6.6	6.8	6.0	22	6.9	5	4.1	3
60-64	6.8	12.7	7.9	29	9.0	6	4.7	3
65-69	9.5	8.6	9.4	35	11.3	8	8.6	6
70-74	8.2	8.3	8.2	28	13.2	9	13.4	9
75-79	6.8	6.3	8.2	23	6.8	4	3.4	2
80-84	6.6	0.6	7.2	14	14.4	6	4.8	2
85+	2.9	2.5	2.7	5	0.0	0	7.2	3

11. TRACHEA, BRONCHUS AND LUNG (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	0.6	0.4	0.0	0	0.0	0	0.0	0
5-9	0.0	0.2	0.0	0	0.0	0	0.0	0
10-14	0.0	0.2	0.0	0	0.0	0	0.9	1
15-19	0.3	0.2	0.2	1	0.0	0	0.0	0
20-24	0.4	0.7	0.0	0	0.0	0	0.0	0
25-29	0.9	0.8	0.5	3	0.9	1	0.0	0
30-34	3.4	2.3	2.1	15	0.7	1	1.5	2
35-39	8.5	9.2	5.0	34	3.5	5	6.2	9
40-44	20.7	20.0	14.8	90	19.2	24	14.0	18
45-49	59.9	52.5	39.9	207	39.7	44	31.5	36
50-54	108.8	109.4	83.7	337	80.8	72	86.2	78
55-59	207.2	199.8	167.0	569	149.2	101	168.8	116
60-64	308.2	318.7	279.8	931	284.1	172	295.5	174
65-69	419.0	429.9	418.9	1276	441.5	261	398.8	235
70-74	550.3	542.2	520.3	1319	501.9	259	499.8	252
75-79	616.8	605.1	618.1	1133	629.0	247	495.5	203
80-84	584.0	555.7	625.7	628	502.4	116	625.0	149
85+	437.5	503.2	497.2	319	470.6	68	399.8	62
FEMALES								
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.3	0.4	0.2	1	0.0	0	0.0	0
20-24	0.6	0.1	0.4	2	0.0	0	0.0	0
25-29	0.2	0.4	0.6	4	0.9	1	0.0	0
30-34	1.4	2.2	2.4	18	1.4	2	0.7	1
35-39	6.6	4.0	6.0	42	6.2	9	6.8	10
40-44	17.8	15.7	15.7	100	13.7	18	12.7	17
45-49	30.0	34.3	39.3	213	36.2	42	31.9	38
50-54	71.9	70.2	75.6	322	71.1	67	72.2	69
55-59	99.6	119.7	118.6	433	117.5	85	118.7	87
60-64	134.1	165.4	175.3	645	192.9	128	177.1	114
65-69	164.3	209.4	236.0	882	250.4	178	258.6	180
70-74	173.5	243.9	262.2	895	275.1	188	312.4	210
75-79	141.9	208.0	288.8	810	340.1	199	301.4	180
80-84	141.2	180.4	226.5	438	251.7	105	311.1	130
85+	86.7	105.5	124.2	229	172.4	69	157.9	66

12. **CONNECTIVE TISSUE** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	0.8	1.4	1.8	11	1.7	2	0.0	0
5-9	0.2	0.2	0.7	4	0.0	0	0.0	0
10-14	0.3	0.4	0.6	3	1.9	2	0.9	1
15-19	1.6	1.0	0.4	2	2.0	2	0.0	0
20-24	1.0	1.8	0.7	4	2.0	2	3.2	3
25-29	1.2	1.3	0.9	6	1.7	2	0.9	1
30-34	1.2	2.3	1.6	12	1.4	2	1.5	2
35-39	2.1	3.7	1.5	10	0.7	1	0.0	0
40-44	3.0	2.8	3.5	21	5.6	7	0.8	1
45-49	2.2	2.3	1.9	10	1.8	2	2.6	3
50-54	4.2	4.0	3.2	13	0.0	0	3.3	3
55-59	5.6	4.8	4.7	16	1.5	1	2.9	2
60-64	6.2	4.1	4.8	16	6.6	4	11.9	7
65-69	6.6	8.8	7.9	24	1.7	1	8.5	5
70-74	9.9	9.5	10.3	26	7.8	4	9.9	5
75-79	15.2	6.9	10.4	19	17.8	7	7.3	3
80-84	15.0	9.8	8.0	8	4.3	1	12.6	3
85+	20.8	30.6	17.2	11	13.8	2	12.9	2
FEMALES								
0-4	1.8	1.1	0.7	4	1.8	2	0.9	1
5-9	0.6	0.4	0.4	2	0.0	0	0.0	0
10-14	0.8	0.0	0.4	2	0.0	0	0.0	0
15-19	1.7	1.4	0.8	4	0.0	0	0.0	0
20-24	1.0	0.8	0.7	4	0.0	0	1.1	1
25-29	0.7	1.0	1.8	12	1.7	2	1.8	2
30-34	2.1	0.8	1.5	11	2.8	4	0.7	1
35-39	1.6	1.1	1.0	7	0.0	0	0.0	0
40-44	1.5	2.3	0.9	6	0.8	1	2.2	3
45-49	2.0	2.0	1.7	9	1.7	2	0.8	1
50-54	2.7	1.3	1.4	6	3.2	3	6.3	6
55-59	3.5	2.6	1.9	7	1.4	1	1.4	1
60-64	3.3	3.2	3.5	13	6.0	4	1.6	1
65-69	4.8	4.0	5.9	22	4.2	3	2.9	2
70-74	7.1	5.0	6.7	23	4.4	3	1.5	1
75-79	3.6	4.7	6.4	18	5.1	3	1.7	1
80-84	9.3	8.3	4.7	9	7.2	3	9.6	4
85+	10.0	12.5	11.4	21	7.5	3	9.6	4

13. MELANOMA OF SKIN (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.2	0.2	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.2	1	0.0 0	0.0 0
15-19	0.0	0.5	1.0	5	2.0 2	2.0 2
20-24	2.5	2.4	2.4	14	1.0 1	3.2 3
25-29	3.7	4.7	3.9	26	4.3 5	10.6 12
30-34	6.9	8.4	9.3	68	5.7 8	5.1 7
35-39	11.3	10.6	13.4	92	11.1 16	9.6 14
40-44	14.5	16.8	18.1	110	28.0 35	17.9 23
45-49	16.1	19.0	22.8	118	27.1 30	23.6 27
50-54	25.7	27.6	31.3	126	37.0 33	29.8 27
55-59	27.3	33.0	45.2	154	47.3 32	50.9 35
60-64	39.6	41.9	57.7	192	64.4 39	51.0 30
65-69	31.9	46.6	60.7	185	57.5 34	100.1 59
70-74	39.7	57.6	81.3	206	79.5 41	79.3 40
75-79	36.5	55.6	85.7	157	91.7 36	102.5 42
80-84	39.6	73.9	84.7	85	99.6 23	121.6 29
85+	39.8	53.6	96.6	62	117.7 17	96.7 15
FEMALES						
0-4	0.2	0.0	0.2	1	0.0 0	0.0 0
5-9	0.2	0.0	0.0	0	0.0 0	0.9 1
10-14	0.0	0.4	0.8	4	1.0 1	0.0 0
15-19	1.3	1.1	1.9	9	2.1 2	2.1 2
20-24	2.7	3.6	3.9	22	4.2 4	1.1 1
25-29	7.0	8.2	8.6	57	10.3 12	5.3 6
30-34	10.8	9.8	13.9	103	9.1 13	11.5 16
35-39	13.5	15.9	16.3	114	19.1 28	18.3 27
40-44	11.3	17.7	21.6	138	22.9 30	26.1 35
45-49	19.9	22.3	22.0	119	26.8 31	21.0 25
50-54	22.2	25.8	21.1	90	22.3 21	20.9 20
55-59	17.4	25.5	30.4	111	33.2 24	34.1 25
60-64	18.8	24.1	27.5	101	22.6 15	42.0 27
65-69	21.2	21.2	28.9	108	35.2 25	41.7 29
70-74	21.3	22.9	33.1	113	36.6 25	43.2 29
75-79	17.9	20.9	34.2	96	34.2 20	38.5 23
80-84	17.8	23.7	38.8	75	28.8 12	55.0 23
85+	24.3	33.3	44.5	82	25.0 10	33.5 14

14. **FEMALE BREAST** (Rates are per 100,000 females per year)

Age	1980-84	1985-89	1990-94	1990-94	1995		1996	
	Rate/yr.	Rate/yr.	Rate/yr.	Total No.	Rate/Yr.	No.	Rate/Yr.	No.
0-4	0.0	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.4	0.0	0.0	0	0.0	0	0.0	0
20-24	0.7	1.3	1.2	7	0.0	0	0.0	0
25-29	10.8	8.2	7.4	49	16.3	19	9.7	11
30-34	32.9	28.6	26.1	193	30.1	43	25.8	36
35-39	67.8	67.7	66.3	465	62.2	91	73.2	108
40-44	109.8	121.7	129.0	823	122.1	160	138.5	186
45-49	183.4	191.5	211.2	1145	209.7	243	246.1	293
50-54	198.2	238.6	251.8	1072	294.9	278	281.5	269
55-59	261.7	295.2	294.1	1074	294.3	213	343.7	252
60-64	291.4	364.0	372.9	1372	375.2	249	368.2	237
65-69	342.8	425.6	422.0	1577	422.0	300	416.6	290
70-74	380.2	447.6	470.2	1605	477.0	326	459.7	309
75-79	421.2	488.6	514.2	1442	485.4	284	525.9	314
80-84	413.0	494.2	516.0	998	505.7	211	504.9	211
85+	426.1	471.2	466.6	860	402.2	161	354.2	148

15. **UTERINE CERVIX (INVASIVE)** (Rates are per 100,000 females per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.2	0.0	0.6	3	0.0 0	2.1 2
20-24	2.0	1.5	1.9	11	0.0 0	6.6 6
25-29	9.8	7.5	8.6	57	8.6 10	5.3 6
30-34	11.7	10.8	10.3	76	10.5 15	10.8 15
35-39	14.4	14.8	12.3	86	12.3 18	16.3 24
40-44	12.1	15.2	14.7	94	13.7 18	13.4 18
45-49	15.9	16.6	17.9	97	14.7 17	16.0 19
50-54	15.0	16.2	14.1	60	12.7 12	18.8 18
55-59	15.8	15.6	11.8	43	13.8 10	16.4 12
60-64	18.8	15.9	16.3	60	19.6 13	9.3 6
65-69	19.9	17.3	12.0	45	14.1 10	14.4 10
70-74	15.3	16.1	12.0	41	10.2 7	13.4 9
75-79	19.0	18.1	15.7	44	15.4 9	18.4 11
80-84	15.4	16.5	21.7	42	19.2 8	7.2 3
85+	18.5	16.8	12.5	23	17.5 7	14.4 6

16. UTERINE CORPUS, AND UTERUS NOT OTHERWISE SPECIFIED

(Rates are per 100,000 females per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
0-4	0.0	0.0	0.2	1	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.0	0.0	0	0.0	0	0.0	0
15-19	0.0	0.2	0.0	0	0.0	0	0.0	0
20-24	0.1	0.0	0.2	1	0.0	0	0.0	0
25-29	1.1	1.1	1.1	7	0.0	0	0.0	0
30-34	3.0	2.0	1.9	14	3.5	5	0.7	1
35-39	5.7	6.2	5.3	37	4.8	7	2.0	3
40-44	11.1	11.7	14.1	90	16.0	21	14.9	20
45-49	27.2	24.3	29.0	157	23.3	27	26.0	31
50-54	51.4	40.4	51.0	217	67.9	64	53.4	51
55-59	71.6	60.4	72.0	263	103.6	75	84.6	62
60-64	90.6	73.0	90.8	334	114.5	76	90.1	58
65-69	89.6	105.1	101.7	380	84.4	60	100.6	70
70-74	110.5	97.9	111.3	380	128.8	88	113.1	76
75-79	104.5	95.6	110.5	310	94.0	55	100.5	60
80-84	81.2	65.9	98.8	191	86.3	36	134.0	56
85+	72.1	46.2	74.3	137	67.5	27	57.4	24

17. OVARY, ADJACENT STRUCTURES (Rates are per 100,000 females per year)

Age	1980-84	1985-89	1990-94	1990-94	1995		1996	
	Rate/yr.	Rate/yr.	Rate/yr.	Total No.	Rate/Yr.	No.	Rate/Yr.	No.
0-4	0.0	0.0	0.2	1	0.0	0	0.0	0
5-9	0.2	1.1	0.6	3	0.0	0	0.9	1
10-14	0.8	0.6	0.6	3	1.0	1	0.0	0
15-19	1.2	3.1	2.5	12	8.5	8	0.0	0
20-24	2.3	2.9	3.5	20	6.3	6	2.2	2
25-29	3.5	5.1	3.3	22	4.3	5	1.8	2
30-34	4.2	3.5	5.7	42	7.7	11	2.9	4
35-39	6.1	8.0	8.6	60	7.5	11	4.1	6
40-44	13.6	13.6	13.2	84	17.6	23	14.9	20
45-49	23.5	22.0	21.4	116	19.0	22	26.0	31
50-54	28.8	28.8	33.1	141	26.5	25	37.7	36
55-59	35.8	34.8	37.5	137	35.9	26	36.8	27
60-64	51.0	46.9	46.8	172	46.7	31	42.0	27
65-69	52.1	54.8	57.8	216	42.2	30	56.0	39
70-74	52.8	53.2	63.0	215	57.1	39	55.1	37
75-79	47.3	57.8	53.1	149	56.4	33	63.6	38
80-84	71.9	57.4	66.7	129	50.3	21	57.4	24
85+	52.9	38.0	57.0	105	55.0	22	33.5	14

18. **PROSTATE** (Rates are per 100,000 males per year)

Age	1980-84	1985-89	1990-94	1990-94	1995		1996	
	Rate/yr.	Rate/yr.	Rate/yr.	Total No.	Rate/Yr.	No.	Rate/Yr.	No.
0-4	0.2	0.0	0.0	0	0.0	0	0.0	0
5-9	0.0	0.0	0.0	0	0.0	0	0.0	0
10-14	0.0	0.2	0.0	0	0.0	0	0.0	0
15-19	0.0	0.0	0.0	0	0.0	0	0.0	0
20-24	0.1	0.0	0.0	0	0.0	0	0.0	0
25-29	0.0	0.0	0.0	0	0.0	0	0.0	0
30-34	0.2	0.0	0.0	0	0.7	1	0.0	0
35-39	0.0	0.2	0.2	1	0.7	1	0.7	1
40-44	1.4	0.9	2.8	17	0.8	1	4.7	6
45-49	4.4	4.7	14.1	73	25.3	28	28.0	32
50-54	20.0	25.2	64.8	261	108.9	97	123.8	112
55-59	72.3	70.0	187.9	640	277.7	188	314.4	216
60-64	165.2	186.1	441.8	1470	566.5	343	587.6	346
65-69	333.3	390.8	822.1	2504	877.9	519	821.3	484
70-74	538.8	618.7	1126.6	2856	1147.3	592	1015.4	512
75-79	791.1	799.0	1286.4	2358	1046.7	411	898.3	368
80-84	939.0	994.3	1290.2	1295	866.2	200	813.7	194
85+	1062.7	1072.6	1212.7	778	927.4	134	767.4	119

19. **TESTIS** (Rates are per 100,000 males per year)

Age	1980-84	1985-89	1990-94	1990-94	1995		1996	
	Rate/yr.	Rate/yr.	Rate/yr.	Total No.	Rate/Yr.	No.	Rate/Yr.	No.
0-4	0.2	0.9	0.8	5	0.0	0	2.6	3
5-9	0.2	0.0	0.2	1	0.0	0	0.8	1
10-14	0.3	0.0	0.0	0	0.0	0	0.0	0
15-19	3.1	2.6	3.8	19	5.0	5	1.0	1
20-24	10.0	8.4	7.7	45	5.1	5	8.4	8
25-29	9.8	12.6	14.0	92	23.2	27	14.1	16
30-34	12.3	13.3	15.4	113	9.2	13	13.7	19
35-39	9.8	10.6	11.7	80	7.7	11	11.0	16
40-44	7.5	6.3	9.9	60	5.6	7	10.9	14
45-49	5.5	4.2	4.6	24	3.6	4	7.0	8
50-54	2.9	3.2	4.7	19	6.7	6	3.3	3
55-59	2.9	2.8	1.8	6	0.0	0	8.7	6
60-64	2.9	0.3	0.9	3	1.7	1	1.7	1
65-69	2.4	1.6	1.3	4	1.7	1	1.7	1
70-74	1.4	0.4	0.4	1	1.9	1	2.0	1
75-79	0.7	0.6	0.6	1	0.0	0	2.4	1
80-84	0.0	5.9	1.0	1	0.0	0	0.0	0
85+	0.0	0.0	0.0	0	0.0	0	0.0	0

20. **BLADDER** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.2	0.2	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.1	0.2	0.4	2	1.0 1	0.0 0
20-24	1.0	0.7	1.0	6	1.0 1	0.0 0
25-29	0.6	0.8	0.8	5	0.9 1	0.0 0
30-34	2.5	2.2	2.7	20	0.7 1	1.5 2
35-39	4.2	4.5	2.8	19	2.1 3	2.7 4
40-44	9.4	9.9	7.4	45	10.4 13	7.0 9
45-49	19.0	18.4	22.6	117	20.7 23	16.6 19
50-54	37.4	40.4	33.5	135	40.4 36	33.2 30
55-59	58.3	67.0	66.0	225	65.0 44	58.2 40
60-64	96.0	112.2	113.9	379	94.1 57	105.3 62
65-69	158.7	157.2	179.6	547	192.8 114	171.4 101
70-74	208.1	214.6	215.0	545	191.9 99	210.2 106
75-79	312.0	295.3	282.0	517	211.4 83	290.5 119
80-84	353.3	352.3	351.7	353	320.5 74	339.7 81
85+	352.3	406.7	332.0	213	332.2 48	367.6 57
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.9 1
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.3	0.0	0.2	1	0.0 0	0.0 0
20-24	0.4	0.3	0.0	0	0.0 0	0.0 0
25-29	0.0	0.7	0.2	1	0.9 1	0.0 0
30-34	0.8	0.0	0.8	6	0.7 1	0.0 0
35-39	2.1	2.1	1.1	8	2.1 3	1.4 2
40-44	3.9	1.9	3.3	21	2.3 3	1.5 2
45-49	7.6	7.4	5.5	30	2.6 3	8.4 10
50-54	13.1	12.6	13.6	58	14.9 14	13.6 13
55-59	16.6	18.1	23.0	84	18.0 13	16.4 12
60-64	28.7	29.9	31.5	116	37.7 25	42.0 27
65-69	44.1	40.7	43.1	161	52.1 37	50.3 35
70-74	47.7	56.4	58.3	199	41.0 28	50.6 34
75-79	61.3	63.0	72.4	203	66.7 39	77.0 46
80-84	74.9	72.8	82.2	159	100.7 42	83.8 35
85+	82.4	94.6	74.9	138	72.4 29	117.3 49

21. **KIDNEY, RENAL PELVIS, URETER** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	3.5	1.8	2.3	14	2.6 3	2.6 3
5-9	0.8	0.6	0.2	1	0.0 0	0.0 0
10-14	0.0	0.2	0.0	0	0.0 0	0.0 0
15-19	0.1	0.0	0.0	0	0.0 0	0.0 0
20-24	0.3	0.0	0.2	1	0.0 0	0.0 0
25-29	0.3	0.6	0.9	6	0.0 0	0.0 0
30-34	1.3	0.9	1.2	9	0.7 1	4.3 6
35-39	1.6	3.4	3.4	23	5.6 8	2.1 3
40-44	7.1	5.7	5.9	36	5.6 7	10.1 13
45-49	14.9	11.5	10.8	56	15.3 17	11.4 13
50-54	19.5	21.7	19.1	77	21.3 19	25.4 23
55-59	27.0	34.6	31.7	108	48.7 33	34.9 24
60-64	43.3	46.6	48.7	162	57.8 35	62.8 37
65-69	56.7	60.2	76.2	232	98.1 58	83.2 49
70-74	69.6	71.8	74.6	189	93.0 48	65.5 33
75-79	59.5	81.6	80.7	148	101.9 40	92.8 38
80-84	78.1	79.5	78.7	79	69.3 16	92.3 22
85+	60.6	64.4	71.7	46	34.6 5	64.5 10
FEMALES						
0-4	1.9	2.8	1.6	9	0.9 1	5.5 6
5-9	0.4	0.8	0.8	4	0.9 1	1.8 2
10-14	0.0	0.2	0.2	1	0.0 0	0.9 1
15-19	0.0	0.3	0.4	2	0.0 0	0.0 0
20-24	0.0	0.3	0.0	0	0.0 0	0.0 0
25-29	0.3	0.7	0.5	3	1.7 2	0.0 0
30-34	0.6	0.9	1.1	8	1.4 2	0.0 0
35-39	1.4	1.6	1.9	13	3.4 5	3.4 5
40-44	3.7	3.7	3.3	21	5.3 7	4.5 6
45-49	5.7	5.2	4.8	26	6.0 7	3.4 4
50-54	4.4	8.9	10.8	45	8.5 8	11.5 11
55-59	11.2	13.1	17.0	62	16.6 12	10.9 8
60-64	15.3	19.0	23.4	86	21.1 14	17.1 11
65-69	26.7	25.1	28.1	105	35.2 25	35.9 25
70-74	26.1	28.5	43.1	147	30.7 21	40.2 27
75-79	30.1	38.1	37.8	106	29.1 17	41.9 25
80-84	27.9	29.5	48.6	94	55.1 23	59.8 25
85+	27.7	30.8	35.3	65	40.0 16	50.3 21

22. **BRAIN, CEREBRAL MENINGES, AND CRANIAL NERVES** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	2.0	3.7	5.2	31	4.3	5	4.4	5
5-9	2.4	2.9	3.9	22	4.2	5	3.3	4
10-14	1.8	2.0	1.7	9	0.9	1	2.7	3
15-19	1.8	2.7	2.4	12	2.0	2	1.0	1
20-24	2.9	2.6	2.4	14	1.0	1	5.3	5
25-29	2.7	2.9	3.3	22	0.0	0	4.4	5
30-34	2.9	4.3	2.7	20	6.4	9	2.9	4
35-39	5.6	3.8	4.1	28	3.5	5	7.6	11
40-44	5.5	5.2	6.1	37	6.4	8	2.3	3
45-49	9.7	7.3	6.6	34	8.1	9	6.1	7
50-54	11.9	10.6	11.9	48	16.8	15	8.8	8
55-59	16.7	15.6	19.4	66	20.7	14	7.3	5
60-64	23.1	21.5	15.0	50	14.9	9	18.7	11
65-69	22.4	33.5	24.3	74	25.4	15	20.4	12
70-74	24.3	30.1	29.2	74	11.6	6	25.8	13
75-79	27.2	23.2	30.0	55	12.7	5	22.0	9
80-84	24.5	21.1	28.9	29	30.3	7	21.0	5
85+	20.9	15.8	23.4	15	34.6	5	25.8	4
FEMALES								
0-4	3.1	4.1	3.3	19	3.6	4	2.8	3
5-9	2.5	2.8	3.2	17	2.6	3	0.9	1
10-14	2.7	2.5	3.6	18	5.7	6	2.8	3
15-19	2.6	1.8	1.7	8	0.0	0	2.1	2
20-24	2.6	1.9	2.8	16	3.1	3	2.2	2
25-29	3.1	3.0	1.5	10	1.7	2	1.8	2
30-34	0.9	2.8	2.8	21	3.5	5	2.2	3
35-39	2.2	3.7	2.3	16	0.7	1	2.7	4
40-44	3.5	3.4	2.4	15	5.3	7	3.0	4
45-49	4.8	4.8	4.4	24	7.8	9	3.4	4
50-54	8.4	4.8	10.3	44	5.3	5	5.2	5
55-59	9.7	12.2	9.3	34	6.9	5	6.8	5
60-64	13.2	11.3	10.3	38	10.6	7	10.9	7
65-69	20.2	17.3	16.3	61	19.7	14	14.4	10
70-74	17.2	20.2	21.4	73	20.5	14	26.8	18
75-79	15.2	19.2	17.1	48	13.7	8	15.1	9
80-84	20.4	15.6	15.0	29	12.0	5	33.5	14
85+	10.0	11.7	13.0	24	10.0	4	21.5	9

23. **THYROID** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.4	2	0.0 0	0.0 0
10-14	0.7	0.2	1.0	5	0.0 0	0.0 0
15-19	0.4	0.2	0.2	1	0.0 0	0.0 0
20-24	1.1	1.0	0.7	4	2.0 2	0.0 0
25-29	1.0	1.0	1.8	12	3.4 4	6.2 7
30-34	1.8	2.7	2.3	17	0.0 0	0.7 1
35-39	2.9	2.6	2.8	19	4.2 6	2.7 4
40-44	3.1	3.5	3.5	21	4.8 6	4.7 6
45-49	2.1	2.1	5.0	26	4.5 5	9.6 11
50-54	5.1	5.4	6.0	24	6.7 6	7.7 7
55-59	4.6	4.7	5.6	19	3.0 2	5.8 4
60-64	4.3	4.9	7.8	26	5.0 3	10.2 6
65-69	4.3	6.1	6.2	19	6.8 4	8.5 5
70-74	5.3	5.9	4.7	12	5.8 3	6.0 3
75-79	7.5	5.9	9.8	18	2.6 1	14.7 6
80-84	1.4	8.4	5.0	5	8.7 2	0.0 0
85+	7.6	3.7	4.7	3	0.0 0	0.0 0
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.4	0.2	0.0	0	0.0 0	0.0 0
10-14	0.3	0.6	0.6	3	0.0 0	0.9 1
15-19	2.2	2.2	2.3	11	1.1 1	4.2 4
20-24	3.2	3.3	3.7	21	5.2 5	5.5 5
25-29	7.1	5.0	6.7	44	8.6 10	6.2 7
30-34	8.3	5.4	7.8	58	5.6 8	12.9 18
35-39	9.0	7.7	11.8	83	11.6 17	12.9 19
40-44	8.5	8.1	11.1	71	5.3 7	14.9 20
45-49	5.4	8.2	11.3	61	21.6 25	10.9 13
50-54	6.5	9.6	13.4	57	14.9 14	12.6 12
55-59	12.6	6.2	9.6	35	6.9 5	6.8 5
60-64	8.2	9.6	9.2	34	13.6 9	9.3 6
65-69	4.9	7.3	11.0	41	12.7 9	12.9 9
70-74	7.8	6.5	10.8	37	5.9 4	10.4 7
75-79	6.7	7.9	9.3	26	5.1 3	8.4 5
80-84	12.6	8.3	6.7	13	7.2 3	12.0 5
85+	5.0	5.6	9.8	18	10.0 4	12.0 5

24. **HODGKIN'S DISEASE** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.3	2	0.0 0	0.9 1
5-9	0.6	0.8	0.5	3	0.0 0	0.0 0
10-14	1.2	1.6	1.2	6	0.9 1	1.8 2
15-19	5.6	6.7	3.6	18	2.0 2	3.9 4
20-24	8.8	5.9	8.9	52	8.1 8	6.3 6
25-29	5.5	7.3	7.6	50	7.7 9	6.2 7
30-34	4.9	4.1	6.1	45	5.0 7	2.9 4
35-39	6.0	5.8	6.0	41	3.5 5	4.1 6
40-44	3.6	3.3	3.8	23	4.0 5	4.7 6
45-49	3.1	4.5	3.3	17	7.2 8	6.1 7
50-54	3.7	5.6	3.0	12	1.1 1	5.5 5
55-59	5.6	3.9	3.5	12	3.0 2	2.9 2
60-64	5.9	4.1	3.9	13	8.3 5	6.8 4
65-69	4.0	5.8	4.9	15	1.7 1	1.7 1
70-74	7.7	4.7	4.3	11	5.8 3	7.9 4
75-79	9.9	5.7	5.5	10	7.6 3	7.3 3
80-84	4.1	11.0	4.0	4	8.7 2	8.4 2
85+	7.6	9.0	9.4	6	0.0 0	0.0 0
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.4	0.0	0	0.0 0	0.0 0
10-14	2.2	1.0	0.8	4	1.9 2	2.8 3
15-19	5.2	5.8	5.0	24	5.3 5	6.3 6
20-24	6.3	8.2	9.2	52	6.3 6	5.5 5
25-29	6.3	5.7	6.7	44	9.4 11	5.3 6
30-34	3.9	3.6	5.5	41	4.9 7	2.9 4
35-39	2.1	2.8	4.0	28	3.4 5	3.4 5
40-44	2.1	2.6	4.1	26	0.8 1	4.5 6
45-49	0.7	1.8	2.2	12	0.9 1	1.7 2
50-54	1.6	2.3	1.6	7	1.1 1	0.0 0
55-59	2.0	2.5	3.0	11	2.8 2	1.4 1
60-64	1.4	2.4	2.5	9	0.0 0	1.6 1
65-69	2.5	1.8	1.1	4	2.8 2	4.3 3
70-74	7.5	4.1	2.3	8	1.5 1	4.5 3
75-79	4.9	2.8	4.6	13	5.1 3	1.7 1
80-84	5.3	4.7	6.7	13	2.4 1	4.8 2
85+	0.7	1.3	3.8	7	5.0 2	9.6 4

25. **NON-HODGKIN'S LYMPHOMA** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr.	1995 No.	1996 Rate/Yr.	1996 No.
MALES								
0-4	0.8	0.7	0.7	4	0.0	0	0.9	1
5-9	1.7	1.7	2.2	12	0.0	0	0.0	0
10-14	1.6	1.4	2.5	13	2.7	3	0.0	0
15-19	2.3	2.0	2.8	14	4.0	4	1.0	1
20-24	2.5	2.9	3.4	20	4.1	4	3.2	3
25-29	2.3	4.8	8.3	53	8.6	10	7.9	9
30-34	4.4	5.8	10.1	74	7.1	10	8.7	12
35-39	4.5	9.1	10.2	70	12.5	18	9.6	14
40-44	9.7	11.2	14.7	89	20.8	26	18.6	24
45-49	14.5	18.2	22.8	118	19.8	22	16.6	19
50-54	16.5	20.9	27.3	110	37.0	33	25.4	23
55-59	30.3	32.2	32.9	112	35.5	24	36.4	25
60-64	48.3	45.9	50.5	168	51.2	31	47.6	28
65-69	49.9	55.1	66.0	201	74.4	44	76.4	45
70-74	65.7	81.3	83.6	212	89.2	46	97.2	49
75-79	83.8	100.8	113.5	208	122.2	48	97.6	40
80-84	92.4	105.5	116.6	117	160.3	37	159.4	38
85+	62.4	100.8	121.6	78	103.8	15	109.6	17
FEMALES								
0-4	0.4	0.8	0.2	1	0.9	1	0.9	1
5-9	0.4	0.4	0.4	2	0.0	0	0.0	0
10-14	0.2	1.1	0.8	4	2.9	3	0.9	1
15-19	1.3	1.2	1.0	5	1.1	1	1.0	1
20-24	1.3	1.6	2.8	16	0.0	0	1.1	1
25-29	1.6	3.7	3.0	20	3.4	4	0.9	1
30-34	3.0	3.9	3.1	23	4.2	6	5.7	8
35-39	4.9	4.9	7.4	52	10.3	15	6.1	9
40-44	7.4	8.6	8.8	56	10.7	14	3.7	5
45-49	10.8	12.2	12.9	70	13.8	16	17.6	21
50-54	14.9	18.0	19.7	84	11.7	11	23.0	22
55-59	25.8	27.3	24.1	88	13.8	10	32.7	24
60-64	32.5	38.5	36.2	133	37.7	25	32.6	21
65-69	43.5	49.5	53.0	198	42.2	30	54.6	38
70-74	50.4	66.0	61.5	210	90.7	62	78.9	53
75-79	62.6	62.7	84.2	236	75.2	44	90.4	54
80-84	64.0	94.5	84.3	163	98.3	41	100.5	42
85+	56.2	68.2	77.6	143	84.9	34	86.2	36

26. **MULTIPLE MYELOMA** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.0	0.0	0.0	0	0.0 0	0.0 0
25-29	0.0	0.0	0.0	0	0.0 0	0.0 0
30-34	0.0	0.4	0.0	0	0.0 0	0.0 0
35-39	1.1	0.8	0.3	2	1.4 2	2.7 4
40-44	1.8	1.3	1.0	6	1.6 2	0.0 0
45-49	4.1	3.5	3.5	18	1.8 2	2.6 3
50-54	6.6	2.7	5.5	22	4.5 4	6.6 6
55-59	9.7	7.9	12.3	42	16.3 11	11.6 8
60-64	9.6	12.2	17.4	58	19.8 12	11.9 7
65-69	22.3	23.6	23.6	72	30.5 18	17.0 10
70-74	25.5	25.7	29.2	74	25.2 13	39.7 20
75-79	34.6	41.0	30.6	56	25.5 10	41.5 17
80-84	41.5	37.2	46.8	47	56.3 13	62.9 15
85+	56.9	33.9	45.2	29	62.3 9	19.3 3
FEMALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.0	0.0	0.0	0	0.0 0	0.0 0
25-29	0.0	0.0	0.0	0	0.0 0	0.0 0
30-34	0.5	0.0	0.1	1	0.0 0	0.0 0
35-39	0.3	0.3	0.1	1	0.0 0	0.0 0
40-44	1.7	0.7	1.1	7	1.5 2	0.0 0
45-49	0.7	2.9	1.5	8	2.6 3	1.7 2
50-54	5.0	3.8	3.1	13	5.3 5	6.3 6
55-59	7.0	6.3	6.6	24	4.2 3	5.5 4
60-64	10.8	11.1	12.0	44	15.1 10	10.9 7
65-69	16.3	14.2	14.7	55	14.1 10	15.8 11
70-74	20.6	20.3	23.1	79	17.6 12	11.9 8
75-79	21.4	23.5	18.2	51	34.2 20	26.8 16
80-84	29.8	36.2	32.1	62	26.4 11	38.3 16
85+	30.2	35.3	28.8	53	17.5 7	23.9 10

27. **LEUKEMIA** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	6.3	9.0	5.2	31	5.1 6	9.6 11
5-9	3.7	3.5	2.7	15	2.5 3	5.8 7
10-14	2.8	3.2	2.9	15	2.7 3	1.8 2
15-19	2.8	2.8	2.2	11	2.0 2	1.0 1
20-24	1.5	1.7	2.9	17	7.1 7	2.1 2
25-29	2.2	2.0	2.4	16	2.6 3	3.5 4
30-34	3.8	3.9	3.0	22	2.8 4	5.1 7
35-39	3.5	4.1	4.7	32	3.5 5	1.4 2
40-44	3.4	5.7	5.6	34	8.0 10	5.4 7
45-49	5.7	9.0	5.8	30	10.8 12	6.1 7
50-54	14.2	13.7	12.7	51	16.8 15	13.3 12
55-59	21.0	23.3	20.3	69	16.3 11	11.6 8
60-64	35.3	27.6	33.1	110	44.6 27	34.0 20
65-69	46.8	41.1	45.6	139	47.4 28	52.6 31
70-74	71.2	65.7	62.3	158	83.3 43	63.5 32
75-79	100.4	87.1	71.5	131	73.9 29	43.9 18
80-84	132.6	110.6	100.6	101	69.3 16	88.1 21
85+	157.1	124.4	112.2	72	110.7 16	122.5 19
FEMALES						
0-4	6.8	7.0	5.3	30	8.1 9	5.5 6
5-9	3.9	2.7	3.8	20	2.6 3	0.9 1
10-14	1.4	2.5	2.8	14	1.9 2	1.9 2
15-19	2.3	1.0	1.7	8	4.2 4	1.0 1
20-24	1.3	1.3	1.9	11	2.1 2	2.2 2
25-29	2.1	2.8	1.4	9	0.9 1	0.9 1
30-34	3.6	2.1	1.0	7	6.3 9	0.7 1
35-39	3.5	4.4	3.0	21	4.1 6	4.1 6
40-44	4.8	4.0	4.5	29	3.8 5	3.0 4
45-49	6.6	6.5	6.3	34	4.3 5	5.0 6
50-54	7.6	6.3	6.6	28	5.3 5	9.4 9
55-59	12.9	10.6	13.4	49	12.4 9	12.3 9
60-64	17.0	18.5	16.9	62	21.1 14	14.0 9
65-69	25.4	23.8	24.1	90	31.0 22	18.7 13
70-74	34.4	33.3	31.3	107	42.4 29	29.8 20
75-79	48.0	39.8	37.8	106	34.2 20	60.3 36
80-84	56.4	61.2	48.1	93	43.1 18	45.5 19
85+	74.0	66.6	70.5	130	35.0 14	40.7 17

28. **KAPOSI'S SARCOMA** (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.0	0.0	0.0	0	0.0 0	0.0 0
5-9	0.0	0.0	0.0	0	0.0 0	0.0 0
10-14	0.0	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.0	0.0	0	0.0 0	0.0 0
20-24	0.1	1.3	0.9	5	1.0 1	1.1 1
25-29	0.8	3.8	4.2	28	2.6 3	4.4 5
30-34	1.1	6.1	7.1	52	9.9 14	2.9 4
35-39	0.4	5.9	7.4	51	4.9 7	5.5 8
40-44	0.2	3.8	5.4	33	1.6 2	4.7 6
45-49	0.3	3.0	4.8	25	4.5 5	0.0 0
50-54	1.0	2.7	3.0	12	4.5 4	4.4 4
55-59	1.0	2.1	2.1	7	3.0 2	1.5 1
60-64	1.6	1.4	1.5	5	3.3 2	1.7 1
65-69	1.7	2.0	3.3	10	0.0 0	0.0 0
70-74	3.9	3.9	2.4	6	1.9 1	2.0 1
75-79	6.1	3.9	2.7	5	5.1 2	4.9 2
80-84	2.7	5.9	6.0	6	8.7 2	12.6 3
85+	9.4	14.1	17.2	11	6.9 1	19.3 3

Note: There were 25 or fewer cancers in females in each 5-year time period

29. UNKNOWN, ILL-DEFINED SITES (Rates are per 100,000 per year)

Age	1980-84 Rate/yr.	1985-89 Rate/yr.	1990-94 Rate/yr.	1990-94 Total No.	1995 Rate/Yr. No.	1996 Rate/Yr. No.
MALES						
0-4	0.2	0.3	0.3	2	0.0 0	0.9 1
5-9	0.0	0.0	0.0	0	0.0 0	0.8 1
10-14	0.0	0.0	0.2	1	0.0 0	0.0 0
15-19	0.1	0.1	0.0	0	0.0 0	0.0 0
20-24	0.3	0.6	0.3	2	0.0 0	1.1 1
25-29	0.6	0.4	0.6	4	0.9 1	0.0 0
30-34	0.5	1.1	1.9	14	1.4 2	0.7 1
35-39	1.9	1.4	2.0	14	2.1 3	2.1 3
40-44	4.4	3.2	6.1	37	2.4 3	7.0 9
45-49	7.2	9.8	7.3	38	8.1 9	7.9 9
50-54	12.0	15.5	14.9	60	16.8 15	14.4 13
55-59	26.8	26.2	27.0	92	29.5 20	16.0 11
60-64	40.3	41.2	37.9	126	33.0 20	40.8 24
65-69	62.6	54.8	56.5	172	45.7 27	45.8 27
70-74	83.1	84.6	77.7	197	54.3 28	75.4 38
75-79	108.8	98.6	99.3	183	84.0 33	87.9 36
80-84	136.0	158.2	154.4	155	125.6 29	151.0 36
85+	200.5	183.1	180.8	116	193.8 28	167.7 26
FEMALES						
0-4	0.6	0.4	0.7	4	0.0 0	0.9 1
5-9	0.0	0.0	0.0	0	0.0 0	0.9 1
10-14	0.2	0.0	0.0	0	0.0 0	0.0 0
15-19	0.0	0.2	0.0	0	0.0 0	0.0 0
20-24	0.4	0.4	0.0	0	1.0 1	0.0 0
25-29	0.4	0.4	0.6	4	1.7 2	0.9 1
30-34	0.3	1.3	0.8	6	1.4 2	0.0 0
35-39	2.2	2.0	2.0	14	0.7 1	0.7 1
40-44	4.4	3.2	2.7	17	3.8 5	5.2 7
45-49	7.2	5.6	5.4	29	3.5 4	5.9 7
50-54	13.9	12.6	8.2	35	12.7 12	12.6 12
55-59	21.1	16.1	18.9	69	12.4 9	19.1 14
60-64	30.5	27.0	28.8	106	24.1 16	31.1 20
65-69	46.2	36.0	34.5	129	35.2 25	43.1 30
70-74	63.6	65.4	52.7	180	52.7 36	53.6 36
75-79	85.3	72.8	83.1	233	59.8 35	77.0 46
80-84	106.7	107.8	107.5	208	67.1 28	90.9 38
85+	149.4	150.7	160.6	296	132.4 53	102.9 43

APPENDIX

ICD-O-2 CODES

LYMPHOMAS:

Hodgkin's disease: Morphology codes 9650-9667

Non-Hodgkin's lymphoma: Morphology codes 9590-9595, 9670-9715

Others (excludes lymphomas)

ORAL: C 000-148.

ESOPHAGUS: C 150-159.

STOMACH: C 160-169.

COLON: C 180-189.

RECTUM, ANUS: C 199, 209-218.

LIVER: C 220-221.

GALLBLADDER: C 239-249.

PANCREAS: C 250-259.

LARYNX: C 320-329.

TRACHEA, BRONCHUS, LUNG: C 339-349.

CONNECTIVE TISSUE: C 470-479 and 490-499.

MELANOMA OF SKIN: C 440-449, Morphology 8720-8780.

PROSTATE: C 619.

FEMALE BREAST: Sex female and C 500-509.

TESTIS: C 620-629.

CERVIX UTERUS: C 530-539.

UTERINE CORPUS, AND UTERUS NOT OTHERWISE SPECIFIED: C 540-549, 559.

OVARY, ADJACENT STRUCTURES: C 569-578.

BLADDER: C 670-679.

KIDNEY, RENAL PELVIS, URETER: C 649-669.

BRAIN, CEREBRAL MENINGES, CRANIAL NERVES: C 700-719, 723-725.

THYROID: C 739.

MULTIPLE MYELOMA: Morphology 9731-9732.

LEUKEMIA: Morphology 9800-9941.

KAPOSI'S SARCOMA: Morphology 9140.

UNKNOWN SITE: C 760-768, 800-809