

PCMH+ Quality Measure (QM)	PCMH+ Year 3 Absolute Quality Benchmarks ¹			
	50th Percentile	60th Percentile	70th Percentile	80th Percentile
Adolescent well-care visits	72.8%	73.2%	75.2%	76.5%
Avoidance of antibiotic treatment in adults with acute bronchitis ²	31.4%	35.7%	42.3%	44.4%
Developmental screening in the first three years of life	43.4%	51.7%	55.0%	69.4%
Diabetes HbA1c Screening	91.4%	91.9%	92.4%	92.9%
Emergency Department (ED) Usage ³	64.61	58.07	53.70	51.04
Medication management for people with asthma	46.0%	49.5%	51.8%	52.6%
PCMH CAHPS	82.5%	83.8%	84.5%	85.6%
Prenatal Care	75.1%	77.3%	79.9%	80.8%
Postpartum Care	50.3%	56.3%	57.3%	59.9%
Prenatal Care & Postpartum Care ⁴				
Well-child visits in the first 15 months of life	87.1%	88.5%	89.4%	91.0%

Notes:

1. Absolute Quality Benchmarks for 2019 are derived from 2017 comparison group quality measure scores. Participating Entities that meet or exceed the following percentiles will receive the corresponding number of points for that measure: 50th Percentile- .25 point, 60th Percentile- .50 point, 70th Percentile- .75 point, 80th Percentile- 1 point.
2. Avoidance of antibiotic treatment in adults with acute bronchitis measure is calculated as $1 - (\text{adults with bronchitis that received an antibiotic} / \text{adults with bronchitis})$. A higher rate indicates appropriate treatment.
3. Emergency Department (ED) Usage measure is calculated as $(\text{numerator} / \text{denominator}) * 1,000$ member months. A lower score indicates more appropriate care.
4. Benchmark percentiles are not displayed for the Prenatal Care & Postpartum Care combined measure. The individual components will be scored against the individual benchmarks and are each worth half of the points for this measure.
5. Absolute Quality Benchmarks displayed above are rounded, but exact values will be used when awarding points.