



# Save Your Child's Teeth with Sealants!

Protect your child's overall health with dental sealants

## What are dental sealants?

- Dental sealants are thin coatings that fill in the grooves of molars (back teeth) where most tooth decay happens, keeping out food and germs that cause cavities.
- Sealants should be applied to molars usually around ages 6-8.

## How are sealants put on the back teeth?

- Teeth that need sealants are first cleaned with a toothbrush before the sealant material is painted on the tooth; it quickly hardens.
- Placing sealants is painless and takes just a few minutes.

## Can tooth decay occur between sealants?

- Sealants prevent germs from getting the food they need to live, so decay is unlikely to continue, even if some germs are trapped under the sealant.
- Sealants are shown to reduce cavities by 86% the first year and 56% after four years.

## How long will sealants last?

- Sealants can last for several years, and will be checked during dental visits.

## Will sealants make teeth feel different?

- A dental sealant may be slightly noticeable until normal chewing wears it into place.

Ask your dentist about getting dental sealants.



This project is supported by the Centers for Disease Control and Prevention of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award totaling \$3,420,000 with 100 percent funded by CDC/HHS. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by CDC/HHS, or the U.S. Government.



CONNECTICUT  
Public Health

[ct.gov/dph/oralhealth](http://ct.gov/dph/oralhealth)

Scan to download  
this document in  
Spanish, Italian,  
Polish, Chinese, or  
Portuguese.



# Help Prevent Cavities!

Choose healthy food and drinks to keep teeth healthy



## Foods and Drinks Unlikely To Cause Cavities

- Water
- Crunchy Vegetables
- Lettuce
- Cheese
- Almond Butter (no added sugar)
- Eggs
- Meat
- Nuts

## Foods and Drinks Less Likely To Cause Cavities

- Milk
- Berries
- Fresh, Crunchy Fruits
- Oatmeal
- Whole Grains
- Popcorn
- Yogurt
- Ice Cream
- Smoothies
- Dark Chocolate

## Foods and Drinks That May Cause Cavities

- Soda & Juice
- Sports Drinks
- Fruit Snacks & Dried Fruit
- Cereal
- Granola & Cereal Bars
- Chips, Pretzels & Crackers
- Cookies, Candy & Desserts
- Citrus Fruits

### Helpful Tips

- Foods high in carbohydrates, sugars and starches help plaque form and can result in cavities.
- Fruit snacks and dried fruit should be avoided as they are full of sugar and stick to teeth, easily causing cavities.
- Milk and juice (no more than 4 oz per day) should be limited to meal or snack time, and be drunk in one sitting with food. Never put a sugary drink in a “sippy cup”.
- Remember to give your child age-appropriate food. Nuts, hot dogs and grapes are common choking hazards in young children.