## The information below must be entered in the Department of Public Health <u>Medical</u> <u>Exemptions portal</u>. More information is available <u>here</u>.

Name of Primary Care Provider grad	nting exemption:	
Please check one (practitioner gran	ting exemption must	be licensed as one of the following):
$\Box$ Physician (MD or DO) $\Box$	Physician Assistant	□ APRN
License number:	State	e issued:
NPI:		
Phone number:	E	Email:
Directions:		
Part 1. Please complete the demog	raphics section on the	e patient/student.
		at apply to this patient/student (indicate all that
<b>Part 3</b> . If no contraindications or pr patient/student requires the exem		rt 2, write a brief explanation of the reason the
Part 4. Sign the Statement of Clinic	al Opinion and date th	ne form.
Attach a copy of the patient/stude	nt's most current imm	nunization record.
Part 1. Patient/Student Informatio	n:	
First name <u> (in full)</u>	Middle initial	Last name
Date of Birth		
Mailing Address	City	
State	Zip	
Parent/Guardian: First Name		Last name
Primary phone number		
School name		
School address		
City		
State	Zip	
Current or Grade student is enterin	g	

**Part 2.** Please mark the vaccine(s), exemption duration, and all contraindications/precautions that apply to this patient/student for each vaccine.

Medical contraindications and precautions for immunizations are based upon the Advisory Committee on Immunization Practices (ACIP) <u>General Best Practices for Immunization</u>, published by the Centers for Disease Control and Prevention.

A **contraindication** is a condition in a recipient that increases the risk for a serious vaccine adverse event (VAE) or compromises the ability of the vaccine to produce immunity.

A **precaution** is a condition in a recipient that might increase the risk for a serious VAE or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations are deferred when a precaution is self-limiting, but can be administered if the precaution condition improves.

CDC Recognized Contraindications and Precautions		
Vaccine	Exemption Duration	ACIP Contraindications and Precautions (Check all that apply)
<ul> <li>□ Diphtheria- Tetanus-and acellular</li> <li>Pertussis</li> <li>(DTaP)</li> <li>□ Temporary through: /</li> <li>mm/ yyyy</li> <li>□ Permanent</li> </ul>	<ul> <li>Contraindications</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Encephalopathy within seven days after receipt of previous dose of DTP or DTaP</li> <li>Precautions</li> <li>Progressive neurologic disorder, including infantile spasms, uncentralled or logic mean beloage the defen DTaP</li> </ul>	
		<ul> <li>uncontrolled epilepsy, progressive encephalopathy: defer DTaP until neurologic status clarified and stabilized</li> <li>GBS &lt;6 weeks after previous dose of tetanus-toxoid–containing vaccine</li> <li>Fever greater than 40.5°C (104.9°F) &lt;48 hours after vaccination of previous dose of DTP or DTaP</li> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid–containing or tetanus-toxoid–containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid–containing vaccine</li> <li>Moderate or acute illness with or without fever</li> </ul>
Hepatitis A	□ Temporary through: / mm/ yyyy □ Permanent	<ul> <li>Contraindications</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Precautions</li> <li>Moderate or severe acute illness with or without fever</li> </ul>

□ Hepatitis B □ Temporary through: /	Temporary	Contraindications
	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	
	mm/ yyyy	Hypersensitivity to yeast
	Permanent	Precautions
		Moderate or severe acute illness with or without fever
□Haemophilus	Temporary	Contraindications
influenzae type throug b (HiB)	through:	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	□ Age <6 weeks
	Permanent	Precautions
		Moderate or severe acute illness with or without fever
□ Inactivated Influenza Virus (IIV)	□ Temporary through: / mm/ yyyy □ Permanent	<ul> <li>Contraindications</li> <li>Severe allergic reaction (e.g., anaphylaxis) after previous dose of influenza vaccine or to vaccine component</li> <li>Precautions</li> <li>GBS &lt;6 weeks after a previous dose of influenza vaccine</li> </ul>
		<ul> <li>Moderate or severe acute illness with or without fever</li> <li>Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a health care provider who is able to recognize and manage severe allergic conditions).</li> </ul>
<ul> <li>Inactivated</li> <li>Polio Vaccine</li> <li>(IPV)</li> </ul>	<ul> <li>Temporary through:</li> <li>/ mm/ yyyy</li> <li>Permanent</li> </ul>	<ul> <li>Contraindications</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Precautions</li> <li>Pregnancy</li> <li>Moderate or acute illness with or without fever</li> </ul>

Attenuated thro Influenza Virus (LAIV) mn	Temporary through:/	Contraindications	
		Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	
	mm∕ yyyy □ Permanent	<ul> <li>Concomitant use of aspirin or aspirin-containing medication in children and adolescents</li> </ul>	
		<ul> <li>LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days.(e)</li> </ul>	
		Pregnancy	
		<ul> <li>Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months.</li> </ul>	
		Persons with active cerebrospinal fluid/oropharyngeal communications/leaks.	
		Close contacts and caregivers of severely immunosuppressed persons who require a protected environment.	
		<ul> <li>Persons with cochlear implants (due to the potential for CSF leak, which might exist for some period of time after implantation.</li> <li>Providers might consider consultation with a specialist concerning risk of persistent CSF leak if an age-appropriate inactivated or recombinant vaccine cannot be used).</li> </ul>	
		Altered Immunocompetence	
		□ Anatomic or functional asplenia (e.g. sickle cell disease)	
		Precautions	
		□ GBS <6 weeks after a previous dose of influenza vaccine	
		Asthma in persons aged 5 years old or older	
		<ul> <li>Medical conditions which might predispose to higher risk of complications attributable to influenza(d)</li> </ul>	
		Moderate or severe acute illness with or without fever	
☐ Meningo- coccal conjugate vaccines (MenACWY)	□ Temporary through: / mm/ yyyy □ Permanent	Contraindications <ul> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component, including yeast</li> </ul>	
		Precautions	
		Moderate or severe acute illness with or without fever	

Measles-	Temporary	Contraindications	
Mumps-Rubella (MMR)	through: /	Severe allergic reaction (e.g., anaphylaxis) after a previous dos to a vaccine component	
	mm/ yyyy	Pregnancy	
	Permanent	Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised)	
		□ Family history of altered immunocompetence (i)	
		Precautions	
		□ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product)	
		History of thrombocytopenia or thrombocytopenic purpura	
		<ul> <li>Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing <u>(k)</u></li> </ul>	
		Moderate or severe acute illness with or without fever	
Pneumo-	Temporary	Contraindications	
coccal (PCV13)	through:	Severe allergic reaction (e.g., anaphylaxis) after a previous dose of	
	/ mm/ yyyy	PCV13 or any diphtheria-toxoid–containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid–	
	□ Permanent	containing vaccine), including yeast	
		Precautions	
		Moderate or acute illness with or without fever	
🗌 Tdap	Temporary	Contraindications	
	through:	□ Severe allergic reaction (e.g., anaphylaxis) after a previous dose	
	/ mm/ yyyy	or to a vaccine component	
	Permanent	Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or Tdap	
		Precautions	
		□ GBS <6 weeks after a previous dose of tetanus-toxoid–containing vaccine	
		Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized	
		<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid—containing or tetanus-toxoid—</li> </ul>	

		<ul> <li>containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> </ul>	
🗆 Varicella	Temporary	Contraindications	
	through:	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	
	mm/ yyyy	□ Known severe immunodeficiency (e.g., from hematologic and	
Permanent	<ul> <li>solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) (g)</li> <li>Pregnancy</li> </ul>		
		Family history of altered immunocompetence (j)	
		Precautions	
	ſ	Recent (<11 months) receipt of antibody-containing blood product (specific interval depends on product)	
		Moderate or acute illness with or without fever	

## Part 3. Other Allergic Reactions/ Other Type of Medical Condition

Complete this section if claiming a medical exemption for a vaccine based on a condition that does meet any of the ACIP criteria for a contraindication or precaution listed in part 2.

Vaccine(s), list all that apply: \_

For each vaccine listed above, select the allergic or other reaction for which medical exemption is being submitted. Please check off any of the following that apply:

- □ This patient has an autoimmune disorder
- □ This patient has a family history of an autoimmune disorder
- □ This patient has a family history of a reaction to a vaccination
- □ This patient has a genetic predisposition to a reaction to a vaccination as determined through genetic testing
- □ This patient has a previous documented reaction that is correlated to a vaccination
- □ Other condition/reaction not listed above (must specify): \_\_\_\_

Please provide an explanation of the reaction/condition listed above:

Part 4. Statement of Clinical Opinion

In accord with the legal requirements of Public Act 21-6, the vaccine(s) indicated above is/are in my clinical opinion medically contraindicated for this patient/student due to the physical condition as explained above.

Clinician's Signature\_\_\_\_\_

Date \_\_\_\_\_

A person may be placed into quarantine or isolation when there are "reasonable grounds to believe to be infected with, or exposed to, a communicable disease or to be contaminated or exposed to contamination or at reasonable risk of having a communicable disease or being contaminated or passing such communicable disease or contamination to other persons if the commissioner determines that such individual or individuals pose a significant threat to the public health and that quarantine or isolation is necessary and the least restrictive alternative to protect or preserve the public health." <u>Conn.</u> Gen. Stat. § 19a-131b(a).