et 10-20-15



Draft Provisional SIM Quality Council Measure Set 10

| # | Consumer Experience Measure | NQF | ACO |
|---|--|------|-----|
| 1 | PCMH – CAHPS measure | 0005 | |
| # | Care coordination/patient safety | NQF | ACO |
| 2 | Plan all-cause readmission | 1768 | |
| 3 | Asthma admission rate(child) | 0728 | |
| 4 | Emergency Department Usage per 1000 | | |
| 5 | Documentation of current medications in the medical record | 0419 | 39 |
| 6 | Annual monitoring for persistent medications (roll-up) | 2371 | |
| 7 | Adult major depressive disorder (MDD): Coordination of care of patients with specific co-morbid conditions | | |

| # | Prevention Measure | NQF | ACO |
|----|---|-----------------|-----|
| 10 | Breast cancer screening | 2372 | 20 |
| 11 | Cervical cancer screening | 0032 | |
| 12 | Chlamydia screening in women | 0033 | |
| 13 | Colorectal cancer screening | 0034 | 19 |
| 14 | Adolescent female immunizations HPV | 1959 | |
| 15 | Weight assessment and counseling for nutrition and physical activity for children/adolescents | 0024 | |
| 16 | Preventative care and screening: BMI screening and follow up | 0421 | 16 |
| 17 | Developmental screening in the first three years of life | 1448 | |
| 18 | Well-child visits in the first 15 months of life | 1392 | |
| 19 | Well-child visits in the third, fourth, fifth and sixth years of life | 1516 | |
| 20 | Adolescent well-care visits | | |
| 21 | Tobacco use screening and cessation intervention | 0028 | 17 |
| 22 | Prenatal Care & Postpartum care | 1517 | |
| 23 | Frequency of Ongoing Prenatal Care (FPC) | 1391 | |
| 24 | Oral health: Primary Caries Prevention | 1419 | |
| 25 | Screening for clinical depression and follow-up plan | 0418 | 18 |
| 26 | Oral Evaluation, Dental Services (Medicaid only) | 2517 | |
| 27 | Behavioral health screening (pediatric, Medicaid only, custom measure) | | |

- Italicized measures = proposed data source is the Electronic Health Record
- NQF #s with strike out are no longer NQF endorsed
- *There are three measure options including AHRQ/PQI and Anthem or NCQA customizations of the AHRQ/PQI
- Highlighted measures = Quality measures from our list that are also on the provisional measure set being proposed by the Department of Social Services for the Medicaid Quality Improvement & Shared Savings Program (MQISSP)

| # | Acute & Chronic Care Measure | NQF | ACO |
|----|---|------|-----|
| 28 | Medication management for people with asthma | 1799 | |
| 29 | Asthma Medication Ratio | 1800 | |
| 30 | DM: Hemoglobin A1c Poor Control (>9%) | 0059 | 27 |
| 31 | DM: HbA1c Screening (possible interim measure until NQF 0059 is | 0057 | |
| | available) | | |
| 32 | DM: Diabetes eye exam | 0055 | 41 |
| 33 | DM: Diabetes foot exam | 0056 | |
| 34 | DM: Diabetes: medical attention for nephropathy | 0062 | |
| 35 | HTN: Controlling high blood pressure | 0018 | 28 |
| 36 | Use of imaging studies for low back pain | 0052 | |
| 37 | Avoidance of antibiotic treatment in adults with acute bronchitis | 0058 | |
| 38 | Appr. treatment for children with upper respiratory infection | 0069 | |

| # | Behavioral Health Measure | NQF | ACO |
|----|---|------|-----|
| 40 | Follow-up care for children prescribed ADHD medication | 0108 | |
| 41 | Metabolic Monitoring for Children and Adolescents on | | |
| | Antipsychotics (pediatric, Medicaid only, custom measure) | | |
| 42 | Depression Remission at 12 Twelve Months | 0710 | 40 |
| 43 | Child and Adolescent Major Depressive Disorder (MDD): Suicide | 1365 | |
| | Risk Assessment | | |
| 44 | Unhealthy Alcohol Use – Screening | | |

| Recommended for Reporting | NQF | ACO |
|--|------|-----|
| Anti-Depressant Medication Management | 0105 | |
| Initiation and Engagement of Alcohol and Other Drug Dependence Treatment | 0004 | |
| Follow up after hospitalization for mental illness, 7 & 30 days | | |
| 30 day readmission (MMDLN) | | |
| ED Use (observed to expected) – New NCQA | | |
| % PCPs that meet Meaningful Use | | 11 |
| Cardiac strss img: Testing in asymptomatic low risk patients | 0672 | |

| Measures for consideration and development | NQF | ACO |
|---|------|-----|
| Gap in HIV medical visits | 2080 | |
| HIV/AIDS: Screening for Chlamydia, Gonorrhea, and Syphilis | 0409 | |
| HIV viral load suppression | 2082 | |
| Annual % asthma patients (2-20) with 1 or more asthma-related ED visits | | |
| ASC admissions: chronic obstructive pulmonary disease (COPD) or asthma in older adults | 0275 | 9 |
| ASC: heart failure (HF) | 0277 | 10 |
| All-cause unplanned admission for MCC | | 38 |
| All-cause unplanned admissions for patients with heart failure | | 37 |
| Preventable hospitalization composite (NCQA)/Ambulatory Care Sensitive Condition composite (AHRQ) | | |
| All-cause unplanned admissions for patients with DM | | 36 |
| Asthma in younger adults admission rate | 0283 | |

Measures Not Currently Recommended for Payment

| | | | _ |
|----|------------------|----|---|
| D | $D \Lambda$ | | П |
| ーレ | \mathbf{n}_{F} | ۱Г | |

| Consumer Experience Measure | NQF | ACO |
|---|-----------------|-----|
| ACO – CAHPS | | 1-7 |
| Preventative Health | NQF | ACO |
| Lead screening in children | | |
| Glaucoma screening | | |
| Childhood immunizations DTaP and IPV | | |
| Childhood immunizations MMR | 0038 | |
| Childhood immunizations VZV | 0038 | |
| Childhood immunizations status combo 2,3,4 | 0038 | |
| Adolescent immunizations TDaP/TD & | 0038 | |
| meningococcal | | |
| Adult BMI assessment | | |
| Influenza immunization | 0041 | 14 |
| Pneumonia vaccination status, older adults** | 0043 | 15 |
| Osteoporosis mngt in women who had a fracture* | * 0053 | |
| Medication Adherence, 81 years & older** | | |
| Topical Fluoride for Children at Elevated Caries Risl | k, 2528 | |
| Dental Services | | |
| Preventive care and screening: screening for high | | 21 |
| blood pressure and follow-up documented | | |
| Pediatric behavioral health screening | 0722 | |
| Maternal depression screening | 1401 | |
| Annual Dental Visit (ADV) | 1388 | |
| Annual Dental Visit (Medicaid, Chip) | | |

| Acute & Chronic Care | NQF | ACO |
|---|-----------------|-----|
| Appr. testing for children with pharyngitis | 0002 | |
| Episiotomy in vaginal deliveries | 0470 | |
| Use of appr. medication for people with asthma | 0036 | |
| Use of spirometry testing in the assessment and diagnosis of COPD | 0577 | |
| Ischemic vascular disease: use of aspirin or another antithrombotic | 0068 | 30 |
| CHF: beta-blocker therapy for left ventricular systolic dysfunction | 0083 | 31 |
| CAD: Persistence of Beta blocker therapy after a heart attack* | 0071 | |
| CAD: Medication adherence | 0543 | |
| Disease mdifying anti-rheumatoid arthritis* | 0054 | |

| Behavioral Health Measure | NQF | ACO |
|--|------|-----|
| Bipolar & Major Depression: Appraisal for Alcohol or Chemical Substance Use | | |
| Depression Utilization of the PHQ-9 Tool | 0712 | |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia | 1879 | |
| Tobacco Use & Help w/ Quitting-Adolescents | | |
| % of adults given a new psychiatric diagnoses, and medication, by a PCP who received a F/U within 30 days (DSS custom) | | |

| **Under | consideration | for | older | adults |
|---------|---------------|-----|-------|--------|

^{*} Recommended for consideration for specialists

| | NOT | 1.00 |
|---|-----------------|------|
| At-Risk Population | NQF | ACO |
| HTN: Serum creatinine testing | 0605 | |
| HTN: Diuretics Persistent use w lab mnitoring | | |
| HTN: Medication adherence | | |
| Ischemic stroke: Warfarin: PT/INR monitoring | 0612 | |
| Ischic vascular dis: complete lipid panel & LDL cntr | l <u>0073</u> | 29 |
| Ischemic vascular disease: MI, CABG, PCI or IVD with complete lipid profile | | |
| Ischemic vascular disease: LDL>=100 and treated with lipid lowering agent, or LDL<100 | 0075 | |
| Ischemic vascular disease: LDL-C screening for patients with cardiovascular conditions | | |
| Heart failure: beta-blocker therapy | 0615 | |
| Digoxin: Persistent use with lab monitoring | 2371 | |
| COPD: Pharmacotherapy mgmt of COPD excerbtio | n 0549 | |
| CAD: all-or-nothing composite: Lipid Control | 0074 | 32 |
| CAD: Angiotensin-converting enzyme inhibitor or angiotensin receptor blocker (ACE-I/ARB) therapy—diabetes or left ventricular systolic dysfunction | 0066 | 33 |
| CAD: Antiplatelet therapy | 2379 | 43 |
| CAD: Symptom management – CAD pt seen within 12 months for angina symptoms | 1 | 44 |
| CAD: Beta blocker therapy prior MI or LVEF | 0070 | 45 |
| CAD: Lipid lowering drug for LDL>30 | | |
| CAD: Post-MI use of ACE-I/ARB w/ h/o, HTN, DM, or HF | 0594 | |
| CAD: Drug-eluting stent: antiplatelet therapy | 2379 | |
| Cardiac strss img: Preoperative eval in low risk surgery patients | 0670 | |
| Care coordination/patient safety | NQF | ACO |
| Falls: screening for future fall risk** | 0101 | 13 |
| Use of high-risk meds in the elderly** | 0022 | |
| % of prescribers that use e-prescribing | - | |
| Adverse event rate – OP procedures | | |
| Risk standardized all condition readmission | 1789 | 8 |
| % adlts w/ inptient "medicine" admissions w/ | | |
| post-admission f/u w/i 7 days | | |
| Potentially avoidable ER rate | | |
| Skilled Nursing Facility 30-day All-Cause Readmission Measure (SNFRM) | 2510 | 35 |
| Pediatric ASC composite admissions | | |
| | | |

| At-Risk Population | NQF | ACO |
|---|------|-----|
| Migraine: Frequent use of acute meds/ receiving prophylactic meds | 0602 | |
| Otitis Media with Effusion: Tympanostomy tube insertion: pediatric hearing test | 0587 | |
| Potentially harmful drug-disease interactions | | |
| DM: High blood pressure control | 0061 | |
| DM: Hemoglobin A1C control (<8%) | 0575 | |
| DM: Tobacco non-use | | |
| DM: Low density lipoprotein (LDL-C) control | 0064 | |
| DM: Low density lipoprotein (LDL-C) screening | 0063 | |
| Diabetes treatment medication for HTN | 0546 | |
| Diabetes: A1C testing | 0057 | |
| Diabetes: A1C testing pediatric | 0061 | |
| DM kidney disease monitoring/urine protein scrning | | |
| Comprehensive diabetes care | | |
| Diabetes: annual lipid profile | | |
| Diabetes: blood pressure management (<140/80) | | |
| Diab: LDL=>100 & lipid lowering agent use | | |
| % OF Metformin use for diabetic and pre-diabetic | | |
| members (UCLA study) | | |
| Diabetes: Medication adherence | 2468 | |
| Epilepsy: Anticonvulsants: Persistent use with lab monitoring | | |
| Epilepsy: Carbamazepine: Persistent use with lab monitoring | | |
| Epilepsy: Phenobarbitol: Persistent use w/ lab monitoring | | |
| Epilepsy: Phenytoin: Persistent use with lab monitoring | | |
| Epilepsy: Valproic acid: Persistent use with lab monitoring | | |
| Cesarean section rate in singleton low risk deliveries | 0471 | |
| Colorectal cancer: follow-up after treatment | | |
| CAD: ACE-I/ARB: Persistent use with lab monitoring | 2371 | |
| Access and Availability of Care | IQF | ACO |
| Children and adolescents access to PCPs | | |
| Adults access to preventive/ambulatory | | |

| Access and Availability of Care | IVQI | ACC |
|--|------|-----|
| Children and adolescents access to PCPs | | |
| Adults access to preventive/ambulatory health services | | |
| | | |

| Resource Efficiency | |
|---------------------------------|-----------------------------|
| Bed days/1000 | ER repeat visit rate |
| Generic dispensing/substitution | % of members with 3 or more |
| | ER visits |
| ER: ambulatory visits/1k | EOC efficiency index |
| Non-emergent ER visits/1000 | Amb Surgery Steerage |
| % of ER visits Non-emergent | Radiology Steerage |

| Obstetrics Measure | NQF | ACO |
|---------------------------|------|-----|
| Elective Delivery* | 0469 | |